

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

WO 03/053912

PCT/US02/40605

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
3 July 2003 (03.07.2003)

PCT


(10) International Publication Number  
WO 03/053912 A1(51) International Patent Classification:  
237/14, A61K 31/18, A61P 25/28(US), MACOR, John, E.; 644 Moose Hill Road, Guilford,  
CT 06437 (US).(21) International Application Number: PCT/US02/40605  
(22) International Filing Date:  
20 December 2002 (20.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 20 December 2001 (20.12.2001) US  
60344,332(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY  
(US/US); P. O. Box 4000, Route 206 and Provincetown  
Road, Princeton, NJ 08543-0000 (US).(72) Inventors: PARKER, Michael, F.; 45 Brimfield Hill Road,  
Hillsgrove, CT 06441 (US); MCELHONNE, Katherine,  
E.; 23 Midway Drive, Cromwell, CT 06416 (US); MATE,  
Robert, A.; 380-91 Hitchcock Road, Waterbury, CT 06705  
(US); BRONSON, Jeanne, J.; 230 Parnelle Hill Road,  
Durham, CT 06422 (US); GAI, Yonghua; 12 Linna Lane,  
Killingworth, CT 06419 (US); BEGGSTROM, Carl, P.;  
22 Paper Mill Drive, Madison, CT 06443 (US); MARCIN,  
Lawrence, R.; 40 Ridgewood Drive, Bethany, CT 06437(84) Designated States (regionally): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Burmian patent (AM, AZ, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EG, ES, FI, FR, GB, GR, HU,  
IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).Published:  
with international search report(81) Designated States (nationally): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EG, ES, FI, FR, GB, GR, HU,  
IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).(74) Agents: BAJAJKO, Suzanne et al.; Bristol-My-  
ers Squibb Company, P.O. Box 4000, Princeton, NJ  
08543-0000 (US).

WO 03/053912 A1

(1) Abstract: There is provided a series of novel  $\alpha$ -(N-sulfonamido)acetamide compounds of the Formula (I) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are defined herein, which are inhibitors of  $\beta$ -amyloid peptide ( $\beta$ -A $\beta$ ) production and are useful in the treatment of Alzheimer's Disease and other conditions affected by anti-amyloid activity.(54) Title:  $\alpha$ -(N-SULFONAMIDO)ACETAMIDE DERIVATIVES AS  $\beta$ -AMYLOID INHIBITORS5  $\alpha$ -(N-SULFONAMIDO)ACETAMIDE DERIVATIVESAS  $\beta$ -AMYLOID INHIBITORSFIELD OF THE INVENTION

10 This invention provides novel  $\alpha$ -(N-sulfonamido)acetamide compounds having drug and bio-affecting properties, their pharmaceutical compositions and method of use. In particular, the invention is concerned with  $\alpha$ -(N-arylsulfonamido)acetamides. These compounds possess unique inhibition of the  $\beta$ -amyloid peptide ( $\beta$ -A $\beta$ ) production, thereby acting to prevent the accumulation of amyloid protein deposits in the brain. More particularly, the present invention relates to the treatment of Alzheimer's Disease (AD).

BACKGROUND OF THE INVENTION

20 Alzheimer's Disease is a progressive, neurodegenerative disorder characterized by memory impairment and cognitive dysfunction. AD is characterized pathologically by the accumulation of senile (neuritic) plaques, neurofibrillary tangles, amyloid deposition in neural tissues and vessels, synaptic loss, and neuronal death. It is the most common form of dementia and it now represents the third leading cause of death after cardiovascular disorders and cancer. The cost of Alzheimer's Disease is enormous (in the U.S., greater than \$100 billion annually) and includes the suffering of the patients, the suffering of families, and the lost productivity of patients and caregivers. As the longevity of society increases, the occurrence of AD will markedly increase. It is estimated that more than 10 million Americans will suffer from AD by the year 2020, if methods for prevention and treatment are not found. Currently, AD is estimated to afflict 10% of the population over age 65 and up to 50% of those over the age

- 2 -

of 85. No treatment that effectively prevents AD or reverses the clinical symptoms and underlying pathophysiology is currently available (for review see Selkoe, D.J. *Annu. Rev. Cell Biol.*, 1994, 10: 373-403).

There have been many theories relating to the etiology and pathogenesis of AD. These theories were either based on analogies with other diseases and conditions (e.g., slow virus and aluminum theories), or based on pathologic observations (e.g., cholinergic, amyloid, or tangle theories). Genetic analysis can potentially differentiate between competing theories. The identification of mutations in the  $\beta$ -amyloid precursor protein ( $\beta$ -APP) of individuals prone to early onset forms of AD and related disorders strongly supports the amyloidogenic theories.

Histopathological examination of brain tissue derived upon autopsy or from neurosurgical specimens in affected individuals reveals the occurrence of amyloid plaques and neurofibrillar tangles in the cerebral cortex of such patients. Similar alterations are observed in patients with Trisomy 21 (Down's syndrome). Biochemical and immunological studies reveal that the dominant proteinaceous component of the amyloid plaque is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids. This protein is designated  $A\beta$ ,  $\beta$ -amyloid peptide, and sometimes  $\beta/A4$ ; referred to herein as  $A\beta$ . In addition to its deposition in amyloid plaques,  $A\beta$  is also found in the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. Compelling evidence accumulated during the last decade reveals that  $A\beta$  is an internal polypeptide derived from a type 1 integral membrane protein, termed  $\beta$ -amyloid precursor protein (APP) (Selkoe, D. *Physiol. Rev.* 2001, 81, 741-766; Wolfe, M. *J. Med. Chem.* 2001, 44, 2039-2060).  $\beta$ -APP is normally produced by many cells both in vivo and in cultured cells, derived from various animals and humans. Several proteolytic fragments of APP are generated by proteinases referred to as secretases. A subset of these proteolytic fragments, designated  $\beta$ -amyloid peptide ( $A\beta$ ), contains 39 to 43 amino acids and is generated by the combined action of  $\beta$ -secretase and  $\gamma$ -secretase.  $\beta$ -secretase is a membrane-bound, aspartyl protease

- 3 -

that forms the N-terminus of the  $A\beta$  peptide. The C-terminus of the  $A\beta$  peptide is formed by  $\gamma$ -secretase, an apparently oligomeric complex that includes presenilin-1 and/or presenilin-2. Presenilin-1 and presenilin-2 are polytopic, membrane-spanning proteins that may contain the catalytic components of  $\gamma$ -secretase (Selfert, D.; Bradley, J. et al. *J. Biol. Chem.* 2000, 275, 34086-34091).

Multiple lines of evidence together strongly suggest that a reduction in brain  $A\beta$  levels will prevent the onset and progression of AD. First,  $A\beta$  is a major constituent of the parenchymal plaques observed in all AD patients and the cerebral vasculature amyloid deposits observed in 90% AD patients (reviewed in Selkoe, D. *Physiol. Rev.* 2001, 81, 741-766; Wolfe, M. *J. Med. Chem.* 2001, 44, 2039-2060). These plaques are formed from the aggregation of soluble  $A\beta$  whose brain levels are highly correlated with the severity of AD neurodegeneration (McLean, C.; Cherry, R. et al. *Annu. Neurol.* 1999, 46, 860-866). Second, mutations in three genes (APP, PS-1, or PS-2) that increase  $A\beta$  cause familial AD (FAD), where AD onset is accelerated by at least a decade. Included in the mutations that increase  $A\beta$  are chromosome 21 Trisomy that causes Down's syndrome. Third, transgenic mice that express one or more of the mutant FAD genes have increased  $A\beta$  levels, form parenchymal plaques and cerebral vascular deposits containing  $A\beta$ , exhibit memory deficits (Chapman, P.; White, G. et al. *Nature Neurosci.* 1999, 2, 271-276) and enhance neurofibrillary degeneration in mice that also overexpress mutant tau (Lewis, J.; Dickson, D. et al. *Science* 2001, 293, 1487-1491). Fourth,  $A\beta$  is toxic to cultured cells (Dahlgren, K.; Manelli, A. et al. *J. Biol. Chem.* 2002 277, 32046-32053), induces neurofibrillary tangles in mice with mutant tau (Gotz, J., Chen, F. et al. *Science* 2001, 293, 1491-1495) and interferes with long-term potentiation, a likely component of memory (Walsh, D., Klyubin, I. et al. *Nature* 2002, 416, 535-539 and references therein). Taken together, these data lead one skilled in the art to conclude that excess  $A\beta$  production and/or reduced  $A\beta$  clearance cause AD. From this it follows that reducing brain  $A\beta$  levels by inhibition of  $\gamma$ -secretase will prevent the onset and progression of AD.

- 4 -

In addition to AD, excess production and/or reduced clearance of A $\beta$  causes cerebral amyloid angiopathy (CAA) (reviewed in Thal, D., Ghebremedhin, E. et al. *J. Neuropath. Exp. Neuro.* 2002, 61, 282-293). In these patients, vascular amyloid deposits cause degeneration of vessel walls and aneurysms that may be responsible for 10-15% hemorrhagic strokes in elderly patients. As in AD, mutations in the gene encoding A $\beta$  lead to an early onset form of CAA, referred to as cerebral hemorrhage with amyloidosis of the Dutch type, and mice expressing this mutant protein develop CAA that is similar to patients.

It is hypothesized that inhibiting the production of A $\beta$  will prevent and reduce neurological degeneration, reducing neurotoxicity and, generally, mediating the pathology associated with A $\beta$  production. Methods of treatment could target the formation of A $\beta$  through the enzymes involved in the proteolytic processing of  $\beta$ -amyloid precursor protein. Compounds that inhibit  $\beta$ - or  $\gamma$ -secretase activity, either directly or indirectly, could control the production of A $\beta$ . Advantageously, compounds that specifically target  $\gamma$ -secretases, could control the production of A $\beta$ . Such inhibition of  $\beta$ - or  $\gamma$ -secretases could thereby reduce production of A $\beta$  which, could reduce or prevent the neurological disorders associated with A $\beta$  protein.

Smith, et al. in International Application WO 00/50391, published August 31, 2000, disclose a series of sulfonamide compounds that can act to modulate production of amyloid  $\beta$  protein as a means of treating a variety of diseases, especially Alzheimer's Disease and other diseases relating to the deposition of amyloid. Japanese Patent No. 11343279, published December 14, 1999 discloses a series of sulfonamide derivatives which are TNF- $\alpha$  inhibitors useful for treating autoimmune diseases.

Nothing in these references can be construed to disclose or suggest the novel compounds of this invention and their use to inhibit  $\beta$ -AP production.


- 5 -

## SUMMARY OF THE INVENTION

A series of  $\alpha$ -(N-sulfonamido)acetamide derivatives have been synthesized. These compounds specifically inhibit the production of  $\beta$ -amyloid peptide ( $\beta$ -AP) from  $\beta$ -amyloid precursor protein ( $\beta$ -APP). The pharmacologic action of these compounds makes them useful for treating conditions responsive to the inhibition of  $\beta$ -AP in a patient, e.g., Alzheimer's Disease (AD) and Down's Syndrome. Therapy utilizing administration of these compounds to patients suffering from, or susceptible to, these conditions involves reducing  $\beta$ -AP available for accumulation and deposition in brains of these patients.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention comprises compounds of Formula I, their pharmaceutical formulations, and their use in inhibiting  $\beta$ -AP production in patients suffering from or susceptible to AD or other disorders resulting from  $\beta$ -AP accumulation in brain tissue. The compounds of Formula I which include nontoxic pharmaceutically acceptable salts and/or hydrates thereof have the following formula and meanings:




20 wherein:

- R<sup>1</sup> is selected from the group consisting of
- (a) a straight or branched-chain C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl optionally substituted with substituents selected from the group consisting of hydroxy, C<sub>1-3</sub> cycloalkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkylthio, and halogen;
- (b) C<sub>2-7</sub> cycloalkyl optionally substituted with hydroxy or halogen;
- R is hydrogen or R<sup>1</sup> and R taken together is C<sub>2-6</sub> alkylene;
- R<sup>2</sup> is selected from the group consisting of

- 6 -

- (a) a straight or branched-chain  $C_{1-4}$ alkyl or  $C_{2-4}$ alkenyl optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-4}$ alkoxy, and  $NR^1R^2$ ;
- (b)  $C_{2-7}$ cycloalkyl/methyl optionally substituted with substituents selected from the group consisting of amino,  $(C_{1-4}$ alkyl)NH-,  $di(C_{1-4}$ alkyl)N-,  $C_{1-4}$ alkylC(=O)NH-, and  $C_{1-4}$ alkylOC(=O)NH-;
- (c) a straight or branched-chain  $C_{1-4}$ alkyl-C(=O)-A;
- (d) -B-naphthyl;
- (e)



D and E are each independently a direct bond, a straight or branched-chain  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl, or  $C_{2-7}$ cycloalkyl;

- Z is selected from the group consisting of hydrogen,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halogen, cyano, hydroxy, -OCHF<sub>3</sub>, -OCF<sub>3</sub>, -CF<sub>3</sub>, and -CHF<sub>3</sub>;

- X and Y are each independently selected from the group consisting of hydrogen, hydroxy, halogen, (halogen)<sub>2</sub>CH-,  $C_{1-4}$ alkylS-,  $C_{1-4}$ alkylS(O)-,  $C_{1-4}$ alkylSO<sub>2</sub>-, nitro, F, S<sub>2</sub>, and cyano; -OR<sup>3</sup>; -NR<sup>4</sup>R<sup>5</sup>; -NR<sup>3</sup>C(=O)R<sup>4</sup>; -NR<sup>3</sup>C(=O)OR<sup>5</sup>; -NHSO<sub>2</sub>C<sub>1-4</sub>alkyl; -N(SO<sub>2</sub>C<sub>1-4</sub>alkyl)<sub>2</sub>; -C(=O)W wherein W is selected from the group consisting of hydroxy,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, phenoxy, and -NR<sup>4</sup>R<sup>5</sup>; -OC(=O)C<sub>1-4</sub>alkyl; -phenyl in which said phenyl is optionally substituted with cyano, halogen,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkylS-, CH<sub>2</sub>C(=O),  $C_{1-4}$ alkylS(O)-, or  $C_{1-4}$ alkylSO<sub>2</sub>-; and heterocyclic group, in which said heterocyclic group is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, oxazolyl, isoxazolyl, thiadiazolyl, and thiazolyl, wherein said heterocyclic group is optionally substituted

- 7 -

- with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}$ alkyl, (halogen) $C_{1-4}$ alkyl, and CO<sub>2</sub>C<sub>1-4</sub>alkyl;
- (f) -B-(heterocycle), in which said heterocycle is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, oxazolyl, isoxazolyl, thiadiazolyl and thiazolyl wherein said heterocycle is optionally substituted with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}$ alkyl, CO<sub>2</sub>C<sub>1-4</sub>alkyl, amino,  $(C_{1-4}$ alkyl)NH-,  $di(C_{1-4}$ alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-( $C_{1-4}$ alkyl)piperazin-1-yl;

(g) -B-(piperidin-4-yl), in which said piperidin-4-yl is optionally

- substituted with substituents selected from the group consisting of a straight or branched-chain  $C_{1-4}$ alkyl, CH<sub>2</sub>C(=O)phenyl, phenyl and phenyl/methyl in which said  $C_{1-4}$ alkyl and said phenyl are optionally substituted with substituents selected from the group consisting of cyano, halogen, benzimidazol-2-yl, pyridyl and tetrahydrofuran-2-yl; and -C(=O)W' wherein W' is selected from the group consisting of  $C_{1-4}$ alkoxy, R<sup>6</sup>, and -NR<sup>4</sup>R<sup>5</sup>;

- 20 A is hydroxy,  $C_{1-4}$ alkoxy or NR<sup>4</sup>R<sup>5</sup>;

B is a straight or branched-chain  $C_{1-4}$ alkyl or  $C_{2-4}$ alkenyl;

R<sup>3</sup> is phenyl or pyridyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkyl, (halogen)<sub>2</sub>CH-, and halogenCH<sub>2</sub>-;

- 25 R<sup>4</sup> and R<sup>5</sup> each are independently hydrogen, a straight or branched-chain  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-7}$ cycloalkyl,  $C_{2-7}$ cycloalkylmethyl,  $C_{1-4}$ alkoxy, phenyl, benzyl, pyridyl, piperidin-4-yl, indan-1-yl, indan-2-yl, tetrahydrofuran-3-yl, or pyrrolidin-3-yl; in which each is optionally substituted with substituents selected from the group consisting of hydroxy, cyano, halogen, (halogen) $C_{1-4}$ alkyl, halogenCH<sub>2</sub>-, hydroxymethyl, benzyloxyethyl, phenyl, pyridyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,



- (halogen)<sub>2</sub>C-O-, (halogen)<sub>2</sub>CH-O-, C<sub>1-4</sub>alkylthio, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl, 4-phenylpiperazin-1-yl, 4-benzylpiperazin-1-yl, 4-pyridylpiperazin-1-yl, CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1-4</sub>alkyl, C(=O)NHC<sub>1-4</sub>alkyl, and C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

- 5 R' and R' taken together may be morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, 1,2,3,4-tetrahydroisoquinolin-2-yl, decalhydroquinolin-1-yl, piperidin-1-yl, piperazin-1-yl, [1,4]-oxazepan-4-yl, azetidin-1-yl, 2,3-dihydro-1*H*-isindol-2-yl, or 2,3-dihydro-1*H*-indol-1-yl; in which each is optionally substituted with substituents selected from the group consisting of hydroxy, cyano, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-, phenyl, pyridyl, benzyl, C<sub>1-4</sub>alkyl, C<sub>2-7</sub>cycloalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylthio, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1-4</sub>alkyl, C(=O)NHC<sub>1-4</sub>alkyl, and C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

- 10 is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-8</sub> alkenyl, benzyl, or phenyl in which each is optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, C<sub>2-7</sub>cycloalkyl, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, (C<sub>1-4</sub>alkyl)(phenyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-

- 20 (C<sub>1-4</sub>alkyl)piperazin-1-yl;

R<sup>2</sup> is hydrogen, a straight or branched-chain C<sub>1-8</sub> alkyl;

- 25 R<sup>3</sup> is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-7</sub>cycloalkyl, phenyl, pyridyl, or furanyl; in which each is optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl;

- 30 R<sup>4</sup> is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-8</sub> alkenyl, benzyl, phenyl, oxazolyl or pyridyl; in which each is optionally substituted with substituents selected from the group consisting of halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, amino, (C<sub>1-4</sub>alkyl)NH-,

di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl; or a non-toxic pharmaceutically acceptable salt thereof.

- 5 The present invention also provides a method for the treatment or alleviation of disorders associated with  $\beta$ -amyloid peptide, especially Alzheimer's Disease, which comprises administering together with a conventional adjuvant, carrier or diluent a therapeutically effective amount of a compound of formula I or a nontoxic pharmaceutically acceptable salt, solvate or hydrate thereof.

- 10 The term "C<sub>1-8</sub> alkyl" as used herein and in the claims (unless the context indicates otherwise) means straight or branched chain alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, 3-methylbutyl, hexyl and the like. The term "C<sub>2-8</sub> alkenyl" used herein and in the claims (unless the context indicates otherwise) means straight or branched chain alkenyl groups such as ethenyl (i.e. vinyl), propenyl, allyl, butenyl, 3-methylbutenyl, pentenyl, hexenyl and the like. Unless otherwise specified, the term "halogen" as used herein and in the claims is intended to include bromine, chlorine, iodine and fluorine while the term "halide" is intended to include bromide, chloride and iodide anion.

- 20 The term "C<sub>2-7</sub> cycloalkyl" means a carbon cyclic ring system such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

The term "C<sub>1-4</sub> haloalkyl" means a straight or branched chain C<sub>1-4</sub> alkyl group containing from 1 to 3 halogen atoms such as trifluoromethyl, fluoroethyl, 1,2-dichloroethyl, trichloroethyl and the like.

- 25 The term "C<sub>2-8</sub> alkylene" means a straight or branched chain alkylene group such as methylene, ethylene, propylene, methylethylene, butylene, methylpropylene, pentylene, methylbutylene and ethylpropylene.

As the compounds of the present invention possess an asymmetric carbon atom, the present invention includes the racemate as well as the individual enantiometric forms of the compounds of Formula I as described herein and in the claims. The use of a single designation such as (R) or (S) is intended to

include mostly one stereoisomer. Mixtures of isomers can be separated into individual isomers according to methods which are known per se, e.g. fractional crystallization, adsorption chromatography or other suitable separation processes. Resulting racemates can be separated into antipodes in the usual manner after introduction of suitable salt-forming groupings, e.g. by forming a mixture of diastereoisomeric salts with optically active salt-forming agents, separating the mixture into diastereomeric salts and converting the separated salts into the free compounds. The possible enantiomeric forms may also be separated by fractionation through chiral high pressure liquid chromatography columns.

The term "nontoxic pharmaceutically acceptable salt" as used herein and in the claims is intended to include nontoxic base addition salts. Suitable salts include those derived from organic and inorganic acids such as, without limitation, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulfonic acid, acetic acid, tartaric acid, lactic acid, sulfonic acid, citric acid, maleic acid, fumaric acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, and the like.


In the method of the present invention, the term "therapeutically effective amount" means the total amount of each active component of the method that is sufficient to show a meaningful patient benefit, i.e., healing of acute conditions characterized by inhibition of  $\beta$ -amyloid peptide production. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously. The terms "treat, treating, treatment" as used herein and in the claims means preventing or ameliorating diseases associated with  $\beta$ -amyloid peptide.

### General Reaction Schemes

The general procedures used to synthesize the compounds of Formula I are described in Reaction Schemes 1-23. Reasonable variations of the described

procedures, which would be evident to one skilled in the art, are intended to be within the scope of the present invention.

### Reaction Scheme 1



The starting ( $\alpha$ -amino)acetamides of Formula II are used in racemic or in enantiomerically pure form and are commercially available or are prepared by well-known literature procedures from commercially available ( $\alpha$ -amino)acids (general reference for amide preparation: R.C. Larock "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 - 976; see also Reaction Scheme 18 for the conversion of the acid of Formula XLVIII to the amide of Formula XLIX). The compound of Formula II is treated with a suitable base and a sulfonylating reagent such as a sulfonyl chloride in an aprotic solvent such as  $\text{CH}_2\text{Cl}_2$  at room temperature to generate the ( $\alpha$ -sulfonamido)acetamide of Formula III. Suitable bases include triethylamine and pyridine.

In one method for conversion of the compound of Formula III to the sulfonamide of Formula I, the compound of Formula III is treated with a suitable base and an alkylating agent in an aprotic solvent with or without heating.


Suitable bases for this reaction include potassium carbonate and cesium carbonate. Alkylating agents include alkyl halides (e.g., alkyl chloride, alkyl bromide, or alkyl iodide) and alkyl sulfonates (tosylates, mesylates, trifluoromethanesulfonates). Preferred solvents include DMF and acetonitrile.

The temperature range for the reaction is typically 20 °C to 100 °C.

An alternative method for conversion of the compound of Formula III to the compound of Formula I involves treatment of the compound of Formula III

with triphenyl phosphine, a dialkyl azodicarboxylate, and an alcohol in an inert solvent with or without heating.


### Reaction Scheme 1-solid support



The compounds of Formula I can also be prepared using solid phase methodology. For example, Fmoc-protected Rink amide resin is treated with piperidine in DMF to effect removal of the Fmoc group. The resin is then coupled with an amino-protected ( $\alpha$ -amino) acid in the presence of a coupling agent such as 1-hydroxybenzotriazole and a dialkyl carbodiimide in an inert solvent such as DMF with or without heating. Deprotection of the  $\alpha$ -amino group affords the polymer-bound amide of Formula IV. In the case of an Fmoc-protected amino acid, the deprotection can be accomplished by treatment with piperidine in DMF.


Reaction of the compound of Formula IV with an appropriate base such as pyridine and a sulfonylating agent such as a sulfonyl chloride in an inert solvent provides the resin-linked sulfonamide of Formula V. Alkylation of the compound of Formula V with an alkyl halide (e.g., alkyl chloride, alkyl bromide, or alkyl iodide) or alkyl sulfonate (e.g., mesylate, tosylate, or trifluoromethanesulfonate) is carried out in the presence of a base in an inert solvent at room temperature. A preferred base is 2,4-butyldiamino-2,3-dimethylpentano-1,3,2-diaaphosphorine. Cleavage from the resin provides the sulfonamide of Formula I. In the case of the Rink amide resin, the cleavage is preferably carried out using trifluoroacetic acid in an inert solvent such as  $\text{CH}_2\text{Cl}_2$ .

### Reaction Scheme 2



The compounds of Formula I can also be prepared as shown in Reaction Scheme 2. Reductive alkylation of the amine of Formula I to provide the amine of Formula VI is effected by treatment with an aldehyde and a hydride reducing agent in the presence of an acid catalyst with or without heating. A preferred reducing agent is sodium cyanoborohydride. A preferred acid catalyst is a Lewis acid such as  $\text{ZnCl}_2$ . The reaction solvent is preferably methanol. The amine of Formula VI is then treated with a sulfonylating agent such as a sulfonyl chloride in the presence of an amine such as triethylamine. This reaction is carried out in an inert solvent such as  $\text{CH}_2\text{Cl}_2$  with or without heating to afford the product of Formula I. The reaction is typically carried out at room temperature.


### Reaction Scheme 3



wherein linker = straight-chain or branched  $\text{C}_1$ -alkyl or  $\text{C}_2$ -alkenyl;  
LG = leaving group

Preparation of compounds of Formula VIII is accomplished as shown in Reaction Scheme 3 by reaction of the compound of Formula VII with an amine in the presence of an acid scavenger such as triethylamine in an inert solvent such as  $\text{CH}_2\text{Cl}_2$  with or without heating. The compound of Formula VII is prepared by the sequence shown in Reaction Scheme 1 or Reaction Scheme 2.

- 14 -

Reaction Scheme 4

5 The compounds of Formula XI and XII are prepared as shown in Reaction

Scheme 4. Reduction of the nitro group of the compound of Formula IX

(prepared by the sequence shown in Reaction Scheme 1 or 2) with hydrogen gas

under pressure in the presence of a palladium catalyst, acid, and in a solvent such

as methanol provided the aniline derivative of Formula X. Monomethylation of

10 the compound of Formula X to provide the compound of Formula XI is

accomplished by reaction with 1.1 equivalents of a methyl halide or a methyl

sulfonate, for example dimethylsulfate, in the presence of a base such as

triethylamine and in an inert solvent such as DMF. The monomethylation

reaction is typically carried out between 20 °C and 40 °C. Preparation of the


15 dimethylamine of Formula XII is effected by treatment of the aniline of Formula

X with an excess of a methyl halide such as methyl iodide or a methyl sulfonate

in the presence of a base, for example cesium carbonate, in a solvent such as

DMF, with or without heating.

- 15 -

Reaction Scheme 5

5 Reaction Scheme 5 outlines the synthesis of esters of Formula XIII, acids

of Formula XIV, and amides of Formula XV. Reaction of a compound of

Formula III with a haloalkylcarboxylate ester, for example *t*-butyl bromoacetate,

in the presence of a base such as potassium carbonate and in an inert solvent such

as DMF affords the ester of Formula XIII. Deprotection of the ester is effected

10 by methods known to those skilled in the art (ref.: T.W. Greene and P.G.M.

Wuts, "Protecting Groups in Organic Synthesis", Wiley Interscience, New York,

1999, pp. 373 - 442). For example, for *t*-butyl esters, cleavage to the acid of

Formula XIV is accomplished by treatment with trifluoroacetic acid in a solvent

such as CH<sub>2</sub>Cl<sub>2</sub>. Conversion of the acid to the amide of Formula XV is carried

15 out using common amide coupling procedures well known to those skilled in the


art (ref.: R.C. Larock "Comprehensive Organic Transformations, VCH

Publishers, New York, 1989, pp. 972 - 976). In a preferred procedure, the acid of

Formula XIV is treated with a primary or secondary amine in the presence of 1-

hydroxybenzotriazole and 1,3-dicyclohexylcarbodiimide in an aprotic solvent

20 such as CH<sub>2</sub>Cl<sub>2</sub> or DMF.

Reaction Scheme 6

5 The preparation of acids of Formula XVII and amides of Formula XVIII is shown in Reaction Scheme 6. Conversion of an ester of Formula XVI (prepared as shown in Reaction Schemes 1 or 2) to an acid of Formula XVII is accomplished using standard ester cleavage conditions well known to those

skilled in the art (ref.: T. W. Greene and P. G.M. Wuts, "Protecting Groups in


10 Organic Synthesis", Wiley Interscience, New York, 1999, pp. 373 - 442). In the case of a methyl ester of Formula XVI, treatment with aqueous sodium hydroxide in a solvent such as methanol or a methanol/THF mixture at 20 °C to 40 °C provides the acid of Formula XVII. Conversion of the acid of Formula XVII to

the amide of Formula XVIII is achieved using common amide coupling procedures well known to those skilled in the art (ref.: R. C. Larock

15 "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 - 976). In a preferred procedure, the acid of Formula XVII is treated with

a primary or secondary amine in the presence of 1-hydroxybenzotriazole and a

20 carbodiimide, for example 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, in a solvent such as DMF or  $CH_2Cl_2$ . A base such as diisopropylethylamine can be added as an acid scavenger.

Reaction Scheme 7

5 The synthesis of piperidine derivatives of Formula XIX, XX, XXI, XXII, and XXIII is described in Reaction Scheme 7. Reaction of a compound of

Formula III with an N-protected piperidine substituted with a 4-haloalkyl or 4-

sulfonyloxyalkyl group, such as 4-(toluenesulfonyloxyethyl)-1-(4-

10 butoxycarbonyl)piperidine, in the presence of a base such as cesium carbonate in a solvent such as DMF, with or without heating, provides the carbonate of

Formula XIX. Cleavage of the carbonate group in the compound of Formula

XIX is carried out under standard conditions well known to those skilled in the

art (ref.: T. W. Greene and P. G.M. Wuts, "Protecting Groups in Organic

15 Synthesis", Wiley Interscience, New York, 1999, pp. 503-550) to provide the piperidine of Formula XX. In the case of a (4-butoxycarbonyl)piperidine

derivative, the cleavage is effected by treatment with trifluoroacetic acid in

$CH_2Cl_2$ .

Conversion of the piperidine of Formula XX to an amide of Formula XXI


is carried out using amide-coupling procedures well known to those skilled in the

20 art (ref.: R. C. Larock "Comprehensive Organic Transformations, VCH

- 18 -

Publishers, New York, 1989, pp. 972 – 976). In a preferred method, the piperidine of Formula XX is treated with an acyl chloride in the presence of an amine such as triethylamine and in an inert solvent such as  $\text{CH}_2\text{Cl}_2$  with or without heating. Alternatively, the piperidine of Formula XX may be coupled with an acid in the presence of coupling agents such as hydroxybenzotriazole and a carbodiimide to provide an amide of Formula XXI. Preparation of the urea of Formula XXII is achieved by treatment of the amine of Formula XX with an isocyanate and a base such as triethylamine in a solvent such as  $\text{CH}_2\text{Cl}_2$  with or without heating. Alkylation of the piperidine of Formula XX provides N-substituted piperidines of Formula XXIII. In a typical procedure, the piperidine is treated with an alkyl halide or an alkyl sulfonate in the presence of a base such as triethylamine and in a solvent such as  $\text{CH}_2\text{Cl}_2$ .

#### Reaction Scheme 8




Alcohols of Formula XXV and amines of Formula XXVI are synthesized by the sequence shown in Reaction Scheme 8. A protected alcohol of Formula XXIV is prepared by the procedure shown in Reaction Schemes 1 or 2.

Deprotection of the alcohol under the appropriate conditions for the chosen protecting group (ref.: T.W. Greene and P.G.M. Wuts, "Protecting Groups in Organic Synthesis", Chapter 2) provides the alcohol of Formula XXV. For example, when the protecting group is a tetrahydropyranyl moiety, the alcohol is liberated by treatment of the compound of Formula XXIV with *p*-toluenesulfonic acid in a solvent such as methanol. The alcohol of Formula XXV is converted to

- 19 -

a leaving group (e.g., a halide or sulfonate) and then treated with a primary or secondary amine to afford an amine of Formula XXVI. For example, the alcohol may be converted to a mesylate derivative by reaction with methanesulfonyl chloride and a base such as triethylamine in  $\text{CH}_2\text{Cl}_2$ . Subsequent reaction of the mesylate with a primary or secondary amine in the presence of a base such as triethylamine in a solvent such as  $\text{CH}_2\text{Cl}_2$  provides the amine of Formula XXVI.

#### Reaction Scheme 9




Amines of Formula XXVII are prepared from amines of Formula XXVII as shown in Reaction Scheme 9. Amines of Formula XXVII wherein D is a direct bond are prepared as in Reaction Scheme 1 or 4. Amines of Formula XXVII wherein D is other than a direct bond are prepared as in Reaction Scheme 8.

Conversion of the amines of Formula XXVII to the amides of Formula XXVIII is carried out using amide-coupling conditions well known to those skilled in the art (ref.: R.C. Larock "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 – 976). For example, reaction of the amine of Formula XXVII with an acid chloride in the presence of a base such as triethylamine in a solvent such as  $\text{CH}_2\text{Cl}_2$  provides the amide of Formula XXVIII. Conversion of the amines of Formula XXVII to carbamate derivatives can be carried out using conditions well known to those skilled in the art. (ref.: T.W. Greene and P.G.M. Wuts, "Protecting Groups in Organic Synthesis", P. 503 -


350). Preparation of sulfonamide derivatives from an amine of Formula XXVII can also be achieved using methods such as that described for the conversion of the intermediate of Formula II to the sulfonamide of Formula III.

#### 5 Reaction Scheme 10



The synthesis of pyridine derivatives of Formula XXX is accomplished as shown in Reaction Scheme 10. The chloropyridine derivative of Formula XXIX is prepared using the chemistry described in Reaction Schemes 1 or 2. Treatment of the compound of Formula XXIX with a primary or secondary amine in a solvent such as THF at temperatures from 20 °C to 100 °C, using sealed, pressurized vessel as appropriate, provides the aminopyridine of Formula XXX.

#### 15 Reaction Scheme 11




Amine-substituted phenol ethers of Formula XXXII are prepared from (O-allyl)phenols as indicated in Reaction Scheme 11. The starting allyl ethers of


Formula XXXI are prepared as shown in Reaction Schemes 1 or 2. Treatment of the compound of Formula XXXI with osmium tetroxide and trimethylamine N-oxide in a solvent such as acetone followed by treatment with sodium periodate gives an intermediate aldehyde that is typically used without purification.

5 Reaction of the unpurified aldehyde with a primary or secondary amine and a reducing agent such as sodium triacetoxyborohydride in a solvent such as ethanol with or without heating affords the amine of Formula XXXII.


#### 10 Reaction Scheme 12



Conversion of the ester of Formula XXXIII to the tertiary alcohol of Formula XXXIV is carried out as shown in Reaction Scheme 12. Reaction of the ester of Formula XXXIII with an excess of a methyl organometallic reagent such as methyl magnesium bromide in a solvent such as THF at a temperature ranging from 0 °C to 25 °C yields the alcohol of Formula XXXIV.


Reaction Scheme 13

5 Preparation of the 1,3,4-oxadiazole of Formula XXXVI is carried out as shown in Reaction Scheme 13 using methods well known to those skilled in the art (ref: Joule, J.A.; Mills, K.; Smith, G.F. *Heterocyclic Chemistry*, 3rd ed., Chapman & Hall: London, 1995; 452-456 and references cited therein). For example, the ester of Formula XXXV is treated with hydrazine in methanol with heating up to the reflux point. The resulting acyl hydrazine intermediate is used without purification in a subsequent reaction with an alkyl acetimidate in pyridine with heating at reflux to provide the oxadiazole of Formula XXXVI.

Reaction Scheme 14


15 Synthesis of the 1,2,4-oxadiazole of Formula XXXVII is achieved as shown in Reaction Scheme 14 using methods well known to those skilled in the art (ref: Joule, J.A.; Mills, K.; Smith, G.F. *Heterocyclic Chemistry*, 3rd ed.,

5 Chapman & Hall: London, 1995; 452-456 and references cited therein). For example, treatment of the acid of Formula XVII with hydrobenzotriazole, a carbodiimide, and acetamidoxime (N'-hydroxy ethanimidamide) in the presence of a base such as triethylamine provides an intermediate that is heated in refluxing pyridine to provide the oxadiazole of Formula XXXVII.


Reaction Scheme 15

10 The 1,2,4-oxadiazole of Formula XXXIX is prepared from the nitrile of Formula XXXVIII (Reaction Scheme 15) using methods well-known to those skilled in the art (ref: Joule, J.A.; Mills, K.; Smith, G.F. *Heterocyclic Chemistry*, 3rd ed., Chapman & Hall: London, 1995; 452-456 and references cited therein). For example, reaction of the nitrile of Formula XXXVIII with hydroxylamine in a solvent such as ethanol at temperatures up to reflux provides an intermediate N-hydroxyamidine that is subsequently treated with acetyl chloride in the presence of a base such as triethylamine in a solvent such as CH<sub>2</sub>Cl<sub>2</sub> to provide the 1,2,4-oxadiazole of Formula XXXIX.




**Reaction Scheme 16**

Reaction Scheme 16 shows the transformation of the amide of Formula XL to the ketone of Formula XLI. The amide of Formula XL, which is prepared as described in Reaction Scheme 6, is treated with a methyl organometallic reagent such as methyl magnesium bromide in a solvent such as THF to provide the ketone of Formula XLI. The range of the reaction temperature is from -20 °C to 25 °C.

**Reaction Scheme 17**

$\beta$ -Amino amides of Formula XLIII are prepared from acrylamides of Formula XLII as shown in Reaction Scheme 17. For example, an acrylamide of Formula XLII, which is prepared as described in Reaction Scheme 9, is treated

with a primary or secondary amine in a solvent such as toluene to provide the  $\beta$ -amino amide of Formula XLIII.


**Reaction Scheme 18**

Preparation of the sulfonamide intermediate of Formula XLIX (a single enantiomer of the compound of Formula III) is outlined in Reaction Scheme 18.

Reaction of the  $\alpha$ -anion of the intermediate of Formula XLIV (ref: Josien, H.; Martin, A.; Chassaing, G. *Tetrahedron Lett.* 1991, 32, 6547) with an alkylating agent such as an alkyl halide (e.g., an alkyl chloride, alkyl bromide, or alkyl iodide) or an alkyl sulfonate (e.g., an alkyl mesylate, alkyl tosylate, or alkyl trifluoromethanesulfonate) provides the intermediate of Formula XLV. The  $\alpha$ -anion of the compound of Formula XLIV is formed by treatment with a strong base such as an alkyl lithium (e.g., *n*-BuLi) or a dialkylamide (e.g., lithium diisopropylamide) in a solvent such as THF with or without a co-solvent such as HMPA. The reaction temperature is typically between -78 °C and 25 °C. Removal of the benzhydrylene protecting group of the compound of Formula


XLV is carried out under conditions well known to those skilled in the art (ref.: T. W. Greene and P. G. M. Wuts, "Protecting Groups in Organic Synthesis", Wiley Interscience, New York, 1999, pp. 587- 588). For example, the compound of Formula XLV is treated with an acid such as HCl in water in a solvent such as THF to effect hydrolysis of the benzhydrydene protecting group. The resulting amine of Formula XLVI is treated with a sulfonylating agent as described for Reaction Scheme 1 to provide the sulfonamide of Formula XLVII. Hydrolysis of the acylsulfonamide of Formula XLVII to afford the acid of Formula XLVIII is carried out by treatment with hydroxide ion, for example in the form of lithium hydroxide, in the presence of additives such as lithium bromide and tetrabutylammonium bromide. The acid of Formula XLVIII is converted to the amide of formula XLIX under conditions that are well known to those skilled in the art (general ref for amide preparation.: R. C. Larock "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 - 976). For example, reaction of the compound of Formula XLVII with ammonium chloride in the presence of 1-hydroxybenzotriazole, a carbodiimide reagent, and an amine base such as diisopropylethylamine provides the amide of Formula XLIX. This reaction is typically run in a polar solvent such as DMF and at a reaction temperature from 0 °C to 40 °C. The amide of Formula XLIX is converted to the compounds of Formula I by the method described in Reaction Scheme 1.

### Reaction Scheme 19



Reaction Scheme 19 illustrates one method for synthesis of an  $\alpha$ -substituted (N-sulfonamido)acetamide intermediate of Formula III starting with an activated glycine derivative of Formula I. Reaction of the compound of Formula I (ref.: Haufe, G.; Laue, K. W.; Tiller, M. U.; Takeuchi, Y.; Shibata, N. *Tetrahedron* 1998, 54, p. 5929-5938; Kroger, S.; Haufe, G. *Amino Acids* 1997, 12, p. 363-372) with an alkylating agent such as an alkyl halide (e.g., an alkyl chloride, alkyl bromide, or alkyl iodide) or an alkyl sulfonate (e.g., an alkyl mesylate, an alkyl tosylate, or an alkyl trifluoromethanesulfonate) in the presence of a base such as potassium carbonate and an additive such as tetrabutylammonium bromide and in an inert solvent such as acetonitrile at a reaction temperature of between 25 °C and 70 °C provides the compound of Formula LI. Removal of the benzhydrydene protecting group is carried out under conditions well known to those skilled in the art (ref.: T. W. Greene and P. G. M. Wuts, "Protecting Groups in Organic Synthesis", Wiley Interscience, New York, 1999, pp. 587- 588). For example, a solution of the compound of Formula LI in a solvent such as diethyl ether is treated with aqueous acid (e.g., aqueous HCl), typically at a reaction temperature of between 0 °C and 30 °C, to provide the amine ester of Formula LII. Conversion of the ester of Formula LII to the amide of Formula II is carried out using procedures well known to those skilled in the art. For example, when the compound of Formula LII is an ethyl ester, hydrolysis of the ester is achieved by treatment of an ethereal solution with an acid such as HCl, typically with heating of the reaction mixture in refluxing solvent. The resulting acid intermediate is then converted to a methyl ester of Formula LII by transformation to the acid chloride under standard conditions, (e.g., treatment with thionyl chloride and methanol), followed by reaction with aqueous ammonia in a solvent such as toluene (ref.: R. C. Larock "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 - 976). The amine of Formula II is converted to the compound of Formula I as described in Reaction Scheme 1.

### Reaction Scheme 20




**5 Preparation of the compound of Formula I, VII is shown in Reaction**

Scheme 20. Alkene **LIII** is prepared as described in Reaction Scheme 18 from an intermediate of Formula **XLIV** and 1-bromo-2-methyl-2-propene). Treatment of the alkene of Formula **LIII** with HF-pyridine in a solvent such as THF at a

reaction temperature between 0 °C and 25 °C affords the fluoralkyl compound of Formula LV. Conversion of the compound of Formula LIV to the amide of Formula LV is accomplished as described in Reaction Scheme 18.

Transformation of the amide of Formula LV to the compound of Formula LVI is carried out as described in Reaction Scheme 1.

### Reaction Scheme 21



5 The syntheses of the compounds of Formula LXXII and Formula LXXIV are

outlined in Reaction Scheme 21. Ethyl 2-amino-4-methyl-4-pentenoate (prepared as in Reaction Scheme 19 from (benzylidene)dimethylamino)acetic acid ethyl ester and 1-bromo-2-methyl-2-propene) is treated with a sulfonylating agent such as a sulfonyl chloride in the presence of a base such as triethylamine in an inert

solvent such as  $\text{CH}_2\text{Cl}_2$ , to afford the ester of Formula L.VII. Reaction of the ester of Formula L.VII with HF-pyridine in a solvent such as THF and at a reaction temperature of between 0 °C and 25 °C provides a mixture of the fluoroalkyl derivative of Formula L.VIII and the lactone of Formula L.IX. These products are separated and carried on individually into subsequent reactions.


The ester of Formula LVIII is hydrolyzed to the acid of Formula LX using methods well known to those skilled in the art (ref.: T. W. Greene and P. G. M. Wuts, "Protecting Groups in Organic Synthesis", Wiley Interscience, New York, 1999, pp. 373 - 442). For example, treatment of the ester of Formula LVIII with aqueous sodium hydroxide in a solvent such as methanol affords the acid of

- 30 -

Formula LX. The acid of Formula LX is converted to the amide of Formula LXI using the procedure described in Reaction Scheme 18 for the preparation of the amide of Formula XLIX. Preparation of the amide of Formula LXII from the compound of Formula LXI is achieved as described in Reaction Scheme 1.

For the lactone of Formula LIX, treatment with aqueous ammonia provides the amide of Formula LXIII. This reaction is typically carried out with heating in a sealed tube. The reaction temperature is between 40 °C and 80 °C. Further conversion of the intermediate of Formula LXIII to the sulfonamide of Formula LXIV proceeded as described in Reaction Scheme 1.

#### Reaction Scheme 22




The synthetic sequence for preparation of a difluoroalkyl amide of Formula LXIX is shown in Reaction Scheme 22. The compound of Formula L is treated with 4-bromo-1-butene in the presence of a base such potassium carbonate in the presence of a tetraalkylammonium halide salt such as tetrabutylammonium bromide in a solvent such as CH<sub>3</sub>CN at a temperature from 20 °C to 70 °C. Removal of the benzhydrydene protecting group as described in Reaction Scheme 19 provides an intermediate amine that is then treated with a sulfonylating reagent such as a sulfonyl chloride to provide the ester of Formula LXV. Alkylation of the sulfonamide nitrogen is accomplished using the procedure described in Reaction Scheme 1 to afford the compound of Formula

- 31 -

LXVI. Conversion of the alkene of Formula LXVI to the aldehyde of Formula LXVII is achieved by reaction of the alkene with osmium tetroxide and trimethylamine N-oxide in a solvent such as acetone, followed by treatment with sodium periodate. The reaction temperature is typically 20 °C to 40 °C. Reaction of the aldehyde of Formula LXVII with a fluorinating agent such as DAST in a solvent such as CH<sub>2</sub>Cl<sub>2</sub> yields the difluoroalkyl derivative of Formula LXVIII. The compound of Formula LXVIII is converted to the amide of Formula LXIX by hydrolysis of the ester to an acid using an base such as sodium hydroxide in a solvent such as methanol. The intermediate acid was converted to the amide using conditions well known to those skilled in the art (ref.: R.C. Larock "Comprehensive Organic Transformations, VCH Publishers, New York, 1989, pp. 972 – 976). For example, reaction of the acid with ammonium chloride in the presence of hydroxybenzotriazole and a carbodiimide reagent and an amine base such as diisopropylethylamine provided the amide of Formula LXIX. This reaction is typically run in a polar solvent such as DMF and at a reaction temperature from 0° C to 40° C.


#### Reaction Scheme 23



The α-amino amide of Formula LXXI is prepared using the reaction showed in Reaction Scheme 23. The amide of Formula LXX is prepared as

- 32 -

described in Reaction Scheme 9. Treatment of the compound of Formula LXX with a secondary or tertiary amine in a solvent such as THF at a reaction temperature between 20 °C and 40 °C affords the amine of Formula LXXI.



In a preferred embodiment, the present invention includes compounds of Formula Ia or a pharmaceutically acceptable salt thereof



wherein:

R<sup>1</sup> is selected from the group consisting of

- (a) a straight or branched-chain C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl optionally substituted with substituents selected from the group consisting of hydroxy, C<sub>2-7</sub> cycloalkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkylthio, and halogen;
- (b) C<sub>2-7</sub> cycloalkyl optionally substituted with hydroxy or halogen;

R<sup>2</sup> is selected from the group consisting of

- (a) a straight or branched-chain C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub> alkoxy, and NR<sup>4</sup>R<sup>5</sup>;

- 33 -

- (b) C<sub>2-7</sub> cycloalkylmethyl optionally substituted with substituents selected from the group consisting of amine, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkylthio, C<sub>1-4</sub> alkoxy(C(=O)NH-, and C<sub>1-4</sub> alkyl(C(=O)NH-);

- (c) a straight or branched-chain C<sub>1-6</sub> alkyl-C(=O)-A;

- (d) -B-naphthyl;

- (e)



D and E are each independently a direct bond, a straight or branched-chain C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, or C<sub>2-7</sub> cycloalkyl;

Z is selected from the group consisting of hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-4</sub> alkoxy, halogen, cyano, hydroxy, -OCHF<sub>2</sub>, -OCF<sub>3</sub>, -CF<sub>3</sub>, and

-CHF<sub>2</sub>;

X and Y are each independently selected from the group consisting of hydrogen, hydroxy, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, C<sub>1-6</sub> alkylS-, C<sub>1-6</sub> alkylS(O)-, C<sub>1-6</sub> alkylSO<sub>2</sub>-, nitro, F, S-, and cyano;

-OR<sup>6</sup>;

-NR<sup>6</sup>R<sup>7</sup>;

-NR<sup>6</sup>C(=O)R<sup>8</sup>;

-NR<sup>6</sup>C(=O)OR<sup>9</sup>;

-NHSO<sub>2</sub>C<sub>1-6</sub>alkyl;

-N(SO<sub>2</sub>C<sub>1-6</sub>alkyl)<sub>2</sub>;

-C(=O)W wherein W is selected from the group consisting of

hydroxy, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, phenoxy, and -NR<sup>6</sup>R<sup>7</sup>;

-OC(=O)C<sub>1-6</sub>alkyl;

-phenyl in which said phenyl is optionally substituted with cyano, halogen, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylS-, CH<sub>2</sub>C(=O)-, C<sub>1-6</sub> alkylS(O)-, or C<sub>1-6</sub> alkylSO<sub>2</sub>-; and

- 34 -

heterocyclic group, in which said heterocyclic group is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, isoxazolyl, thiadiazolyl, and thiazolyl, wherein said heterocyclic group is optionally substituted with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}$ alkyl, (halogen) $C_{1-4}$ alkyl, and  $CO_2C_{1-4}$ alkyl;

- (i) -B-(heterocycle), in which said heterocycle is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, isoxazolyl, thiadiazolyl and thiazolyl wherein said heterocycle is optionally substituted with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}$ alkyl,  $CO_2C_{1-4}$ alkyl, amino,  $(C_{1-4}alkyl)NH_2$ ,  $di(C_{1-4}alkyl)N_2$ , morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4- $(C_{1-4}alkyl)piperazin-1-yl$ ;

- (8) -B-(piperidin-4-yl), in which said piperidin-4-yl is optionally substituted with substituents selected from the group consisting of a straight or branched-chain  $C_{1-6}$ alkyl,  $CH_2C(=O)phenyl$ , phenyl and phenylmethyl in which said  $C_{1-6}$ alkyl and said phenyl are optionally substituted with substituents selected from the group consisting of cyano, halogen, benzimidazol-2-yl, pyridyl and tetrahydrofuran-2-yl; and -C(=O)W' wherein W' is selected from the group consisting of  $C_{1-4}$ alkoxy,  $R^1$ , and -NR<sup>1</sup>R<sup>2</sup>;

- 25 A is hydroxy,  $C_{1-4}$ alkoxy or NR<sup>1</sup>R<sup>2</sup>;  
B is a straight or branched-chain  $C_{1-6}$ alkyl or  $C_{3-6}$ alkenyl;  
R<sup>1</sup> is phenyl or pyridyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkyl, (halogen) $C_{1-4}$ alkyl, and halogen $CH_2$ ;  
30 R<sup>1</sup> and R<sup>2</sup> each are independently hydrogen, a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-6}$ alkenyl,  $C_{3-6}$ cycloalkyl,  $C_{3-6}$ cycloalkylmethyl,

- 35 -

- $C_{1-4}$ alkoxy, phenyl, benzyl, pyridyl, piperidin-4-yl, inden-1-yl, inden-2-yl, tetrahydrofuran-3-yl, or pyrrolidin-3-yl; in which each is optionally substituted with substituents selected from the group consisting of hydroxy, cyano, halogen, (halogen) $C_{1-4}$ alkoxy, (halogen) $CH_2$ , hydroxymethyl, benzyloxy, methyl, phenyl, pyridyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, (halogen) $C_{1-4}$ alkoxy, (halogen) $CH_2O$ ,  $C_{1-4}$ alkylthio, amino,  $(C_{1-4}alkyl)NH_2$ ,  $di(C_{1-4}alkyl)N_2$ , morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, 4- $(C_{1-4}alkyl)piperazin-1-yl$ , 4-phenylpiperazin-1-yl, 4-benzylpiperazin-1-yl, 4-pyridylpiperazin-1-yl,  $CO_2H$ ,  $CO_2C_{1-4}alkyl$ ,  $C(=O)NHC_{1-4}alkyl$ , and  $C(=O)N(C_{1-4}alkyl)_2$ ;

- R<sup>1</sup> and R<sup>2</sup> taken together may be morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, 1,2,3,4-tetrahydroisoquinolin-2-yl, decalyl, decahydroisoquinolin-1-yl, piperidin-1-yl, piperazin-1-yl, [1,4]-oxazepan-4-yl, azetidin-1-yl, 2,3-dihydro-1H-isindol-2-yl, or 2,3-dihydro-1H-indol-1-yl; in which each is optionally substituted with substituents selected from the group consisting of hydroxy, cyano, halogen, (halogen) $C_{1-4}$ alkoxy, (halogen) $CH_2$ , halogen $CH_2$ , phenyl, pyridyl, benzyl,  $C_{1-4}alkyl$ ,  $C_{3-6}$ cycloalkyl,  $C_{1-4}alkoxy$ ,  $C_{1-4}alkylthio$ , amino,  $(C_{1-4}alkyl)NH_2$ ,  $di(C_{1-4}alkyl)N_2$ ,  $CO_2H$ ,  $CO_2C_{1-4}alkyl$ ,  $C(=O)NHC_{1-4}alkyl$ , and  $C(=O)N(C_{1-4}alkyl)_2$ ;

- 20 R<sup>3</sup> is a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-6}$ alkenyl, benzyl, or phenyl in which each is optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, amino,  $(C_{1-4}alkyl)NH_2$ ,  $di(C_{1-4}alkyl)N_2$ ,  $(C_{1-4}alkyl)(phenyl)N_2$ , morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4- $(C_{1-4}alkyl)piperazin-1-yl$ ;
- 25 R<sup>1</sup> is hydrogen, a straight or branched-chain  $C_{1-6}$ alkyl;  
R<sup>2</sup> is a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl, phenyl, pyridyl, or furanyl; in which each is optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $(C_{1-4}alkyl)NH_2$ ,  $di(C_{1-4}alkyl)N_2$ , morpholin-4-yl, thiomorpholin-4-yl,


pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-6</sub>alkyl)piperazin-1-yl;

R<sup>2</sup> is a straight or branched-chain C<sub>1-6</sub>alkyl, C<sub>3-6</sub> alkenyl, benzyl, phenyl, oxazolyl or pyridyl, in which each is optionally substituted with

substituents selected from the group consisting of halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, amino, (C<sub>1-6</sub>alkyl)NH-, di(C<sub>1-6</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-6</sub>alkyl)piperazin-1-yl; or a non-toxic pharmaceutically acceptable salt thereof.

In another preferred embodiment, the invention includes compounds of Formula Ia or a pharmaceutically acceptable salt thereof wherein R<sup>2</sup> is phenyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkyl, (halogen)<sub>2</sub>C-, and halogenCH<sub>2</sub>-.

In yet another preferred embodiment, the invention includes compounds of Formula Ia or a pharmaceutically acceptable salt thereof wherein R<sup>2</sup> is



20


### BIOLOGICAL TESTING METHODS

Compounds of Formula (I) are expected to possess  $\gamma$ -secretase inhibitory activity. The detection of  $\gamma$ -secretase activity requires assays capable of reliable, accurate and expedient detection of  $\gamma$ -secretase cleavage products, particularly AB. The  $\gamma$ -secretase inhibitory activity of the compounds of the present invention is demonstrated using assays for such activity, for example, using the assays described below. Compounds within the scope of the present invention have been shown to inhibit the activity of  $\gamma$ -secretase, as determined using assays for such activity.

Compounds provided by this invention should also be useful as standards and reagents in determining the ability of a potential pharmaceutical to inhibit AB production. These would be provided in commercial kits comprising a compound of this invention.

#### In vitro binding assay to identify $\gamma$ -secretase inhibitors.

Competitive binding assays can be used to identify molecules that inhibit the binding of a radiolabeled  $\gamma$ -secretase inhibitor and therefore inhibit  $\gamma$ -secretase activity. For example, [<sup>3</sup>H]-Compound A can be used for binding assays with membranes from THP-1 cells (Seiffert, D.; Bradley, J. et al., *J. Biol. Chem.* 2000, **275**, 34086-34091). Compound A is (2R,3S) N1-[(3S)-hexahydro-1-(3-phenoxycyclopropyl)-2-oxo-1H-azepin-3-yl]-2-(2-methylpropyl)-3-(propyl)-butanediamide, the synthesis of which is described in U.S. patent US6331408 (12/18/2001); PCT Publication WO 00/28331; PCT Publication WO 00/07995; and Seiffert, D., Bradley, J. et al., *J. Biol. Chem.* 2000, **275**, 34086-34091.



Compound A

20

For these assays, THP-1 cells were grown in spinner cultures in RPMI 1640 containing L-glutamine and 10  $\mu$ M  $\beta$ -mercaptoethanol to a density of 5 x 10<sup>6</sup> cells/ml. Cells were harvested by centrifugation and cell pellets were quick frozen in dry ice/ethanol and stored at -70 °C prior to use. The pellets of approximately 2 x 10<sup>4</sup> THP-1 cells were homogenized using a Brikman Polytron at setting 6 for 10 sec. The homogenate was centrifuged at 48,000 x g for 12 min, and the resulting pellet was washed by repeating the homogenization

and centrifugation. The final cell pellet was resuspended in buffer to yield a protein concentration of approximately 0.5 mg/ml. Assays were initiated by the addition of 150  $\mu$ l of membrane suspension to 150  $\mu$ l of assay buffer containing 0.064  $\mu$ Ci of radioligand and various concentrations of unlabeled compounds.

- 5 Binding assays were performed in duplicate in polypropylene 96-well plates in a final volume of 0.3 ml containing 50 mM Hepes, pH 7.0, and 5% dimethyl sulfoxide. Nonspecific binding was defined using incubations with 300 nM compound A (Seifert, D., Bradley, J. et al., *J. Biol. Chem.* 2000, 275, 34086-34091). After incubating at 23 °C for 1.3 hr, bound ligand was separated from free radioligand by filtration over GFF glass fiber filters presoaked in 0.3% ethylenimine polymer solution. Filters were washed three times with 0.3 ml of ice cold phosphate-buffered saline, pH 7.0, containing 0.1% Triton X-100. Filter-bound radioactivity was measured by scintillation counting.  $IC_{50}$  values were then determined and used to calculate  $K_i$  values using the Cheng-Prusoff correction for  $IC_{50}$  values. Compounds were scored as active  $\gamma$ -secretase inhibitors if  $K_i$  values were less than 10  $\mu$ M.

- 15 Examples of the results obtained when the invention compounds are subjected to the above described assay are shown in Table 1. In the table, an inhibitory concentration ( $IC_{50}$ ) of less than or equal to 50 nM is represented by +++; between 50 nM and 500 nM by ++; between 500 nM and 10000 nM by +.

**TABLE 1: Examples of activity in the *in vitro* binding Assay**

EXAMPLE	ACTIVITY RATING <sup>a</sup>
96	+++
123	+++
159	+++
315	++
341	++
357	++

EXAMPLE	ACTIVITY RATING <sup>a</sup>
362	+++
365	+++
366	+++
367	+
376	++
379	+++
385	+++
389	+++
394	+++
403	++
405	+++
408	+
409	++
437	+++
441	+++
443	++
445	+++
447	+++
450	++
451	+
452	++
457	++
464	+
474	+++
476	+++
479	++
486	+++

<sup>a</sup> Activity based on  $IC_{50}$  values:



- 40 -

+++	=	<50 nM
++	=	50 - 500 nM
+	=	>500 nM and <10,000 nM

##### 5 In vitro assay to identify $\gamma$ -secretase inhibitor based on the inhibition of A $\beta$ formation from membrane preparations.

An isolated membrane fraction which contains functionally active  $\gamma$ -secretase and  $\beta$ -APP substrates can generate  $\gamma$ -secretase cleavage products including A $\beta$  (Roberts, S.B.; Hendrick, J. P.; Vinitzky, A.; Lewis, M.; Smith, D. W.; Pak, R. PCT Publication WO 01/0175435; Fichteler, K.; Kosika, M.; Fuchs, M. Patent Application No. DE 99-19941039; Shearman, M.; Behr, D. et al., *Biochemistry*, 2000, 39, 8698-8704; Zhang, L. Song, L. et al., *Biochemistry* 2001, 40, 5049-5053). An isolated membrane fraction can be prepared from human derived cell lines such as HeLa and H4 which have been transfected with wild type or mutant forms of  $\beta$ -APP or a human alkaline phosphatase  $\beta$ -APP fusion construct, and stably express high levels of  $\gamma$ -secretase substrates. The endogenous  $\gamma$ -secretase present in the isolated membranes prepared at 0-4 °C cleaves the  $\beta$ -APP substrates when the membranes are shifted from 0-4 to 37 °C. Detection of the cleavage products including A $\beta$  can be monitored by standard techniques such as immunoprecipitation (Citron, M.; Diehl, T.S. et al., *Proc. Natl. Acad. Sci. USA*, 1996, 93,13170-13175), western blot (Klaffki, H. W.; Ambrowski, D. et al., *J. Biol. Chem.*, 1996, 271, 28655-28659), enzyme linked immunosorbent assay (ELISA) as demonstrated by Seubert, P.; Vigo-Pelfrey, C. et al., *Nature*, 1992, 359, 325-327, or by a preferred method using time-resolved fluorescence of the homogeneous sample containing membranes and A $\beta$  (Roberts, S.B.; Hendrick, J. P.; Vinitzky, A.; Lewis, M.; Smith, D. W.; Pak, R. PCT Publication WO 01/0175435; Shearman, M.; Behr, D. et al., *Biochemistry*, 2000, 39, 8698-8704). The A $\beta$  present in a homogeneous sample containing membranes can be detected by time-resolved fluorescence with two antibodies that recognize different epitopes of A $\beta$ . One of the antibodies

- 41 -

recognizes an epitope that is present in A $\beta$  but not present in the precursor fragments; preferably the antibody binds the carboxyl terminus of A $\beta$  generated by the  $\gamma$ -secretase cleavage. The second antibody binds to any other epitope present on A $\beta$ . For example, antibodies that bind the N-terminal region (e.g., 26D6-B2-B3<sup>®</sup> SIBIA Neurosciences, La Jolla, CA) or bind the C-terminal end (e.g., 9S3.2<sup>®</sup> antibody, Biosolutions, Newark, DE) of the A $\beta$  peptide are known. The antibodies are labeled with a pair of fluorescent adducts that transfer fluorescent energy when the adducts are brought in close proximity as a result of binding to the N- and C-terminal ends or regions of A $\beta$ . A lack of fluorescence is indicative of the absence of cleavage products, resulting from inhibition of  $\gamma$ -secretase. The isolated membrane assay can be used to identify candidate agents that inhibit the activity of  $\gamma$ -secretase cleavage and A $\beta$  production.

A typical membrane-based assay requires 45  $\mu$ g membrane protein per well in a 96- or 384-well format. Membranes in a neutral buffer are combined with the test compound and shifted from 0-4 to 37 °C. Test agents may typically consist of synthetic compounds, secondary metabolites from bacterial or fungal fermentation extracts, or extracts from plant or marine samples. All synthetic agents are initially screened at doses ranging from 10-100  $\mu$ M or in the case of extracts at sufficient dilution to minimize cytotoxicity. Incubation of the membranes with the test agent will continue for approximately 90 minutes at which time fluorescence labeled antibodies are added to each well for A $\beta$  quantitation. The time-resolved fluorescence detection and quantitation of A $\beta$  is described elsewhere (Roberts, S.B.; Hendrick, J. P.; Vinitzky, A.; Lewis, M.; Smith, D. W.; Pak, R. PCT Publication WO 01/0175435; Shearman, M.; Behr, D. et al., *Biochemistry*, 2000, 39, 8698-8704). Results are obtained by analysis of the plate in a fluorescence plate reader and comparison to the mock treated membranes and samples in which known amounts of A $\beta$  were added to construct a standard concentration curve. A positive acting compound is one that inhibits the A $\beta$  relative to the control sample by at least 50% at the initial tested concentration. If a compound is found to be active then a dose response

experiment is performed to determine the lowest dose of compound necessary to elicit the inhibition of the production of A $\beta$ . Compounds were scored as active  $\gamma$ -secretase inhibitors if K<sub>i</sub> values were less than 10  $\mu$ M.

Examples of the results obtained when the invention compounds are subjected to the above described assay are shown in Table 2. In the table, an inhibitory concentration (IC<sub>50</sub>) of less than or equal to 50 nM is represented by +++; between 50 nM and 500 nM by ++; between 500 nM and 10000 nM by +.

TABLE 2: Examples of activity in the *in vitro* assay based on the inhibition of A $\beta$  formation from membrane preparations

EXAMPLE	ACTIVITY RATING <sup>a</sup>
1	+++
2	+++
3	+++
4	+++
5	+++
6	+++
7	+++
8	+++
9	+++
10	+++
11	+++
12	+++
13	+++
14	+++
15	+++
16	+++
17	+++

EXAMPLE	ACTIVITY RATING <sup>a</sup>
18	++
19	++
20	++
21	+++
22	+++
23	+++
24	+++
25	+++
26	+++
27	++
28	++
29	+++
30	++
31	++
32	+++
33	+++
34	+++
35	++
36	++
37	+++
38	+++
39	+++
40	+++
41	+++
42	+++
43	+++
44	+++
45	+++

- 44 -

EXAMPLE	ACTIVITY RATING <sup>a</sup>
46	+++
47	+++
48	+++
49	++
50	+++
51	+++
52	+++
59	++
61	+++
83	+
85	+
87	+++
89	+++
95	+++
103	+++
113	++
122	+
133	+++
153	++

<sup>a</sup> Activity based on IC<sub>50</sub> values:

+++ = &lt;50 nM

++ = 50 - 500 nM

+ = &gt;500 nM and &lt;10,000 nM

In vitro assays to identify  $\gamma$ -secretase inhibitor based on the inhibition of A $\beta$  formation in cultured cells.

Cultured human cell lines, such as HEK293 and H4 cells, which express

10 APP and  $\gamma$ -secretase activity or transfected derivative cell lines that overexpress

- 45 -

wild-type APP, mutant APP, or APP fusion proteins will secrete A $\beta$  peptides into the culture media that can be quantified as previously outlined (Dovey, H., John, V. et al., *J. Neurochem.* 2001, 76, 173-181). The incubation of these

cultured cells with  $\gamma$ -secretase inhibitors decreases the production of A $\beta$  peptides.

5 For instance, H4 cells stably transfected to overexpress the HPLA-APP fusion protein described above were grown as above, detached, and adjusted to  $2 \times 10^5$  cells/ml. 100  $\mu$ l of the resulting suspension was then added to each well of a 96-well plate. After 4 hrs, the media was removed and replaced with 100  $\mu$ l serum-free media containing various dilutions of the test compound. Plates were then incubated for 18 hrs at 37 °C and a 100  $\mu$ l aliquot of the tissue culture

10 supernatant was removed for determination of A $\beta$  levels using time-resolved fluorescence of the homogenous sample as outlined above. Alternately, the other methods described above for A $\beta$  determination could be used. The extent of A $\beta$  inhibition was used to calculate the IC<sub>50</sub> value for the test compound.

15 Compounds of the present invention are considered active when tested in the above assay if the IC<sub>50</sub> value for the test compound is less than 50  $\mu$ M.

Examples of the results obtained when the invention compounds are subjected to the above described assay are shown in Table 3. In the table, an inhibitory concentration (IC<sub>50</sub>) of less than or equal to 50 nM is represented by +++; between 50 nM and 500 nM by ++; between 500 nM and 50000 nM by +.

TABLE 3: Examples of activity in the *in vitro* assay based on the inhibition of A $\beta$  formation in cultured cells

EXAMPLE	ACTIVITY RATING <sup>a</sup>
1	+++
5	+++
19	++
26	+++

- 46 -

EXAMPLE	ACTIVITY RATING <sup>a</sup>
38	+++
41	+++
51	+++
55	+++
61	+++
72	+++
80	+++
89	+++
96	+++
101	+++
123	+++
127	++
143	+++
147	++
158	+++
171	++
193	+++
203	+++
205	++
207	+++
245	+++
246	+++
249	++
254	+++
256	+++
260	+++
272	+++
280	++

- 47 -

EXAMPLE	ACTIVITY RATING <sup>a</sup>
282	+++
288	++
301	++
302	+++
321	++
322	+++
329	+++
330	++
331	+
340	+++
341	++
342	+++
349	+++
352	++
358	++
359	+++
366	+++
367	+
378	+++
383	+++
394	+++
403	++
416	+++
418	+++
424	+++
433	+++
434	+++
439	+++

- 48 -

EXAMPLE	ACTIVITY RATING <sup>a</sup>
442	+++
472	+++
481	+
492	++
495	+++
497	+++

<sup>a</sup> Activity based on IC<sub>50</sub> values:

- +++ = <50 nM  
 ++ = 50 - 500 nM  
 + = >500 nM and <10,000 nM

- Compounds of the present invention have been demonstrated to have an IC<sub>50</sub> value less than 10  $\mu$ M in one or all of the above assays. Therefore, the compounds of Formula I or pharmaceutical compositions thereof are useful in the treatment, alleviation or elimination of disorders or other disorders associated with the inhibition of  $\beta$ -amyloid peptide.

- In addition to cleaving APP,  $\gamma$ -secretase cleaves other substrates, including: the Notch family of transmembrane receptors (reviewed in: Selkoe, D. *Physiol. Rev.* 2001, 81, 741-766; Wolfe, M. *J. Med. Chem.* 2001 44, 2039-2060); LDL receptor-related protein (May, P., Reddy, Y.K., Herz, J. *J. Biol. Chem.* 2002, 277, 18736-18743); ErbB-4 (Ni, C.Y., Murphy, M.P., Golde, T.E., Carpenter, G. *Science* 2001, 294, 2179-2181); E-cadherin (Marambaud, P., Shioi, I., et al., *EMBO J.* 2002, 21, 1948-1956); and CD44 (Okamoto, I., Kawano, Y., et al., *J. Cell Biol.* 2001, 155, 755-762). If inhibition of cleavage of non-APP substrates causes undesirable effects in humans, then desired  $\gamma$ -secretase inhibitors would preferentially inhibit APP cleavage relative to unwanted substrates. Notch cleavage can be monitored directly by measuring the amount of cleavage product or indirectly by measuring the effect of the cleavage product on

- 49 -

transcription (Mizutani, T., Taniguchi, Y., et al. *Proc. Natl. Acad. Sci. USA* 2001, 98, 9026-9031).

In vivo assays for the determination of A $\beta$  reduction by  $\gamma$ -secretase inhibitors.

- In vivo assays are available to demonstrate the inhibition of  $\gamma$ -secretase activity. In these assays, animals, such as mice, that express normal levels of APP and  $\gamma$ -secretase or are engineered to express higher levels of APP and hence A $\beta$  can be used to demonstrate the utility of  $\gamma$ -secretase inhibitors (Dovey, H., John, V., et al., *J. Neurochem.* 2001, 76, 173-181). In these assays,  $\gamma$ -secretase inhibitors were administered to animals and A $\beta$  levels in multiple compartments, such as plasma, cerebral spinal fluid, and brain extracts, were monitored for A $\beta$  levels using methods previously outlined. For instance, Tg2576 mice, which overexpress human APP, was administered  $\gamma$ -secretase inhibitors by oral gavage at doses that will cause measurable A $\beta$  lowering, typically less than 100 mg/kg. Three hours after dosing plasma, brain, and CSF were collected, frozen in liquid nitrogen, and stored at -80 °C until analysis. For A $\beta$  detection, plasma was diluted 15-fold in PBS with 0.1% Chaps while CSF was diluted 15-fold in 1% Chaps with protease inhibitors (5  $\mu$ g/ml leupeptin, 30  $\mu$ g/ml aprotinin, 1 mM phenylmethylsulfonylfluoride, 1  $\mu$ M pepstatin). Brains were homogenized in 1% Chaps with protease inhibitors using 24 ml solution/g brain tissue. Homogenates were then centrifuged at 100,000 x g for 1 hr at 4 °C. The resulting supernatants were then diluted 10-fold in 1% Chaps with protease inhibitors. A $\beta$  levels in the plasma, CSF, and brain lysate were measured using time-resolved fluorescence of the homogeneous sample or one of the other methods previously described.
- A  $\gamma$ -secretase inhibitor is considered active in one of the above in vivo assays if it reduces A $\beta$  by at least 50% at a dosage of 100 mg/kg.
- Therefore, the compounds of Formula I or pharmaceutical compositions thereof are useful in the treatment, alleviation or elimination of disorders or other disorders associated with the inhibition of  $\beta$ -amyloid peptide.

In another embodiment, this invention includes pharmaceutical compositions comprising at least one compound of Formula I in combination with a pharmaceutical adjuvant, carrier or diluent.

In still another embodiment, this invention relates to a method of treatment or prevention of disorders responsive to the inhibition of  $\beta$ -amyloid peptide in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound of Formula I or a nontoxic pharmaceutically acceptable salt, solvate or hydrate thereof.

In yet another embodiment, this invention relates to a method for treating Alzheimer's Disease and Down's Syndrome in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound of Formula I or a non-toxic pharmaceutically acceptable salt, solvate or hydrate thereof.

For therapeutic use, the pharmacologically active compounds of Formula I will normally be administered as a pharmaceutical composition comprising as the (or an) essential active ingredient at least one such compound in association with a solid or liquid pharmaceutically acceptable carrier and, optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques.

The pharmaceutical compositions include suitable dosage forms for oral, parenteral (including subcutaneous, intramuscular, intradermal and intravenous) bronchial or nasal administration. Thus, if a solid carrier is used, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The solid carrier may contain conventional excipients such as binding agents, fillers, tableting lubricants, disintegrants, wetting agents and the like. The tablet may, if desired, be film coated by conventional techniques. If a liquid carrier is employed, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule, sterile vehicle for injection, an aqueous or non-aqueous liquid suspension, or may be a dry product for reconstitution with water or other suitable vehicle before use. Liquid preparations may contain conventional additives such as suspending agents, emulsifying

agents, wetting agents, non-aqueous vehicle (including edible oils), preservatives, as well as flavoring and/or coloring agents. For parenteral administration, a vehicle normally will comprise sterile water, at least in large part, although saline solutions, glucose solutions and like may be utilized. Injectable suspensions also may be used, in which case conventional suspending agents may be employed. Conventional preservatives, buffering agents and the like also may be added to the parenteral dosage forms. The pharmaceutical compositions are prepared by conventional techniques appropriate to the desired preparation containing appropriate amounts of the active ingredient, that is, the compound of Formula I according to the invention. See, for example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, 17th edition, 1985.

The dosage of the compounds of Formula I to achieve a therapeutic effect will depend not only on such factors as the age, weight and sex of the patient and mode of administration, but also on the degree of  $\beta$ -AP inhibition desired and the potency of the particular compound being utilized for the particular disorder of disease concerned. It is also contemplated that the treatment and dosage of the particular compound may be administered in unit dosage form and that the unit dosage form would be adjusted accordingly by one skilled in the art to reflect the relative level of activity. The decision as to the particular dosage to be employed (and the number of times to be administered per day) is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect.

A suitable dose of a compound of Formula I or pharmaceutical composition thereof for a mammal, including man, suffering from, or likely to suffer from any condition related to  $\beta$ -AP production as described herein, generally the daily dose will be from about 0.05 mg/kg to about 10 mg/kg and preferably, about 0.1 to 2 mg/kg when administered parenterally. For oral administration, the dose may be in the range from about 1 to about 75 mg/kg and preferably from 0.1 to 10 mg/kg body weight. The active ingredient will preferably be administered in equal doses from one to four times a day. However, usually a small dosage is administered, and the dosage is gradually

increased until the optimal dosage for the host under treatment is determined. In accordance with good clinical practice, it is preferred to administer the instant compounds at a concentration level that will produce an effective anti-amyloid effect without causing any harmful or untoward side effects. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the choice of compound of be administered, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms.

The following examples are given by way of illustration and are not to be construed as limiting the invention in any way inasmuch as many variations of the invention are possible within the spirit of the invention.

#### DESCRIPTION OF THE SPECIFIC EMBODIMENTS

In the following examples, all temperatures are given in degrees

Centigrade. Melting points were recorded on a Thomas Scientific Unimelt

capillary melting point apparatus and are uncorrected. Proton magnetic resonance (<sup>1</sup>H NMR) spectra were recorded on a Bruker Avance 300, a Bruker Avance 400, or a Bruker Avance 500 spectrometer. All spectra were determined in the

solvents indicated and chemical shifts are reported in  $\delta$  units downfield from the internal standard tetramethylsilane (TMS) and interproton coupling constants are reported in Hertz (Hz). Splitting patterns are designated as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad peak; dd, doublet of doublet; br d, broad doublet; dt, doublet of triplet; br s, broad singlet; dq, doublet of quartet. Infrared (IR) spectra using potassium bromide (KBr) or sodium chloride film were determined on a Jasco FT/IR-410 or a Perkin Elmer 2000 FT-IR spectrometer from 4000  $\text{cm}^{-1}$  to 400  $\text{cm}^{-1}$ , calibrated to 1601  $\text{cm}^{-1}$  absorption of


a polystyrene film and reported in reciprocal centimeters ( $\text{cm}^{-1}$ ). Optical

rotations [ $\alpha$ ]<sub>D</sub> were determined on a Rudolph Scientific Autopol IV polarimeter in the solvents indicated; concentrations are given in mg/mL. Low resolution mass

spectra (MS) and the apparent molecular ( $MH^+$ ) or ( $M-H$ )<sup>+</sup> was determined on a Finnegan SSQ7000. High resolution mass spectra were determined on a Finnegan MAT900. Liquid chromatography (LC)/mass spectra were run on a Shimadzu LC coupled to a Water Micromass ZQ.

The following abbreviations are used: DMF (dimethylformamide); THF (tetrahydrofuran); DMSO (dimethylsulfoxide), Leu (leucine); TFA (trifluoroacetic acid); DAST [(diethylamino)sulfur trifluoride], HPLC (high pressure liquid chromatography); rt (room temperature); aq. (aqueous).


#### 10 Exemplification of Reaction Scheme 1




#### 15 (2R)-2-(4-Chlorobenzensulfonylamino)-4-methylpentanoic acid amide:

To a solution of (*D*)-leucinamide hydrochloride (0.25 g, 1.5 mmol), and Et<sub>3</sub>N (0.43 mL, 3.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (150 mL) was added 4-chlorobenzene-sulfonyl chloride (380 mg, 1.8 mmol). The resulting solution was stirred at rt for 18 h. The reaction was then diluted with CH<sub>2</sub>Cl<sub>2</sub> (200 mL) and washed with H<sub>2</sub>O, 0.5 N HCl, brine, and dried over MgSO<sub>4</sub> to afford the titled compound (410 mg) as a white solid in 90% yield. MS (ESI), ( $M+H$ )<sup>+</sup> 305.2; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  7.77 (d, 2H, *J* = 8.7), 7.62 (d, 2H, *J* = 8.7), 6.90 (br s, 1H), 3.67 (m, 1H), 1.54 (m, 1H), 1.31 (m, 2H), 0.81 (d, 3H, *J* = 7.0), 0.71 (d, 3H, *J* = 7.0).

- 54 -

Method A for conversion of III to I:5 (2R)-2-[N-(4-Chlorobenzenesulfonyl)-N-(4-methoxybenzyl)amino]-4-methylpentanoic acid amide (Example 1):


(2R)-2-(4-Chlorobenzenesulfonylamino)-4-methylpentanoic acid amide (300 mg, 1 mmol),  $K_2CO_3$  (170 mg, 1.2 mmol), and 4-methoxybenzyl chloride (170 mg, 1.1 mmol) in DMF (25 mL) was heated to 60 °C for 18 h. The reaction was then diluted with EtOAc (150 mL) and washed with  $H_2O$ , brine, dried over  $MgSO_4$ , and concentrated to give a crude white wax. Further purification by flash chromatography ( $SiO_2$ , 25% EtOAc/hexanes) afforded the titled compound (297 mg) as a white solid in 70% yield. [ $\alpha$ ] $_D^{25}$  = +44.2 (c 1.00, MeOH); MS (ESI), (M-H)<sup>-</sup> 422.9; <sup>1</sup>H NMR ( $CDCl_3$ )  $\delta$  7.63 (d, 2H, J = 7.0), 7.42 (d, 2H, J = 7.0), 7.25 (d, 2H, J = 8.0), 6.79 (d, 2H, J = 8.0), 6.25 (br s, 1H), 5.35 (br s, 1H), 4.36 (dd, 2H, J = 5.0, 1.5), 4.26 (t, 1H, J = 7.2), 3.78 (s, 3H), 1.83 (m, 1H), 1.18-1.34 (m, 2H), 0.75 (d, 3H, J = 7.0), 0.67 (d, 3H, J = 7.0); IR (KBr) 3480, 2959, 1693, 1674, 1514, 1333, 1158  $cm^{-1}$ .

20 Method B for conversion of III to I:


- 55 -

Methyl 6-dimethylaminonicotinate:

A solution of methyl 6-chloronicotinate (4.0 g, 23 mmol) in dimethylamine/MeOH (2 M, 80 mL, 160 mmol) in a pressure vessel was stirred at 95 °C for 2 h, cooled to rt and concentrated. The residue was dissolved in EtOAc (250 mL), washed with water (2 x 150 mL), dried over  $Na_2SO_4$ , and concentrated to afford the title compound as a tan solid (4.1 g, 98%). MS (ESI), (M+H)<sup>+</sup> 181.24; <sup>1</sup>H NMR ( $CDCl_3$ )  $\delta$  8.79 (s, 1H), 7.99 (d, 1H, J = 9.2), 6.45 (d, 1H, J = 9.2), 3.85 (s, 3H), 3.15 (s, 6H).

10 2-Dimethylamino-5-hydroxymethylpyridine:

A solution of methyl 6-dimethylamino-nicotinate (4.14 g, 23.0 mmol) in anhydrous ether (80 mL) at 0 °C was treated with lithium aluminum hydride (1 M in ether, 20 mL, 20 mmol). The mixture was stirred at rt for 0.5 h, cooled again to 0 °C and quenched slowly with sat. aq.  $NaHCO_3$  (10 mL). The resulting mixture was stirred at rt for 0.5 h, filtered, and washed with ether. The combined filtrates were dried over  $Na_2SO_4$ , and concentrated to give the title compound as a beige waxy solid (3.5 g, 100%). MS (ESI), (M+H)<sup>+</sup> 153.4; <sup>1</sup>H NMR ( $CDCl_3$ )  $\delta$  8.06 (d, 1H, J = 2.4), 7.47 (dd, 1H, J = 2.4, 8.8), 6.45 (d, 1H, J = 8.8), 4.50 (s, 2H), 3.06 (s, 6H), 1.98 (br s, 1H).






(2*R*)-2-[N-(4-Chlorobenzenesulfonyl)-N'-(2-dimethylamino)pyridin-5-yl]amino]-4-fluoro-4-methylpentanoic acid amide (TFA salt (Example 459):


- To a cloudy solution of (2*R*)-2-[(4-chlorobenzenesulfonylamino)-4-fluoro-4-methylpentanoic acid amide (prepared as in Reaction Scheme 20 or from  $\gamma$ -fluoro-D-Leu-OH methyl ester, Parageorgiou et. al., Bioorg. & Med. Chem. Lett. 1994, Vol. 4, p.p. 267-272; 0.060 g, 0.18 mmol), 2-dimethylamino-5-hydroxymethylpyridine (71 mg, 0.46 mmol), triphenylphosphine (122 mg, 0.464 mmol) in  $\text{CH}_2\text{Cl}_2$  (9.5 mL) at rt was added dropwise diisopropyl azodicarboxylate (75  $\mu\text{L}$ , 0.46 mmol). The resulting pale yellow solution was stirred at rt for 2 h and concentrated under vacuum. The residue was dissolved in methanol and purified by reverse phase preparative HPLC (YMAC S5, ODS, MeOH-water-TFA) to afford the title compound as a white foam (90 mg, 85%). MS (ESI), (M+H)<sup>+</sup> 457.2; <sup>1</sup>H NMR ( $\text{CDCl}_3$ )  $\delta$  8.11 (s, 1H), 7.95 (d, 1H,  $J = 9.6$ ), 7.77 (d, 2H,  $J = 6.8$ ), 7.51 (d, 2H,  $J = 6.8$ ), 6.76 (d, 2H,  $J = 9.6$ ), 6.34 (s, 1H), 6.02 (s, 1H), 4.58 (br d, 1H,  $J = 8.4$ ), 4.46 (d, 1H,  $J = 16.0$ ), 4.06 (d, 1H,  $J = 1.6$ ), 3.29 (s, 6H), 2.50 (m, 1H), 1.39 (m, 1H), 1.25 (d, 3H,  $J = 22.0$ ), 1.17 (d, 3H,  $J = 22.0$ ).

#### Exemplification of Reaction Scheme 1 – Solid Support




- Polymer-bound D-Leu-NH<sub>2</sub>: Fmoc-protected Rink amide resin (30 g, 0.61 mmol/g, 18 mmol) was treated with piperidine/DMF solution (250 mL). The mixture was shaken at rt for 24 h, drained, washed with DMF (5 x 200mL),  $\text{CH}_2\text{Cl}_2$  (5 x 200 mL) and dried under vacuum. The resin was then treated with Fmoc-D-Leu-OH (22 g, 62 mmol), 1-hydroxybenzotriazole hydrate (2.5 g, 18 mmol), 1,3-diisopropylcarbodiimide (9.8 mL, 62 mmol), and DMF (250 mL).

- The mixture was shaken for 20 h, drained, washed with DMF (4 x 200 mL), DMF-water (1:1, 3 x 200 mL), DMF (3 x 200 mL), MeOH (3 x 200 mL),  $\text{CH}_2\text{Cl}_2$  (3 x 200mL) and dried. The completion of reaction and the loading of the resin-bound Fmoc-D-Leu-NH<sub>2</sub> (0.56 mmol/g) were determined by the treatment of 52 mg of the resin with 10% (v/v) TFA/ $\text{CH}_2\text{Cl}_2$  (2 mL) to give 11 mg of Fmoc-D-Leu-NH<sub>2</sub>. The resin-bound Fmoc-D-Leu-NH<sub>2</sub> was deprotected with 20% (v/v) piperidine/DMF solution (250 mL) to give polymer-bound D-Leu-NH<sub>2</sub> (20 g).



- Polymer-bound (R)-2-(4-Chlorobenzenesulfonylamino)-4-methylpentanoic acid amide:

- The above polymer-bound D-Leu-NH<sub>2</sub> (20 g) was treated with  $\text{CH}_2\text{Cl}_2$  (150 mL), pyridine (100 mL) and 4-chlorophenylsulfonyl chloride (20.0 g, 94.8 mmol). The mixture was shaken for 24 h, drained, washed with DMF (4 x 200 mL),  $\text{CH}_2\text{Cl}_2$  (4 x 200 mL) and concentrated to give polymer-bound (R)-2-(4-chlorobenzenesulfonylamino)-4-methylpentanoic acid amide as a yellow resin (22 g). The completion of the reaction and the loading of the resin (0.57 mmol/g) were determined by the treatment of 50 mg of the resin with 10% (v/v) TFA/ $\text{CH}_2\text{Cl}_2$  (2 mL) to give 8.7 mg of (R)-2-(4-chlorobenzenesulfonylamino)-4-methylpentanoic acid amide.



- 58 -

(2*R*)-2-[*N*-(4-Chlorobenzensulfonyl)-*N*-(4-methylbenzyl)amino]-4-methylpentanoic acid amide (Example 60):


To a mixture of polymer-bound (2*R*)-2-[*N*-(4-

- 5 chlorobenzensulfonyl)amino]-4-methylpentanoic acid amide (loading 0.45 mmol/g, 50.0 mg, 0.0225 mmol), 4-methylbenzyl bromide (44 mg, 0.24 mmol) and DMF (1.5 mL) was added 2-*tert*-butylimino-2-diethylamino-1,3-dimethylperhydropyridine-1,3,2-diaza-phosphorine (0.10 mL, 0.34 mmol). The resulting mixture was shaken at rt for 2 days, then was drained and washed with DMF (4 x 2 mL), MeOH (4 x 2 mL) and CH<sub>2</sub>Cl<sub>2</sub> (4 x 2 mL).

- 10 The resin was then treated with 10% (v/v) TFA/CH<sub>2</sub>Cl<sub>2</sub>. The mixture was shaken for 1 h, filtered and washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 0.5 mL). The combined filtrates were concentrated under vacuum to afford the title compound as a beige solid (7.7 mg, 100%, HPLC purity > 95%). HRMS (ESI), (M+H)<sup>+</sup> for

- 15 C<sub>26</sub>H<sub>28</sub>SClN<sub>3</sub>O<sub>3</sub>, calcd: 407.1206, found: 407.1201; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.64 (d, 2H, *J* = 8.0), 7.44 (d, 2H, *J* = 8.0), 7.22 (d, 2H, *J* = 8.0), 7.08 (d, 2H, *J* = 8.0), 6.29 (br s, 1H), 5.34 (br s, 1H), 4.53 (d, 1H, *J* = 15.2), 4.34 (d, 1H, *J* = 15.2), 4.27 (t, 1H, *J* = 7.2), 2.32 (s, 3H), 1.84 (m, 1H), 1.30 (m, 1H), 1.21 (m, 1H), 0.75 (d, 3H, *J* = 6.8), 0.67 (d, 3H, *J* = 6.8); IR (KBr) 3467, 3367, 2956, 2869, 1694, 1670, 1340, 1160 cm<sup>-1</sup>.


Exemplification of Reaction Scheme 2



- 59 -

(2*R*)-2-(4-Methoxybenzylamino)-4-methylpentanoic acid amide:

- A solution of *D*-leucineamide hydrochloride (2.8 g, 16.8 mmol), and *p*-anisaldehyde (2.29 g, 16.8 mmol) in methanol (150 mL) was treated with anhydrous ZnCl<sub>2</sub> (338 mg, 5 mmol). The resulting suspension was then treated with NaCNBH<sub>3</sub> (1.05 g, 16.8 mmol) portion wise and heated at reflux for 3 h. The reaction was cooled to rt, quenched with saturated NaHCO<sub>3</sub> (3 mL), diluted with EtOAc (500 mL), and washed with brine. Concentration afforded the crude benzyl amine as a white wax, which was carried on without further purification (3.57 g, 84%). MS (ESI), (M+H)<sup>+</sup> 251.4; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.20 (d, 2H, *J* = 6.6), 7.10 (br s, 2H), 6.88 (d, 2H, *J* = 8.4), 5.30 (br s, 1H), 3.80 (s, 3H), 3.63 (dd, 2H, *J* = 4.5, 12), 1.44-1.65 (m, 3H), 0.95 (d, 3H, *J* = 6.3), 0.80 (d, 3H, *J* = 6.3).




- 15 (2*R*)-2-[*N*-(4-Chlorobenzensulfonyl)-*N*-(4-methoxybenzyl)amino]-4-methylpentanoic acid amide (Example 1):

- (2*R*)-2-[*N*-(4-Methoxybenzylamino)-4-methylpentanoic acid amide (3.57 g, 14.3 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and treated with Et<sub>3</sub>N (4.2 mL, 29 mmol) and 4-chlorobenzensulfonyl chloride (3.6 g, 17 mmol) at rt for 18 h. The solvents were removed and the residue was taken into EtOAc (500 mL). The organic solution was washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, and concentrated. The resulting material was then further purified by flash chromatography (SiO<sub>2</sub>, 1% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound (2.4 g) as a slightly colored solid in 40% yield. MS (ESI), (M+H)<sup>+</sup> 422.9; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.63 (d, 2H, *J* = 7.0), 7.42 (d, 2H, *J* = 7.0), 7.25 (d, 2H, *J* = 8.0), 6.79 (d, 2H, *J* = 8.0), 6.25 (br s, 1H), 5.35 (br s, 1H), 4.36 (dd, 2H, *J* = 5.0, 15), 4.26 (t, 1H, *J* =

7.2), 3.78 (s, 3H), 1.83 (m, 1H), 1.18-1.34 (m, 2H), 0.75 (d, 3H,  $J=7.0$ ), 0.67 (d, 3H,  $J=7.0$ ); IR (KBr) 3480, 2959, 1693, 1674, 1514, 1333, 1158  $\text{cm}^{-1}$ .


### Exemplification of Reaction Scheme 3



(2R)-2-[N-(4-Morpholinohexyl)-N-(4-chlorobenzenesulfonyl)amino]-4-methylpentanoic acid amide (Example 25):


- 10 A solution of (2R)-2-[N-(4-bromohexyl)-N-(4-chlorobenzenesulfonyl)amino]-4-methylpentanoic acid amide (Example 24; prepared as described in Reaction Scheme 1; 0.20 g, 0.44 mmol),  $\text{Et}_3\text{N}$  (0.25 mL, 1.7 mmol), and morpholine (150 mg, 1.7 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was stirred at rt for 18 h. The reaction was then concentrated to give a crude white wax which was purified by flash chromatography ( $\text{SiO}_2$ , 85%  $\text{EtOAc}/15\%$  hexanes/10% MeOH) to afford the title compound (112 mg) as a white solid in 54% yield. MS (ESI),  $(\text{M}+\text{H})^+$  474.4;  $^1\text{H NMR}$  ( $\text{DMSO}-d_6$ )  $\delta$  7.82 (d, 2H,  $J=8.0$ ), 7.64 (d, 2H,  $J=8.0$ ), 7.42 (br s, 1H), 6.99 (s, 1H), 4.25 (m, 1H), 3.51-3.60 (br s, 4H), 3.18-3.41 (m, 2H), 2.25-2.35 (br s, 4H), 2.27 (m, 2H), 1.5-1.62 (m, 9H), 0.80 (d, 6H,  $J=6.0$ ).

### Exemplification of Reaction Scheme 4



5 (2R)-2-[N-(4-Chlorobenzenesulfonyl)-N-(4-aminobenzyl)amino]-4-methylpentanoic acid amide (Example 48):


- 10 (2R)-2-[N-(4-Chlorobenzenesulfonyl)-N-(4-aminobenzyl)amino]-4-methylpentanoic acid amide (Compound of Example 24; prepared as described in Reaction Scheme 1; 2.8 g, 6.6 mmol) was suspended with 10% Pd/C (1 g) and conc. HCl (1 mL) in MeOH (100 mL) and placed under a hydrogen atmosphere at 40 psi for 1 h. The suspension was filtered through Celite and then concentrated to give the title compound as a tan solid (2.4 g, 88% yield). MS (ESI),  $(\text{M}+\text{H})^+$  410.1;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.80 (d, 2H,  $J=8.5$ ), 7.63 (d, 2H,  $J=8.5$ ), 7.52 (br s, 1H), 7.46 (d, 1H,  $J=8.0$ ), 7.26 (d, 1H,  $J=8.0$ ), 7.02 (br s, 1H), 4.70 (dd, 2H,  $J=5.0, 1.8$ ), 4.30-4.41 (m, 1H), 3.67 (br s, 2H), 1.28-1.33 (m, 3H), 0.86 (d, 3H,  $J=7.0$ ), 0.57 (d, 3H,  $J=7.0$ ).



20 (2R)-2-[N-(4-Chlorobenzenesulfonyl)-N-(4-methylaminobenzyl)amino]-4-methylpentanoic acid amide (Example 51):

- 62 -

A solution of (2*R*)-2-[*N*-(4-chlorobenzenesulfonyl)-*N*-(4-aminobenzyl)amino]-4-methyl-pentanoic acid amide (Example 48, 400 mg, 1 mmol), Et<sub>3</sub>N (0.16 mL, 1.1 mmol), dimethylsulfate (139 mg, 1.1 mmol) in 25 mL of toluene was stirred at rt for 18 h. The reaction was concentrated, then taken into EtOAc and washed with H<sub>2</sub>O, brine, dried over K<sub>2</sub>CO<sub>3</sub> and concentrated to give a crude mixture of starting material and product. The material was further purified by flash chromatography (SiO<sub>2</sub>, 35% EtOAc/hexanes) to afford the title compound, 195 mg, in 46% yield. MS (ESI), (M+H)<sup>+</sup> 424.1; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.65 (d, 2H, *J* = 8.0), 7.58 (d, 2H, *J* = 8.2), 7.47 (d, 2H, *J* = 8.0), 7.31 (d, 2H, *J* = 8.5), 6.24 (br s, 1H), 5.16 (br s, 1H), 4.50 (dd, 2H, *J* = 50, 17), 4.27 (t, 1H, *J* = 10), 2.44 (s, 3H), 1.74-1.83 (m, 1H), 1.25-1.33 (m, 1H), 0.93-1.01 (m, 1H), 0.74 (d, 3H, *J* = 7.0), 0.63 (d, 3H, *J* = 7.0).




(2*R*)-2-[*N*-(4-Chlorobenzenesulfonyl)-*N*-(4-dimethylaminobenzyl)amino]-4-methylpentanoic acid amide (Example 65):

(2*R*)-2-[*N*-(4-Chlorobenzene-sulfonyl)-*N*-(4-aminobenzyl)amino]-4-methyl-pentanoic acid amide (Example 48, 0.10 g, 0.22 mmol) was dissolved in DMF (5 mL). To this solution was added iodomethane (62 mg, 0.44 mmol), and cesium carbonate (220 mg, 0.66 mmol). The reaction was then stirred at 40 °C for 18 h. The reaction was poured into EtOAc and water. The organic was collected, dried over MgSO<sub>4</sub> and concentrated to an oily residue. The residue was further purified (Biotage 40S, loaded in CH<sub>2</sub>Cl<sub>2</sub>, eluted in 25% EtOAc/hexanes) to yield a yellow powder (15 mg, 16%). MS(ESI), (M+H)<sup>+</sup> 438.1; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 500 MHz) δ 7.74 (dd, 2H, *J* = 1.9, 6.7), 7.54 (dd, 2H, *J* = 1.9, 6.8), 7.43 (s, 1H),

- 63 -


7.16 (d, 2H, *J* = 8.0), 7.01 (s, 1H), 6.61 (d, 2H, *J* = 8.8), 4.59 (q, 2H, *J* = 16, 25), 4.34 (dd, 1H, *J* = 5.0, 9.3), 2.85 (s, 6H), 1.27-1.47 (m, 3H), 0.80 (d, 3H, *J* = 5.9), 0.52 (d, 3H, *J* = 6.1).

### 5 Exemplification of Reaction Scheme 5



[*N*-(1*R*)-1-Carbamoyl-3-methyl-butyl]-*N*-(4-chlorobenzenesulfonyl)amino} acetic acid tert-butyl ester (Example 40):


(2*R*)-2-(4-Chlorobenzenesulfonyl)amino)-4-methylpentanoic acid amide (3.00 g, 9.87 mmol) was dissolved in DMF (50 mL). To the solution was added potassium carbonate (6.0 g, 39 mmol) and bromoacetic acid (*tert*-butyl ester (6.0 mL, 39 mmol). The solution was heated to 70 °C for 3 h. The reaction was quenched with EtOAc and saturated NaHCO<sub>3</sub>. The organic layer was washed with brine, dried over MgSO<sub>4</sub>, and concentrated. The crude oil was further purified on a Biotage 40M (loaded in CH<sub>2</sub>Cl<sub>2</sub>, eluted in 30% EtOAc/hexanes) to afford a white powder (1.2 g, 35%). MS(ESI), (M+H)<sup>+</sup> 446.3; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.76 (d, 2H, *J* = 8.0), 7.52 (d, 2H, *J* = 8.0), 6.61 (br s, 1H), 5.45 (s, 1H), 4.15-4.18 (m, 1H), 3.09-3.24 (m, 2H), 2.50-2.58 (m, 4H), 2.31-2.39 (m, 2H), 1.92-1.99 (m, 1H), 1.15-1.59 (m, 8H), 1.00-1.04 (m, 7H), 0.71-0.74 (m, 6H).



- 64 -

[N-[(1*R*)-1-Carbamoyl-3-methyl-butyl]-N-(4-chlorobenzenesulfonyl)amino]acetic acid (Example 59):

- Trifluoroacetic acid (15 mL) was added to a solution of [N-[(1*R*)-1-carbamoyl-3-methyl-butyl]-N-(4-chlorobenzenesulfonyl)amino]acetic acid tert-butyl ester (0.50 g, 1.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 mL). The reaction was stirred at rt for 4 h. The reaction was then concentrated to a white solid (0.40 g, 92%), which was used without further purification. MS(ESI), (M+H)<sup>+</sup> 363.1; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 500MHz) δ 7.90 (dd, 2H, *J* = 2.0, 6.8), 7.65 (dd, 2H, *J* = 2.0, 6.8), 7.60 (s, 1H), 7.06 (s, 1H), 4.32 (d, 1H, *J* = 18), 4.12 (t, 1H, *J* = 8.0), 4.02 (d, 1H, *J* = 18), 1.55-1.65 (m, 1H), 1.35-1.45 (m, 2H), 0.78 (d, 3H, *J* = 6.1), 0.73 (d, 3H, *J* = 6.1).




- 15 (2*R*)-2-[N-(4-Chlorobenzenesulfonyl)-N-(cyclopropylcarbamoylmethyl)amino]-4-methyl-pentanoic acid amide (Example 88):

- To a solution of [N-[(1*R*)-1-carbamoyl-3-methyl-butyl]-N-(4-chlorobenzenesulfonyl)amino]acetic acid (Example 59, 175 mg, 0.480 mmol), cyclopropylamine (41 μL, 0.58 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was added. 1-hydroxybenzotriazole (47 mg, 0.72 mmol), and 1,3-dicyclohexylcarbodiimide (144 mg, 0.720 mmol). The reaction was stirred for 18 h at rt, and then was poured into an EtOAc/water mixture. The organic layer was separated, dried over MgSO<sub>4</sub>, and concentrated to a clear oil residue. The residue was further purified by Biogage 40S (eluted in 40% EtOAc in hexanes) to afford a white solid (54 mg, 29%). MS(ESI), (M+H)<sup>+</sup> 402.2; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500MHz) δ 7.85 (dd, 2H, *J* = 1.9, 8.9), 7.50 (dd, 2H, *J* = 2.0, 8.7), 7.40 (br s, 1H), 6.55 (br s, 1H), 6.30 (br s,

- 65 -


1H), 4.23 (dd, 1H, *J* = 2.9, 8.9), 3.92 (d, 1H, *J* = 17), 3.83 (d, 1H, *J* = 17), 2.68-2.73 (m, 1H), 1.75-1.83 (m, 1H), 1.50-1.57 (m, 1H), 1.40-1.49 (m, 1H), 0.88 (d, 3H, *J* = 6.4), 0.87 (d, 3H, *J* = 6.7), 0.80 (d, 2H, *J* = 7.0), 0.51 (t, 2H, *J* = 4.0).

5 Exemplification of Reaction Scheme 6

- 10 4-{[N-[(1*R*)-1-Carbamoyl-3-methyl-butyl]-N-(4-chlorobenzenesulfonyl)amino]-methyl}-benzoic acid (Example 89):

- A solution of the compound of Example 61 [4-{[N-[(1*R*)-1-carbamoyl-3-methyl-butyl]-N-(4-chlorobenzenesulfonyl)amino]-methyl}-benzoic acid methyl ester, 354 mg, 0.782 mmol] was dissolved in methanol (4 mL). A solution of 5 N NaOH (1 mL) was added, followed by enough THF (1 mL) to achieve homogeneity. After 1 h, an additional aliquot of 5 N NaOH (1 mL) was added, and stirring was continued for 2.5 h. The solution was acidified to pH 2 with 1 N HCl and extracted with CHCl<sub>3</sub> (2x). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give a white solid (343 mg, 100%). MS (ESI), (M+H)<sup>+</sup> 439.17; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300MHz) δ 7.91 (d, 2H, *J* = 8.2), 7.81-7.84 (m, 3H), 7.56 (d, 2H, *J* = 8.6), 7.49 (d, 2H, *J* = 8.2), 6.55 (br s, 1H), 5.10 (d, 1H, *J* = 15.4), 4.23 (dd, 1H, *J* = 4.6, 9.7), 4.05 (d, 1H, *J* = 15.4), 2.04-2.14 (m, 1H), 1.20-1.31 (m, 1H), 0.80-0.89 (m, 1H), 0.74 (d, 3H, *J* = 6.6), 0.68 (d, 3H, *J* = 6.6).

- 66 -




(2R)-2-((N-(4-chlorobenzenesulfonyl)-N'-(4-(morpholine-4-carbonyl)-benzyl)amino)-4-methyl-pentanoic acid amide (Example 101):

- 5 To a 0 °C solution of 4-((N'-(1R)-1-carbamoyl-3-methyl-butyl)-N-(4-chlorobenzenesulfonyl)amino)-methyl-benzoic acid (50.0 mg, 0.114 mmol) in DMF (0.3 mL) was added morpholine (12.9 mg, 0.148 mmol), followed by 1-hydroxybenzotriazole (18.5 mg, 0.137 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (26.2 mg, 0.137 mmol), and  $\text{Pr}_3\text{NBE}$  (26  $\mu\text{L}$ , 0.15 mmol). After 2 h, the solution was warmed to rt. After 4 h, the solution was poured into 10% aq. citric acid and extracted with EtOAc (2x). The combined organic layers were washed sequentially with water and sat. aq.  $\text{NaHCO}_3$ , then dried ( $\text{MgSO}_4$ ) and concentrated. Flash column chromatography ( $\text{SiO}_2$ , 40 to 100% EtOAc/hexanes) gave the title compound as a white solid (46.0 mg, 79%).
- 15 MS (ESI),  $(\text{M}+\text{H})^+$  508.22;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300MHz)  $\delta$  7.68 (d, 2H,  $J=8.6$ ), 7.29-7.47 (m, 6H), 6.38 (br s, 1H), 5.75 (br s, 1H), 4.65 (d, 1H,  $J=16.0$ ), 4.42 (d, 1H,  $J=16.0$ ), 4.32 (t, 1H,  $J=7.5$ ), 3.30-3.85 (br m, 8H), 1.69-1.78 (m, 1H), 1.28-1.37 (m, 1H), 1.08-1.14 (m, 1H), 0.76 (d, 3H,  $J=6.5$ ), 0.63 (d, 3H,  $J=6.6$ ).

- 67 -


#### Exemplification of Reaction Scheme 7



- 5 4-((N'-(1R)-1-carbamoyl-3-methyl-butyl)-N-(4-chlorobenzenesulfonyl)amino)-methyl-piperidine-1-carboxylic acid tert-butyl ester (Example 92):

- To a solution of (2R)-2-(4-chlorobenzenesulfonyl)amino)-4-methyl-pentanoic acid amide (4.2 grams, 14 mmol) in DMF (50 mL) was added cesium carbonate (13.6 grams, 417 mmol). To this reaction was added 4-(toluene-4-sulfonyloxy)methyl-piperidine 1-carboxylic acid tert-butyl ester (ref.: Glissen, C.; Bormans, G.; De Groot, T.; Verbruggen, A. *J. Labeled Compd. Radiopharm.* 1999, 42, 1289; 10.4 g, 282 mmol). The reaction was stirred at 70 °C for 18 h. The reaction was then quenched with sat. aq.  $\text{NaHCO}_3$  and extracted with EtOAc. The organic layer was collected, washed with brine, dried over  $\text{MgSO}_4$ , and concentrated to a clear oil. The oil was then purified on a Biotage 40S (eluted with 30% EtOAc in hexanes) to afford a white solid (3.0 g, 44%). MS(ESI),  $(\text{M}+\text{H})^+$  502.1;  $^1\text{H NMR}$  ( $\text{DMSO}-d_6$ , 500MHz)  $\delta$  7.86 (dd, 2H,  $J=2.0$ , 6.8), 7.65 (dd, 2H,  $J=2.0$ , 6.8) 7.37 (br s, 1H), 7.07 (br s, 1H), 4.19 (t, 1H,  $J=7.6$ ), 3.92 (br s, 2H), 3.35 (dd, 1H,  $J=1.5$ , 6.8), 3.05 (dd, 1H,  $J=1.5$ , 8.1), 1.85 (br s, 1H), 1.50-1.70 (m, 4H), 1.38 (s, 9H), 1.10-1.20 (m, 1H), 0.80-1.00 (m, 3H), 0.82 (d, 6H,  $J=7.6$ ).
- 20


- 68 -



(2R)-2-[(N-(4-Chlorobenzesulfonyl)-N-(piperidin-4-ylmethyl)amino)-4-methyl-pentanoic acid amide (Example 126):

To a solution of 4-[(N-((1R)-1-carbamoyl-3-methyl-butyl)-N-(4-chlorobenzesulfonyl)amino)-methyl]-piperidine-1-carboxylic acid *tert*-butyl ester (Example 92, 2.6 grams, 5.2 mmol) in  $\text{CH}_2\text{Cl}_2$  (25 mL) was added trifluoroacetic acid (10 mL). The reaction was stirred at rt for 1 h and then was concentrated to give a white solid (1.6 grams, 84%). MS(ESI), ( $M+H$ )<sup>+</sup> 402.15;

<sup>1</sup>H NMR ( $\text{DMSO}-d_6$ , 500MHz),  $\delta$  7.87 (d, 2H,  $J$  = 8.5), 7.66 (d, 2H,  $J$  = 8.6), 7.41 (s, 1H), 7.04 (s, 1H), 4.17 (t, 1H,  $J$  = 7.3), 3.40-3.50 (m, 1H), 3.20-3.25 (m, 1H), 3.03-3.10 (m, 1H), 2.65-2.80 (m, 2H), 1.85-2.00 (m, 1H), 1.20-1.85 (m, 2H), 1.45-1.60 (m, 1H), 1.30-1.40 (m, 1H), 1.10-1.30 (m, 4H), 0.75-0.90 (m, 1H), 0.82 (d, 3H,  $J$  = 7.3), 0.80 (d, 3H,  $J$  = 7.0).




(2R)-2-[(N-(4-Chlorobenzesulfonyl)-N-(1-(pyridine-4-carbonyl)-piperidin-4-ylmethyl)amino)-4-methyl-pentanoic acid amide (Example 278):

To a solution of (2R)-2-[(N-(4-chlorobenzesulfonyl)-N-(piperidin-4-ylmethyl)amino)-4-methyl-pentanoic acid amide (Example 126, 0.10 g, 0.22 mmol) and  $\text{Et}_3\text{N}$  (0.06 mL, 0.5 mmol) in  $\text{CH}_2\text{Cl}_2$  (3.0 mL) was added

- 69 -

isomonicinoyl chloride hydrochloride (56 mg, 0.32 mmol). The reaction was stirred at rt for 18 h and then was poured into a mixture of  $\text{EtOAc}$  and sat. aq.  $\text{NaHCO}_3$ . The organic solution was separated and washed with brine, dried over  $\text{MgSO}_4$ , and concentrated to an oily residue. The residue was purified on a

Biologage 10M (eluted with 80%  $\text{EtOAc}/\text{hexanes}$ ) to give a white solid (36 mg, 30%). MS(ESI), ( $M+H$ )<sup>+</sup> 509.20; <sup>1</sup>H NMR ( $\text{CDCl}_3$ , 500MHz)  $\delta$  8.66 (br s, 2H), 7.80 (d, 1H,  $J$  = 8.6), 7.73 (d, 2H,  $J$  = 8.5), 7.51 (d, 2H,  $J$  = 7.6), 7.41 (br s, 1H), 6.64 (br s, 1H), 5.35 (br s, 1H), 4.70 (br s, 1H), 4.10 (br s, 1H), 3.71 (br s, 1H), 3.33 (br s, 1H), 3.02 (dd, 2H,  $J$  = 4.8, 1.6), 2.70-2.85 (br s, 1H), 1.50-2.09 (m, 5H), 1.18-1.33 (m, 4H), 0.73 (d, 3H,  $J$  = 6.7), 0.68 (d, 3H,  $J$  = 6.5).



4-[(N-((1R)-1-Carbamoyl-3-methyl-butyl)-N-(4-chlorobenzesulfonyl)amino)-methyl]-piperidine-1-carboxylic acid phenethylamide (Example 256):

To a solution of (2R)-2-[(N-(4-chlorobenzesulfonyl)-N-(piperidin-4-ylmethyl)amino)-4-methyl-pentanoic acid amide (Example 126, 0.10 g, 0.22 mmol) and  $\text{Et}_3\text{N}$  (32  $\mu\text{L}$ , 0.25 mmol) in  $\text{CH}_2\text{Cl}_2$  (3.0 mL) was added (2-isocyanato-ethyl)-benzene (0.040 mL, 0.30 mmol). The reaction was stirred at rt


for 18 h and then was poured into sat. aq.  $\text{NaHCO}_3$  and extracted with  $\text{EtOAc}$ .

The organic layer was washed with brine, dried over  $\text{MgSO}_4$ , and concentrated to an oily residue. The residue was further purified on a Biologage system (eluted with 75%  $\text{EtOAc}/\text{hexanes}$ ) to afford the desired product as a white solid (67 mg, 52%). MS(ESI), ( $M+H$ )<sup>+</sup> 549.00; <sup>1</sup>H NMR ( $\text{CDCl}_3$ , 500MHz)  $\delta$  7.71 (d, 2H,  $J$  = 8.6), 7.71 (d, 2H,  $J$  = 8.9), 7.15-7.35 (m, 5H), 6.64 (s, 1H), 5.86 (s, 1H), 4.15 (dd,

- 70 -

1H,  $J = 5.2, 9.5$ ), 3.88 (d, 1H,  $J = 13$ ), 3.76 (d, 1H,  $J = 13$ ), 3.46 (t, 2H,  $J = 6.7$ ), 3.21-3.29 (m, 1H), 2.97 (dd, 1H,  $J = 4.6, 14$ ), 2.65-2.85 (m, 4H), 1.75-1.95 (m, 3H), 1.00-1.30 (m, 5H), 0.75-0.80 (m, 1H), 0.72 (d, 3H,  $J = 6.7$ ), 0.67 (d, 3H,  $J = 6.7$ ).

5



(2R)-2-[(N-(4-Chlorobenzesulfonyl)-N'-(1-(2-(4-cyanophenyl)-2-oxo-ethyl)-piperidin-4-ylmethyl)-amino)-4-methyl-pentanoic acid amide (Example 286):

10

To a solution of (2R)-2-[(N-(4-chlorobenzesulfonyl)-N'-(piperidin-4-ylmethyl)amino)-4-methyl-pentanoic acid amide (Example 126, 0.050 g, 0.12 mmol) and Et<sub>3</sub>N (0.040 mL, 0.30 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL) was added 4-(2-chloro-acetyl)-benzonitrile (55 mg, 0.30 mmol). The reaction was stirred at rt for

18 h and then was concentrated to residue. The residue was purified on a Biotage


15

system (eluted with 80% EtOAc/hexanes) to produce 29 mg (48%) of the desired product as a white solid. MS(ESI), (M+H)<sup>+</sup> 545.16; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500MHz) δ 7.72 (d, 2H,  $J = 8.5$ ), 7.50-7.65 (m, 2H), 7.50 (d, 2H,  $J = 7.0$ ), 7.35-7.45 (m, 2H), 6.67 (s, 1H), 5.32 (s, 1H), 4.14 (dd, 1H,  $J = 5.0, 9.0$ ), 3.52 (br s, 1H), 3.28 (t, 1H,  $J = 14$ ), 2.97 (dd, 1H,  $J = 3.5, 14$ ), 2.82 (br s, 1H), 1.00-2.00 (m, 10H), 0.71 (d, 3H,  $J = 6.5$ ), 0.66 (d, 3H,  $J = 6.5$ ).

20

- 71 -

### Exemplification of Reaction Scheme 8



5

(2R)-2-[(N-(4-Chlorobenzesulfonyl)-N'-(4-(tetrahydro-pyran-2-yloxy)methyl)-benzyl)-amino)-4-methyl-pentanoic acid amide:

(2R)-2-(4-Chlorobenzesulfonylamino)-4-methyl-pentanoic acid amide

(6.35 g, 196 mmol), Cs<sub>2</sub>CO<sub>3</sub> (5.62 g, 196 mmol), and 2-[(4-

bromomethyl)benzyl]oxy]tetrahydropyran (5.62 g, 196 mmol) in acetonitrile (200

10

mL) were heated to reflux for 1 h. The reaction was filtered hot with suction through Celite. The filtrate was reduced *in vacuo* to a white foam (9.5 g, 96%).

The foam was used as is in the next reaction. MS (ESI), (M+H)<sup>+</sup> 510.9, <sup>1</sup>H NMR

(CDCl<sub>3</sub>) δ 7.83 (d, 2H,  $J = 8.0$ ), 7.75 (d, 2H,  $J = 8.0$ ), 7.39 (d, 2H,  $J = 8.0$ ), 7.24


(d, 2H,  $J = 8.0$ ), 6.25 (br s, 1H), 5.35 (br s, 1H), 4.82 (d, 1H,  $J_{ab} = 12$ ), 4.65 (m,

1H), 4.52 (d, 1H,  $J_{ab} = 12$ ), 4.30 (d, 1H,  $J_{ab} = 16$ ), 4.20 (d, 1H,  $J_{ab} = 16$ ), 3.74 (m,

2H), 3.46 (m, 1H), 1.89 (m, 1H), 1.66 (m, 6H), 0.97 (d, 3H,  $J = 7.0$ ), 0.94 (d, 3H,

$J = 7.0$ ).

15



20




- 72 -

(2*R*)-2-[N-(4-chlorobenzenesulfonyl)-N-(4-hydroxymethyl)benzylamino]-4-methyl-pentanoic acid amide (Example 95):

To a solution of (2*R*)-2-[N-(4-chlorobenzenesulfonyl)-N-(4-

(tetrahydropyran-2-yl)oxymethyl)benzylamino]-4-methyl-pentanoic acid amide (9.5 g, 186  $\mu$ mol) in methanol (200 mL) was added a catalytic amount of *p*-toluenesulfonic acid. The mixture was stirred overnight at rt. The solvent was removed *in vacuo*. The resulting foam was dissolved in  $\text{CH}_2\text{Cl}_2$  (100 mL) washed with 1 N NaOH,  $\text{H}_2\text{O}$ , brine, and dried over  $\text{MgSO}_4$ . The filtrate solvent was removed *in vacuo*. The resulting foam was crystallized from hot hexane affording the product as a white solid (7.7g) in 92 % yield. MS (ESI), ( $\text{M}+\text{H}^+$ ) 425.17,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.68 (d, 2H,  $J=7.0$ ), 7.46 (d, 2H,  $J=7.0$ ), 7.33 (d, 2H,  $J=8.0$ ), 7.28 (d, 2H,  $J=8.0$ ), 6.26 (br s, 1H), 5.35 (br s, 1H), 4.67 (br s, 2H), 4.59 (d, 1H,  $J_{\text{ab}}=16$ ), 4.37 (d, 1H,  $J_{\text{ab}}=16$ ), 4.26 (t, 1H, 7.0), 1.86-1.80 (m, 2H), 1.34-1.28 (m, 1H), 1.16-1.10 (m, 1H), 0.96 (d, 3H,  $J=7.0$ ), 0.93 (d, 3H,  $J=7.0$ ).




Methanesulfonic acid 4-[(N-((1*R*)-1-carbamoyl-3-methyl-butyl)-N-(4-chlorobenzene-sulfonyl)-amino)methyl]benzyl ester:

To a stirred solution of (2*R*)-2-[N-(4-chloro-benzenesulfonyl)-N-(4-hydroxymethyl-benzyl)amino]-4-methyl-pentanoic acid amide (1.5 g, 3.5  $\mu$ mol) in  $\text{CH}_2\text{Cl}_2$  (15 mL) cooled to 0 °C was added  $\text{Et}_3\text{N}$  (0.74 mL, 5.3  $\mu$ mol). A solution of methanesulfonyl chloride (0.29 mL, 3.5  $\mu$ mol) in 5 mL  $\text{CH}_2\text{Cl}_2$  was added dropwise and the reaction was allowed to stir at 0 °C for 1 h. The reaction mixture was diluted with 25 mL  $\text{CH}_2\text{Cl}_2$ , quickly washed with 1 N HCl, brine, and dried by passing the organic phase through a cotton plug. The solvent was

- 73 -


removed *in vacuo* affording the title compound in quantitative yield. The resulting foam was used as is in subsequent reactions. MS (ESI), ( $\text{M}+\text{H}^+$ ) 409.15  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.70 (d, 2H,  $J=8.0$ ), 7.48 (d, 2H,  $J=8.0$ ), 7.41 (d, 2H,  $J=8.0$ ), 7.38 (d, 2H,  $J=8.0$ ), 6.27 (br s, 1H), 5.32 (br s, 1H), 5.24 (s, 2H), 4.64 (d, 1H,  $J_{\text{ab}}=16$ ), 4.43 (d, 1H,  $J_{\text{ab}}=16$ ), 4.33 (t, 1H,  $J=6$ ), 2.90 (s, 3H), 1.90 (m, 1H), 1.60 (m, 2H), 0.96 (d, 3H,  $J=7.0$ ), 0.91 (d, 3H,  $J=7.0$ )



(2*R*)-2-[N-(4-chlorobenzenesulfonyl)-N-(4-dimethylaminomethyl)benzyl]amino]-4-methyl-pentanoic acid amide (Example 110):

To a stirred solution of methanesulfonic acid 4-[(N-((1*R*)-1-carbamoyl-3-methyl-butyl)-N-(4-chlorobenzenesulfonyl)amino)-methyl]-benzyl ester (150 mg, 0.298  $\mu$ mol) in (3 mL)  $\text{CH}_2\text{Cl}_2$  at 0 °C was added 1 equivalent of  $\text{Et}_3\text{N}$ , followed by dimethylamine (0.3 mL, 2 M in THF). The reaction was stirred overnight at rt. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{MgSO}_4$ , and concentrated to give an amber glass. Purification by flash chromatography ( $\text{SiO}_2$ , 10%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) afforded the title compound (95 mg) in 71 % yield. MS (ESI), ( $\text{M}+\text{H}^+$ ) 452.23,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.94 (d, 2H,  $J=8.0$ ), 7.74 (d, 2H,  $J=8.0$ ), 7.63 (d, 2H,  $J=8.0$ ), 7.38 (d, 2H,  $J=8.0$ ), 6.23 (br s, 1H), 5.35 (br s, 1H), 4.22 (d, 1H,  $J_{\text{ab}}=16$ ), 4.14 (d, 1H,  $J_{\text{ab}}=16$ ), 3.28-3.23 (m, 3H), 2.17 (br s, 6H), 1.95 (m, 1H), 1.55 (m, 2H), 0.96 (d, 3H,  $J=7.0$ ), 0.93 (d, 3H,  $J=7.0$ ).


- 74 -

Exemplification of Reaction Scheme 9

5 (2R)-2-[N-(4-Acetylamino-benzyl)-N-(4-chlorobenzensulfonyl)amino]-4-methyl-pentanoic acid amide (Example 163):

A solution of the compound of Example 48 [(2R)-2-[N-(4-chlorobenzensulfonyl)-N-(4-aminobenzyl)amino]-4-methyl-pentanoic acid amide (250 mg, 0.60 mmol) and Et<sub>3</sub>N (120 mg, 1.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was treated with acetyl chloride (56 mg, 0.72 mmol). After stirring for 18 h, the reaction was concentrated, chromatographed using silica gel flash


15 chromatography (1% methanol/CH<sub>2</sub>Cl<sub>2</sub>) to afford the titled compound (110 mg, 41%). MS (ESI), (M-H)<sup>-</sup> 422.9; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.67 (d, 2H, J = 8.0), 7.28-7.46 (m, 6H), 7.12 (br s, 1H), 6.24 (br s, 1H), 5.19 (br s, 1H), 4.48 (dd, 2H, J = 5.0, 15), 4.27 (t, 1H, J = 7.0), 2.18 (s, 3H), 1.80-2.01 (m, 1H), 1.12-1.32 (m, 2H), 0.75 (d, 3H, J = 7.0), 0.67 (d, 3H, J = 7.0).



- 75 -

5 (2R)-2-[N-(4-Chlorobenzensulfonyl)-N-(4-[(2-dimethylamino-acetyl)-methyl-amino]-methyl)-benzyl)-amino]-4-methyl-pentanoic acid amide (Example 272):

(2R)-2-[N-(4-Chlorobenzensulfonyl)-N-(4-methylamino-methyl)-benzyl)-amino]-4-methyl-pentanoic acid amide (75 mg, 0.17 mmol), (α-dimethylamino)acetic acid (18 mg, 0.17 mmol), 1-hydroxybenzotriazole (24 mg, 0.17 mmol), and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (33 mg, 0.17 mmol) were combined in 3 mL CH<sub>2</sub>Cl<sub>2</sub> and stirred overnight. The reaction mixture was diluted with 5 mL CH<sub>2</sub>Cl<sub>2</sub> and washed with 1 N NaOH and brine. The organic phase was dried by filtering through cotton and the solvent was removed *in vacuo*. Purification via preparative HPLC afforded the title compound (61 mg) in 68% yield. MS (ESI), 523.4 (M+H)<sup>+</sup> <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.02 (d, 2H, J = 8.0), 7.71 (d, 2H, J = 8.0), 7.37 (d, 2H, J = 8.0), 7.28 (d, 2H, J = 8.0), 6.23 (br s, 1H), 5.51 (br s, 1H), 4.46 (s, 2H), 4.70 (d, 1H, J<sub>ab</sub> = 16), 4.33 (d, 1H, J<sub>ab</sub> = 16), 3.25 (t, 1H, J = 6.0), 2.69 (s, 3H), 2.63 (s, 2H), 2.20 (s, 6H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J = 7.0), 0.94 (d, 3H, J = 7.0).

Exemplification of Reaction Scheme 10


20 (2R)-2-[N-(4-Chlorobenzensulfonyl)-N-(2-dimethylamino-pyridin-5-ylmethyl)amino]-4-methyl-pentanoic acid amide TFA salt (Example 254):

A solution of (2R)-2-[N-(4-chlorobenzensulfonyl)-N-(2-chloropyridin-5-ylmethyl)amino]-4-methyl-pentanoic acid amide (prepared via Reaction Scheme 1, 18 mg, 41 μmol) in dimethylamine/THF (2 M, 20 mL, 40 μmol) was stirred at 95 °C for 30 h in a pressure vessel. Five mL of reaction mixture (25% of total

- 76 -

reaction volume) was purified by reverse phase preparative HPLC (YMC S5, ODS, MeOH-water-TFA) to afford the title compound as a white foam (17 mg, 30% yield). HRMS (ESI, (M-H)<sup>+</sup> for C<sub>26</sub>H<sub>25</sub>SClN<sub>2</sub>O<sub>3</sub> calcd: 437.1426, found: 437.1420; <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.04 (s, 1H), 8.03 (d, 1H, J = 9.8), 7.76 (d, 2H, J = 7.6), 7.54 (d, 2H, J = 7.6), 6.83 (d, 1H, J = 9.8), 6.62 (br s, 1H), 6.40 (br s, 1H), 4.64 (d, 1H, J = 15.9), 4.29 (m, 1H), 4.18 (d, 1H, J = 15.9), 3.30 (s, 6H), 1.84 (m, 1H), 1.29 (m, 1H), 0.93 (m, 1H), 0.77 (d, 3H, J = 6.5), 0.72 (d, 3H, J = 6.5).

### Exemplification of Reaction Scheme 11




(2R)-2-[N-(4-allyloxy-3-fluorobenzyl)-N-(4-chlorobenzesulfonyl)amino]-4-methyl-pentanoic acid amide:

To a solution of (2R)-2-(4-chlorobenzesulfonyl)amino)-4-methyl-pentanoic acid amide (1.00 g, 3.29 mmol), and Cs<sub>2</sub>CO<sub>3</sub> (1.29 g, 3.95 mmol) in DMF (25 mL) was added 1-allyloxy-4-bromomethyl-2-fluorobenzene (ref.: Graham, Samuel L, et al., Eur. Pat. Appl. (1992): EP 487270; 0.88 g, 3.67 mmol). The resulting solution was stirred at rt for 18 h. The reaction was then diluted with 9:1 EtOAc:hexanes (350 mL) and washed with H<sub>2</sub>O (4 x 200 mL), brine, and dried over Na<sub>2</sub>SO<sub>4</sub> to afford the titled compound (393 mg) as a white solid in 26% yield. MS (ESI), (M+H)<sup>+</sup> 469.1; <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.66 (d, 2H, J = 8.1), 7.45 (d, 2H, J = 8.1), 7.11 (d, 1H, J = 12.0), 6.98 (m, 1H), 6.84 (t, 1H, J = 8.0), 6.22 (br s, 1H), 6.04 (m, 2H), 5.42 (m, 1H), 5.16 (br s, 1H), 4.59 (m, 2H), 4.40 (m, 3H), 1.83 (m, 1H), 1.32 (m, 1H), 1.14 (m, 1H), 0.76 (d, 3H, J = 7.0), 0.68 (d, 3H, J = 7.0).

- 77 -

(2R)-2-[N-(4-Chlorobenzesulfonyl)-N-[3-fluoro-4-(2-morpholin-4-yl-ethoxy)-benzyl]-amino)-4-methyl-pentanoic acid amide (Example 42):




A mixture of the allyloxy intermediate (0.39 g, 0.84 mmol) from above, osmium tetroxide (0.01 g, 0.04 mmol), and trimethylamine N-oxide (0.140 g, 1.81 mmol) was dissolved in acetone (10 mL) and stirred for 4 h at rt. The solution was concentrated *in vacuo* and redissolved in 1.5:1 dioxane:H<sub>2</sub>O (15 mL). Sodium periodate (0.22 g, 1.0 mmol) was added and the solution was stirred at rt for 18 h. The reaction was then diluted with EtOAc (200 mL) and washed with H<sub>2</sub>O, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give (2R)-[N-(4-chlorobenzesulfonyl)-N-[3-fluoro-4-(2-oxo-ethoxy)-benzyl]-amino)-4-methyl-pentanoic acid amide as a crude beige solid. This crude material was taken onto the next step without further purification. (2R)-2-[N-(4-Chlorobenzesulfonyl)-N-[3-fluoro-4-(2-oxo-ethoxy)-benzyl]-amino)-4-methyl-pentanoic acid amide (0.16 g, 0.34 mmol) and morpholine (0.090 g, 1.0 mmol) was dissolved in EtOH (5 mL) and heated to 80 °C for approximately 15 min. The oil bath was removed and sodium triacetoxyborohydride (0.290 g, 1.36 mmol) was added and the slurry was stirred at rt for 16h. The solution was concentrated to dryness, taken up in brine, extracted with EtOAc (2 x 100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo* to give a crude orange residue. Further purification by Prep HPLC (20 x 100mm YMC S5 ODS C-18 column, 25 mL/min, 0-100 % MeOH/H<sub>2</sub>O 0.1 % TFA 15 min) afforded as a TFA salt the titled compound (69.5 mg) as a pale yellow solid in 31 % yield. [α]<sub>D</sub><sup>23</sup> +23 (c 6.4, CH<sub>2</sub>Cl<sub>2</sub>); LCMS (M+H)<sup>+</sup> 542.25; <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.71 (d, 2H, J = 8.0), 7.50 (d, 2H, J = 8.0), 7.16 (d, 1H, J =

- 78 -

12.0), 7.05 (d, 1H,  $J=8.0$ ), 6.87 (t, 1H,  $J=8.0$ ), 6.38 (br s, 1H), 5.91 (br s, 1H), 4.41 (ABq, 2H,  $J=16$ ,  $J_w=17.6$ ), 4.45 (m, 2H), 4.27 (t, 1H,  $J=8.0$ ), 4.03 (m, 4H), 3.70 (m, 2H), 3.51 (m, 2H), 3.10 (m, 2H), 1.83 (m, 1H), 1.29 (m, 1H), 1.05 (m, 1H), 0.75 (d, 3H,  $J=8.0$ ), 0.68 (d, 3H,  $J=8.0$ ).

5

### Exemplification of Reaction Scheme 12




10 (2R)-2-{N-(4-Chlorobenzesulfonyl)-N-[4-(1-hydroxy-1-methyl-ethyl)-benzyl]-amino}-4-methyl-pentanoic acid amide (Example 287):

A solution of the compound of Example 61 [4-{N-[(1R)-1-carbamoyl-3-methyl-butyl]-N-(4-chlorobenzesulfonyl)amino]-methyl]-benzoic acid methyl ester, 101 mg, 0.221 mmol] was cooled to 0 °C in THF (2 mL). A solution of methyl magnesium bromide (1.4 M in toluene/THF, 0.50 mL, 0.71 mmol) was added dropwise. The dark yellow solution was stirred at 0 °C, and after 30 min, additional methyl magnesium bromide solution (0.25 mL, 0.353 mmol) was added. After 1 h, the solution was allowed to warm to rt. After 3.5 h, the reaction was quenched by the addition of sat. aq.  $\text{NH}_4\text{Cl}$ , and the mixture was extracted with EtOAc (2x). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. Flash column chromatography ( $\text{SiO}_2$ , 20 to 100% EtOAc/hexanes) provided the title compound as a white foam (62 mg, 62%). MS (ESI,  $(\text{M}+\text{H})^+$  453.16;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300MHz)  $\delta$  7.61 (d, 2H,  $J=8.7$ ), 7.40 (d, 2H,  $J=8.7$ ), 7.37 (d, 2H,  $J=8.4$ ), 7.26 (d, 2H,  $J=8.4$ ), 6.28 (br s, 1H), 5.25 (br s, 1H), 4.49 (d, 1H,  $J=15.9$ ), 4.41 (d, 1H,  $J=15.9$ ), 4.33 (t, 1H,  $J=6.6$ ), 1.73-1.80 (m, 1H),

- 79 -

1.55 (s, 6H), 1.28-1.35 (m, 1H), 1.20-1.25 (m, 1H), 0.77 (d, 3H,  $J=6.5$ ), 0.66 (d, 3H,  $J=6.6$ ).

### Exemplification of Reaction Scheme 13



5

(2R)-2-{N-(4-Chlorobenzesulfonyl)-[4-(5-methyl-1,3,4-oxadiazol-2-yl)-benzyl]-amino}-4-methyl-pentanoic acid amide (Example 436):


10 Step 1: A solution of the compound of Example 61 [4-{N-[(1R)-1-carbamoyl-3-methyl-butyl]-N-(4-chlorobenzesulfonyl)amino]-methyl]-benzoic acid methyl ester, 0.500 g, 1.10 mmol] was diluted with methanol (10 mL) and hydrazine (2 mL) was added. The starting material slowly dissolved over 5 min. After 30 min, the solution was heated at reflux. After 22 h, the solution was cooled to rt. Water (15 mL) was added, and a white precipitate formed. The mixture was extracted with EtOAc (2x). The combined organic layers were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to give the corresponding acyl hydrazide as a white foam, which was carried directly on to the cyclization step without purification.

20 Step 2: The crude acyl hydrazide (0.150 g, 0.331 mmol) was dissolved in pyridine (2.2 mL) and ethyl acetoimidate hydrochloride (60.0 mg, 0.364 mmol) was added. The mixture was heated at reflux for 1.25 h. The solution was cooled to rt and concentrated to remove pyridine. The residue was taken up in EtOAc, and was washed sequentially with water, 1 N HCl (2x), sat. aq.  $\text{NaHCO}_3$ , and brine. The solution was dried ( $\text{MgSO}_4$ ) and concentrated. Flash column chromatography ( $\text{SiO}_2$ , 50 to 100% EtOAc/hexanes) provided the listed

- 80 -

compound as a white solid (138 mg, 88% for 2 steps).  $[\alpha]_D^{25} +11.1$  (c 7.0 mg/mL,  $\text{CHCl}_3$ ); MS (ESI):  $(M+H)^+$  477.22;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300MHz)  $\delta$  7.94 (dd, 2H,  $J = 1.8, 8.4$ ), 7.69 (dd, 2H,  $J = 1.8, 8.7$ ), 7.45-7.50 (m, 4H), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.65 (d, 1H,  $J = 15.9$ ), 4.46 (d, 1H,  $J = 15.9$ ), 4.31 (dd, 1H,  $J = 6.6, 7.8$ ), 2.61 (s, 3H), 1.75-1.85 (m, 1H), 1.28-1.35 (m, 1H), 1.08-1.15 (m, 1H), 0.76 (d, 3H,  $J = 6.6$ ), 0.64 (d, 3H,  $J = 6.6$ ).

#### Exemplification of Reaction Scheme 14




(2*R*)-2-(*N*-(4-Chlorobenzenesulfonyl)-*N*-(4-(3-methyl-1,2,4-oxadiazol-5-yl)-benzyl)-amino)-4-methyl-pentanoic acid amide (Example 437):

**Step 1:** To a rt solution of the compound of Example 89 [4-((*N*-(1*R*)-carbamoyl-3-methyl-butyl)-*N*-(4-chlorobenzene-sulfonyl)-amino)-methyl]-benzoic acid, 520 mg, 1.2 mmol] in DMF (2.4 mL) and  $\text{CH}_2\text{Cl}_2$  (7.1 mL) was added 1-hydroxybenzotriazole (192 mg, 1.42 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (272 mg, 1.42 mmol), and  $\text{Pr}_3\text{NEt}$  (0.31 mL, 1.8 mmol). *N*-Hydroxyacetamide (105 mg, 1.42 mmol) was also added. After 21 h, starting material was evident, so additional portions of all reagents were added periodically to push the reaction forward. After 3 d, the mixture was concentrated and partitioned between sat. aq.  $\text{NaHCO}_3$  and EtOAc (2x). The combined organic layers were washed with brine, dried ( $\text{MgSO}_4$ ), and concentrated to a yellow oil, which was carried on to the next step without purification.

- 81 -

**Step 2:** The crude acetamidoxime was dissolved in toluene (10 mL) and the solution was heated at reflux. After 1 h, pyridine (2 mL) was added and heating was continued for another 15 h. The mixture was concentrated and diluted with EtOAc. The organic phase was washed sequentially with water, 1 N HCl (2x), sat. aq.  $\text{NaHCO}_3$ , and brine, then was dried ( $\text{MgSO}_4$ ) and concentrated. Flash column chromatography ( $\text{SiO}_2$ , 10 to 40% EtOAc/hexanes) gave the title compound as a pale yellow solid (238 mg, 42% for two steps).  $[\alpha]_D^{25} +9.30$  (c 5.93,  $\text{CHCl}_3$ ); MS (ESI):  $(M+H)^+$  477.18;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300MHz)  $\delta$  8.04 (d, 2H,  $J = 8.4$ ), 7.70 (dd, 2H,  $J = 1.8, 8.4$ ), 7.45-7.52 (m, 4H), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.67 (d, 1H,  $J = 16.2$ ), 4.47 (d, 1H,  $J = 15.9$ ), 4.31 (t, 1H,  $J = 7.2$ ), 2.47 (s, 3H), 1.75-1.85 (m, 1H), 1.28-1.35 (m, 1H), 1.08-1.15 (m, 1H), 0.76 (d, 3H,  $J = 6.6$ ), 0.64 (d, 3H,  $J = 6.6$ ).

#### Exemplification of Reaction Scheme 15




(2*R*)-2-(*N*-(4-Chlorobenzenesulfonyl)-*N*-(4-(5-methyl-1,2,4-oxadiazol-3-yl)-benzyl)-amino)-4-methyl-pentanoic acid amide (Example 465):

**A solution of the compound of Example 6** [(2*R*)-2-(*N*-(4-chlorobenzenesulfonyl)-*N*-(4-cyanobenzyl)amino)-4-methyl-pentanoic acid amide (0.20 g, 0.47 mmol) in ethanol (6 mL) was treated with hydroxylamine (50% solution in water, 0.050 mL, 0.71 mmol). The reaction was heated to 80 °C for 18 h. The reaction was concentrated to a residue and recrystallized from EtOAc/hexanes to produce a white solid (136 mg, 51%). This solid (0.18 mmol)

- was then dissolved in chloroform and treated with  $\text{Et}_3\text{N}$  (0.030 mL, 0.24 mmol) and acetyl chloride (0.020 mL, 0.18 mmol). The reaction was stirred at rt for 2 h and then was poured into EtOAc and brine. The organic layer was separated, dried over  $\text{MgSO}_4$ , and concentrated to residue. The residue was taken up in toluene and heated at reflux for 24 h. The reaction was concentrated to a residue and purified on a Biotage system (eluted in 1:1 EtOAc/hexanes) to afford the desired product as a white solid (35 mg, 39% yield). MS(ESI),  $(\text{M}+\text{H})^+$  477.13;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 500MHz)  $\delta$  7.98 (d, 2H,  $J=8.2$ ), 7.68 (d, 2H,  $J=8.9$ ), 7.45 (d, 4H,  $J=8.5$ ), 6.21 (s, 1H), 5.19 (s, 1H), 4.62 (d, 1H,  $J=1.5$ ), 4.48 (d, 1H,  $J=1.6$ ), 4.31 (t, 1H,  $J=7.0$ ), 2.65 (s, 3H), 1.75-1.85 (m, 1H), 1.20-1.35 (m, 4H), 1.10-1.17 (m, 1H), 0.85-0.90 (m, 1H), 0.75 (d, 3H,  $J=6.7$ ), 0.64 (d, 3H,  $J=6.4$ ).

#### Exemplification of Reaction Scheme 16




(2*R*)-2-[*N*-(4-Acetylbenzyl)-*N'*-(4-chlorobenzenesulfonyl)amino]-4-methyl-pentanoic acid amide (Example 273):

- A solution of the compound of Example 251 [4-[(*N*-(1*S*)-1-carbamoyl-3-methyl-butyl)-*N'*-(4-chlorobenzenesulfonyl)-amino]-methyl]-*N'*-methoxy-*N'*-methyl-benzamide, 0.100 g, 0.207 mmol] was cooled to 0 °C in THF (2.1 mL). A solution of methyl magnesium bromide (1.4 M in toluene/THF, 0.178 mL, 0.249 mmol) was added dropwise. The resulting solution was stirred at 0 °C for 3 h, at which time additional methyl magnesium bromide solution (0.178 mL, 0.249 mmol) was added. After another 30 min, a final portion of MeMgBr solution (0.3 mL) was added. After a final 15 min, the reaction was quenched by the addition

- of sat. aq.  $\text{NH}_4\text{Cl}$  and 1 N HCl, and the mixture was extracted with EtOAc (2x). The combined organic layers were washed with sat. aq.  $\text{NaHCO}_3$  and brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. Flash column chromatography ( $\text{SiO}_2$ , 20 to 60% EtOAc/hexanes) gave the desired compound as an off-white foam (79 mg, 87%).  $[\alpha]_D^{20} +20.4$  (c 7.57,  $\text{CHCl}_3$ ); MS (ESI),  $(\text{M}+\text{H})^+$  437.13;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300MHz)  $\delta$  7.87 (d, 2H,  $J=8.4$ ), 7.67 (dd, 2H,  $J=1.8, 8.7$ ), 7.42-7.46 (m, 4H), 6.21 (br s, 1H), 5.28 (br s, 1H), 4.64 (d, 1H,  $J=1.5$ ), 4.45 (d, 1H,  $J=1.5$ ), 4.31 (t, 1H,  $J=6.6$ ), 2.58 (s, 3H), 1.73-1.80 (m, 1H), 1.25-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.74 (d, 3H,  $J=6.5$ ), 0.65 (d, 3H,  $J=6.6$ ).

#### Exemplification of Reaction Scheme 17




(2*R*)-2-[*N'*-(4-Chlorobenzenesulfonyl)-*N*-(4-(3-piperidin-1-yl-propionyl)amino)-benzyl]-amino]-4-methyl-pentanoic acid amide (Example 274):

- To a solution of *N'*-(4-[(*N*-(1*S*)-1-carbamoyl-3-methyl-butyl)-*N'*-(4-chlorobenzenesulfonyl)amino]-methyl)-phenyl]-acrylamide (0.10 g, 0.22 mmol) in toluene (5 mL) was added piperidine (20 mg, 0.24 mmol). The mixture was heated at a gentle reflux for 1 h and then the solvent was removed *in vacuo*. Purification by flash chromatography ( $\text{SiO}_2$ , 10%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) afforded the title compound (105 mg) in 86% yield. MS (ESI),  $(\text{M}+\text{H})^+$  449.16,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.69 (d, 2H,  $J=8.0$ ), 7.63 (d, 2H,  $J=8.0$ ), 7.38 (d, 2H,  $J=8.0$ ), 7.23 (d, 2H,  $J=8.0$ ), 6.25 (br s, 1H), 5.35 (br s, 1H), 4.75 (d, 1H,  $J_s=1.6$ ),

4.38 (d, 1H,  $J_{\text{AB}} = 16$ ), 3.25 (t, 1H,  $J = 6.0$ ), 2.65 (t, 2H,  $J = 6.0$ ), 2.56-2.44 (m, 6H), 1.95 (m, 1H), 1.68-1.45 (m, 8H), 0.98 (d, 3H,  $J = 7.0$ ), 0.94 (d, 3H,  $J = 7.0$ )

### Exemplification of Reaction Scheme 18




(2R)-2-(benzhydrylidene-amino)-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)-4-fluorobutan-1-one:

10 To a  $-78^{\circ}\text{C}$  solution of N-2-(benzhydrylidene-amino)-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)ethanone (ref: Josien, H.; Martin, A.; Chassaign, G. Tetrahedron Lett. 1991, 32, 6547; 30.0g, 68 mmol) in HMMPA (60 mL) and THF (300 mL) was added n-BuLi (1.6 M in hexane, 42.4 mL, 68 mmol) dropwise, maintaining the

15 temperature below  $-65^{\circ}\text{C}$ . The reaction was allowed to come to rt, at which time a solution of 1-bromo-3-fluoroethane (17.4 g, 137 mmol) in THF (30 mL) was added dropwise at rt. After 18 h the reaction was poured over  $\text{H}_2\text{O}/\text{HOAc}$  (200 mL/2 mL), diluted with EtOAc, and the organic layers were washed with saturated  $\text{NH}_4\text{Cl}$ , brine, dried over  $\text{MgSO}_4$ , and concentrated. The resulting orange oil was then further purified by silica gel chromatography (25% EtOAc/hexanes) to afford a white solid which was recrystallized from 15% EtOAc/hexanes to give the desired material (24.3 g, 70%). MS (ESI) (M + H<sup>+</sup>) 483.27;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.66 (d, 2H,  $J = 7.2$ ), 7.13-7.44 (m, 8H), 4.82-4.83 (m, 2H), 4.39-4.81 (m, 2H), 3.84-3.87 (m, 1H), 3.28 (ABq, 2H,  $J = 18$ , 10) 2.33-2.41 (m, 2H), 2.02-2.04 (m, 2H), 1.84-1.87 (m, 2H), 1.32-1.39 (m, 2H), 1.10 (s, 3H), 0.91 (s, 3H).


25



(2R)-2-Amino-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)-4-fluorobutan-1-one:

5 A solution of (2R)-2-(benzhydrylidene-amino)-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)-4-fluorobutan-1-one (20.0 g, 41.0 mmol) in THF (400 mL) was treated with 1 N HCl (200 mL). After 3 h, the reaction was diluted with  $\text{H}_2\text{O}$  and extracted with  $\text{Et}_2\text{O}$ . The aqueous phase was then neutralized by the addition of 0.5 N NaOH. The basic phase was then extracted with  $\text{CH}_2\text{Cl}_2$ , dried over  $\text{MgSO}_4$ , and concentrated to give a white solid (11.9 g, 90%).  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  4.56-4.71 (m, 2H), 4.23-4.31 (m, 1H), 3.40-3.49 (m, 3H), 3.11 (d, 2H,  $J = 4.4$ ), 1.17-2.23 (m, 8H), 1.13 (s, 3H), 0.93-1.12 (m, 3H).

10




(2R)-2-(4-chlorobenzensulfonylamino)-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)-4-fluorobutan-1-one:

20 To a solution of (2R)-2-amino-1-((1'S),(5'S)-10',10'-dimethyl-3',3'-dioxo-3,3'-thia-4'-aza-tricyclo-[5.2.1.0^{1,4}]dec-4'-yl)-4-fluorobutan-1-one: (12 g, 36 mmol) and  $\text{Et}_3\text{N}$  (10.4 mL, 72.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (350 mL) was added 4-chlorobenzensulfonyl chloride (9.1 g, 43 mmol) in one portion. After 18 h the reaction was concentrated and the resulting residue was taken into EtOAc and washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{MgSO}_4$ , and concentrated. The material was


- 86 -

- then further purified by silica gel chromatography (30% EtOAc/hexanes) to afford the titled compound (16.0 g, 92%) as a white wax. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.79 (d, 2H, *J* = 8.0), 7.43 (d, 2H, *J* = 8.0), 5.69 (br d, 8.0), 4.42-4.77 (m, 4H), 3.71-3.72 (m, 1H), 3.10 (ABq, 2H, *J* = 9, 4.4), 2.11-2.29 (m, 2H), 1.33-1.99 (m, 6H), 1.04 (s, 3H), 0.91 (s, 3H).



(2R)-2-((4-chlorobenzensulfonyl)amino)-4-fluorobutanoic acid:

- 10 To a rapidly stirred solution of (2R)-2-(4-chlorobenzensulfonyl)amino)-1-((1*S*),(5*S*)-10',10'-dimethyl-3',3'-dioxo-3',4'-thia-4',-aza-tricyclo-[5.2.1.0<sup>2,3</sup>]dec-4'-yl)-4-fluorobutan-1-one (16 g, 32 mmol) in acetonitrile (200 mL) was added LiBr (13.9 g, 16 mmol), tetrabutylammonium bromide (4.13 g, 12.8 mmol), and LiOH (5.45 g, 0.130 mol). After 4.5 h the reaction was concentrated to half volume then diluted with H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The aqueous layer was acidified with 1 N HCl and extracted with EtOAc. The EtOAc extracts were combined, dried over MgSO<sub>4</sub>, and concentrated to give a white solid of which 9.4 g was taken directly towards the next step. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 8.39 (d, 1H, *J* = 9.0), 7.76 (d, 2H, *J* = 6.8), 7.64 (d, 2H, *J* = 6.8), 7.00 (br s, 1H), 4.29-4.48 (m, 2H), 3.80-3.88 (m, 1H), 1.66-1.96 (m, 2H).




25

- 87 -

(2R)-2-((4-chlorobenzensulfonyl)amino)-4-fluorobutanoic acid amide:

- To a solution of (2R)-2-(4-chlorobenzensulfonyl)amino)-4-fluorobutanoic acid (9.0 g, 31 mmol) in DMF (250 mL) was added consecutively 1-hydroxybenzotriazole hydrate (6.2 g, 46 mmol), N,N-diisopropylethylamine (23 mL, 124 mmol), ammonium chloride (3.34 g, 62 mmol), and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (8.8 g, 46 mmol) under N<sub>2</sub>. The resulting solution was stirred at rt for 18 h. The solution was poured over ice water (500 mL) and the solid was filtered off and dried. The material was then precipitated from 10% EtOAc/hexanes to afford a clean white solid (4.5g) in 50% yield. [α]<sub>D</sub><sup>20</sup> = -21.0 (c 1.00, DMF); MS (ESI) (M - H)<sup>+</sup> 293.01; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 8.12 (d, 1H, *J* = 8.8), 7.77 (d, 2H, *J* = 7.0), 7.62 (d, 2H, *J* = 7.0), 7.38 (br s, 1H), 7.03 (br s, 1H), 4.22-4.47 (m, 2H), 3.71-3.85 (m, 1H), 1.65-1.92 (m, 2H).




(2R)-2-((4-chlorobenzensulfonyl)-(4-cyanobenzyl)-amino)-4-fluorobutyramide

(Example 360):

- (2R)-2-(4-chlorobenzensulfonyl)amino)-4-fluorobutyramide (20 mg, 0.7 mmol) was converted to the title compound as in Reaction Scheme 1, method A to afford the titled compound (208 mg) in 73% yield. MS (ESI) (M - H)<sup>+</sup> 407.99; [α]<sub>D</sub><sup>20</sup> = +39.13 (c 1.00, MeOH); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.72 (d, 2H, *J* = 8.4), 7.58 (d, 2H, *J* = 8.4), 7.50 (d, 2H, *J* = 8.4), 7.45 (d, 2H, *J* = 8.4), 6.29 (br s, 1H), 5.21 (br s, 1H), 4.19-4.67 (m, 5H), 2.17-2.28 (m, 1H), 1.49-1.61 (m, 1H).

25




Exemplification of Reaction Scheme 192-(4-Chlorobenzenesulfonylamino)-6-fluoro-hexanoic acid amide (III):

A mixture of (benzhydrylidene-amino)acetic acid ethyl ester (8.6 g, 32 mmol), 4-bromo-1-fluorobutane (10.0 g, 64.5 mmol),  $K_2CO_3$  (13.4 g, 96.9 mmol), tetrabutylammonium bromide (2.1 g, 6.5 mmol), and acetonitrile (300 mL) was heated at reflux for 72 h. The reaction was cooled to rt and filtered through a sintered glass funnel. The filtrate was concentrated *in vacuo*. The residue was dissolved in diethyl ether (250 mL) and a white solid precipitated. The solid was removed by vacuum filtration. A solution of 1 N HCl (100 mL) was added to the filtrate, which contained the crude product (2-(benzhydrylideneamino)-6-fluoro-hexanoic acid ethyl ester). The resulting biphasic mixture was stirred vigorously for 3 h. The mixture was transferred to a separatory funnel. The aqueous layer was collected. The organic layer was extracted with 1 N HCl (30 mL). The combined aqueous layers were washed with 200 mL of diethyl ether.

Concentrated HCl (10.8 mL) was added to the aqueous portion and the resulting solution was heated at reflux for 6 h. The reaction mixture was cooled to rt and concentrated *in vacuo*. Toluene was added to the residue and the mixture was reconcentrated *in vacuo* to afford 2-amino-6-fluoro-hexanoic acid hydrochloride as a white solid. The crude amino acid salt was used without purification or characterization. 2-Amino-6-fluorohexanoic acid hydrochloride (32.3 mmol, theoretically) was suspended in anhydrous methanol (300 mL) and cooled to 0 °C. Thionyl chloride (10.3 mL, 129 mmol) was slowly added over 5 min. The resulting solution was allowed to warm to rt and stir for 18 h. The reaction mixture was concentrated *in vacuo* to afford methyl 2-amino-6-fluoro-hexanoic

acid hydrochloride. Toluene (100 mL) and 28% ammonia in water (75 mL) were added to the crude amino ester. The resulting biphasic mixture was stirred vigorously at rt for 24 h. The reaction mixture was concentrated *in vacuo*. The residual solid was suspended in toluene (200 mL) and reconcentrated *in vacuo* to afford 6-fluorohexanoic acid amide (II) as a white solid. The crude amino acid amide was dissolved in anhydrous DMF (50 mL) and  $CH_2Cl_2$  (350 mL) and reacted with 4-chlorobenzenesulfonylchloride (82 g, 32.3 mmol) and  $Et_3N$  (13.5 mL, 96.9 mmol). After 2 h, a second portion of 4-chlorobenzenesulfonylchloride (1.70 g, 8.1 mmol) was added. After an additional 18 h, the resulting mixture was poured into 1 N HCl (500 mL). The organic layer was collected and washed with water (2 x 500 mL). Hexane (600 mL) was added to the organic layer. A white precipitate formed. The solid was collected by vacuum filtration, rinsed with cold ethanol (50 mL), and dried *in vacuo* to afford 4.95 g (48% yield, 6 steps) of 2-(4-chlorobenzenesulfonylamino)-6-fluoro-hexanoic acid amide (III). LCMS ( $M+Na$ )<sup>+</sup> 345.2; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) 7.99 (d, 1H, *J* = 8.8), 7.77 (d, 2H, *J* = 8.8), 7.62 (d, 2H, *J* = 8.8), 7.29 (s, 1H), 6.95 (s, 1H), 4.34 (dt, 2H, *J*<sub>d</sub> = 47.5, *J*<sub>t</sub> = 6.1), 3.65 (dt, 1H, *J*<sub>d</sub> = 5.6, *J*<sub>t</sub> = 8.6), 1.60-1.39 (m, 4H), 1.36-1.15 (m, 2H). Anal. Calcd for  $C_{16}H_{18}ClFN_2O_5S$ : C, 44.65; H, 4.99; N, 8.67. Found: C, 44.61; H, 5.08; N, 8.75.


2-[(4-Chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-6-fluoro-hexanoic acid amide (Example 333):

2-(4-Chlorobenzenesulfonylamino)-6-fluoro-hexanoic acid amide (0.500 g, 1.55 mmol) was converted to the title compound (360 mg, 50% yield) as in

- 90 -

- Reaction Scheme 1, method A. LCMS (M+Na)<sup>+</sup> 459.9; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  7.82 (d, 2H, *J* = 8.8), 7.79 (d, 2H, *J* = 8.5), 7.63 (d, 2H, *J* = 8.8), 7.58 (d, 2H, *J* = 8.3), 7.52 (s, 1H), 7.09 (s, 1H), 4.82 (ABq, 2H,  $\Delta\nu$  = 37.2, *J*<sub>AB</sub> = 17.6), 4.34 (dd, 1H, *J* = 8.0, 6.6), 4.25 (dt, 2H, *J*<sub>a</sub> = 47.2, *J* = 5.7), 1.58 (m, 1H), 1.49-1.12 (m, 5H); Anal. Calcd for C<sub>26</sub>H<sub>21</sub>ClF<sub>3</sub>N<sub>3</sub>O<sub>5</sub>S: C, 54.85; H, 4.83; N, 9.59. Found: C, 54.92; H, 4.76; N, 9.54.

#### Exemplification of Reaction Scheme 20




(2*R*)-2-(4-Chlorobenzensulfonylamino)-1-((1'*S*),(5'*S*)-10',10'-dimethyl-3',3'-dioxo-3'λ<sup>6</sup>-thia-4'-aza-tricyclo-[5.2.1.0<sup>1,3</sup>]dec-4'-yl)-4-fluoro-4-methyl-pentan-1-one:

- 15 To a solution of (2*R*)-2-(4-chlorobenzensulfonylamino)-1-((1'*S*),(5'*S*)-10',10'-dimethyl-3',3'-dioxo-3'λ<sup>6</sup>-thia-4'-aza-tricyclo-[5.2.1.0<sup>1,3</sup>]dec-4'-yl)-4-methyl-4-penten-1-one [500 mg, 1 mmol, prepared as in Reaction Scheme 18 from *N*-2-(benzhydrylidene-amino)-1-((1'*S*),(5'*S*)-10',10'-dimethyl-3',3'-dioxo-3'λ<sup>6</sup>-thia-4'-aza-tricyclo-[5.2.1.0<sup>1,3</sup>]dec-4'-yl) ethanone (ref. Josien, H.; Martin, A.; Chassaign, G. *Tetrahedron Lett.* 1991, 32, 6547) and 1-bromo-2-methyl-2-propene] in THF (5 mL) at 0 °C was added hydrofluoric acid/pyridine (10 mL). The reaction mixture was allowed to warm to rt and stir for 18 h. The reaction contents were carefully added to a saturated aqueous solution of NaHCO<sub>3</sub> (300 mL). The aqueous mixture was extracted with EtOAc (3 x 100 mL). The combined organic layers were sequentially washed with 1 N HCl (200 mL) and brine (100 mL). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to afford 490 mg (94%) of the title compound as a white

- 91 -


solid: <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  7.83 (d, 2H, *J* = 8.8), 7.45 (d, 2H, *J* = 8.8), 5.37 (d, 1H, *J* = 8.1), 4.65 (m, 1H), 3.64 (t, 1H, *J* = 6.4), 3.43 (ABq, 2H,  $\Delta\nu$  = 5.4, *J*<sub>AB</sub> = 13.7), 2.19-1.83 (m, 7H), 1.41-1.31 (m, 8H), 1.04 (s, 3H), 0.94 (s, 3H).



(2*R*)-2-(4-Chlorobenzensulfonylamino)-4-fluoro-4-methyl-pentanoic acid  
amide:

- 10 (2*R*)-2-(4-Chlorobenzensulfonylamino)-1-((1'*S*),(5'*S*)-10',10'-dimethyl-3',3'-dioxo-3'λ<sup>6</sup>-thia-4'-aza-tricyclo-[5.2.1.0<sup>1,3</sup>]dec-4'-yl)-4-fluoro-4-methyl-pentan-1-one was converted to the title compound in two steps as in Reaction Scheme 18 (165 mg, 55% yield): LCMS (M+Na)<sup>+</sup> 345.1; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  8.10 (d, 1H, *J* = 9.2), 7.77 (d, 2H, *J* = 8.5), 7.62 (d, 2H, *J* = 8.9), 7.34 (s, 1H), 6.92 (s, 1H), 3.85 (m, 1H), 1.89 (m, 1H), 1.74 (m, 1H), 1.31 (d, 3H, *J* = 21.7), 1.29 (d, 3H, *J* = 21.9).

#### Exemplification of Reaction Scheme 21




Ethyl 2-(4-chlorobenzensulfonylamino)-4-methyl-4-pentenoate:

- 20 A solution of ethyl 2-amino-4-methyl-4-pentenoate (2.84 g, 18.1 mmol, prepared as in Reaction Scheme 19 from (benzhydrylideneamino)acetic acid ethyl ester and 1-bromo-2-methyl-2-propene) in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) was reacted with 4-chlorobenzensulfonyl chloride (4.20 g, 19.9 mmol) and Et<sub>3</sub>N (3.78 mL, 27.2

- 92 -

mmol). After 4 h, the resulting mixture was poured into 1 N aqueous HCl (500 mL) and extracted with EtOAc (3 x 150 mL). The organic layer was washed with brine (50 mL), dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The crude concentrate was purified using silica gel column chromatography (10:1 to 5:1 gradient, hexanes/EtOAc) to afford 3.04 g (25% yield over 3 steps) of ethyl 2-(4-chlorobenzenesulfonylamino)-4-methyl-4-pentenoate: LCMS (M+Na)<sup>+</sup> 354.2; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 7.77 (d, 2H, J = 9.1), 7.46 (d, 2H, J = 8.8), 5.07 (d, 1H, J = 9.0), 4.84 (s, 1H), 4.73 (s, 1H), 4.05 (m, 1H), 3.95 (q, 2H, J = 7.1), 2.40 (m, 2H), 1.66 (s, 3H), 1.13 (t, 3H, J = 7.1).




2-(4-Chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoic acid ethyl ester and 4-Chloro-N-(5,5-dimethyl-2-oxo-tetrahydro-furan-3-yl)-benzenesulfonamide:

- 15 Hydrogen fluoride-pyridine (10 mL) was added to a 0 °C solution of ethyl 2-(4-chloro-benzenesulfonylamino)-4-methyl-4-pentenoate (1.0 g, 3.0 mmol) in THF (15 mL). The reaction mixture was allowed to warm to rt. After 5 h, an additional portion (10 mL) of hydrogen fluoride-pyridine was added. The mixture was stirred for 24 h, then a third portion of hydrogen fluoride-pyridine (10 mL) was added. After a total of 53 h, the reaction was quenched with ice chips (20 mL). The crude mixture was poured into ice water (500 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 200 mL). The combined organic layers were washed with sat. aq. NaHCO<sub>3</sub> (100 mL) and concentrated *in vacuo*. The crude concentrate was purified using silica gel column chromatography (10:1 to 5:1 gradient, hexanes/EtOAc) to afford 0.395 g (37% yield) of ethyl 2-(4-chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoate and 0.425 g (46% yield) of 4-chloro-N-(5,5-dimethyl-2-oxo-tetrahydro-furan-3-yl)-
- 20
- 25

- 93 -

benzenesulfonamide. Data for ethyl 2-(4-chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoate: LCMS (M+Na)<sup>+</sup> 374.1, <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 7.78 (d, 2H, J = 8.9), 7.47 (d, 2H, J = 8.5), 5.19 (d, 1H, J = 7.9), 4.08 (m, 1H), 3.93 (m, 2H), 2.09-1.94 (m, 2H), 1.42 (d, 3H, J = 21.6), 1.37 (d, 3H, J = 21.6), 1.12 (t, 3H, J = 7.0). Data for 4-chloro-N-(5,5-dimethyl-2-oxo-tetrahydro-furan-3-yl)-benzenesulfonamide: LCMS (M+Na)<sup>+</sup> 326.0, <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) 8.41 (d, 1H, J = 9.1), 7.86 (d, 2H, J = 8.6), 7.67 (d, 2H, J = 8.8), 4.57 (m, 1H), 2.22 (dd, 1H, J = 12.4, 9.0), 1.72 (t, 1H, J = 12.0), 1.33 (s, 3H), 1.31 (s, 3H).




2-(4-Chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoic acid amide:

- 15 A solution of 2-(4-chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoic acid ethyl ester (457 mg, 1.30 mmol) in MeOH (20 mL) was treated with 10 N NaOH (780 µL, 7.8 mmol) at rt for 18 h. The crude reaction mixture was concentrated *in vacuo*. The residue was treated with water (50 mL) and 1 N HCl (20 mL). The aqueous solution was extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine (50 mL), dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to afford a white solid containing 2-(4-chlorobenzenesulfonylamino)-4-fluoro-4-methyl-pentanoic acid. A mixture of the crude solid, 1-hydroxybenzotriazole (263 mg, 1.95 mmol), diisopropylethylamine (670 mg, 5.2 mmol), ammonium chloride (140 mg, 2.6 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (373 mg, 1.95 mmol), and DMF (20 mL) was stirred at rt for 24 h. The crude mixture was poured into water (500 mL). The aqueous solution was extracted with EtOAc/hexane (90:10, 3 x 150 mL). The combined organic layers were washed with brine (50 mL), dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude concentrate was purified using silica gel column chromatography (95:5, chloroform/MeOH) to
- 20
- 25


- 94 -

afford 0.426 g (100% yield) of the title compound: LCMS (M+N<sup>+</sup>) 345.3; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.10 (d, 1H, J = 9.2), 7.77 (d, 2H, J = 8.5), 7.62 (d, 2H, J = 8.9), 7.34 (s, 1H), 6.92 (s, 1H), 3.85 (m, 1H), 1.89 (m, 1H), 1.74 (m, 1H), 1.31 (d, 3H, J = 21.7), 1.29 (d, 3H, J = 21.9).



2-[(4-Chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-4-fluoro-4-methyl-pentanoic acid amide (Example 357):

10 2-(4-Chlorobenzesulfonylamino)-4-fluoro-4-methyl-pentanoic acid amide was converted to the title compound as in Reaction Scheme 1, method A. LCMS (M+N<sup>+</sup>) 460.2; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.83 (d, 2H, J = 8.5), 7.75 (d, 2H, J = 8.3), 7.68 (s, 1H), 7.64 (d, 2H, J = 8.6), 7.49 (d, 2H, J = 8.1), 7.20 (s, 1H), 4.67 (ABq, 2H, Δν = 28.3, J<sub>a</sub> = 17.3), 4.54 (dd, 1H, J = 9.3, 3.2), 2.23 (m, 1H), 1.42 (m, 1H), 1.25 (d, 3H, J = 21.6), 1.21 (d, 3H, J = 21.7).



2-[(4-Chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-4-hydroxy-4-methyl-pentanoic acid amide (Example 443):


A sealed vial containing a mixture of 4-chloro-N-(5, 5-dimethyl-2-oxo-tetrahydro-furan-3-yl)-benzenesulfonylformamide (0.20 g, 0.66 mmol) and 28%

- 95 -

ammonia in water (3 mL) was heated in a microwave reactor at 80 °C for 40 min. The reaction mixture was cooled to rt and concentrated to dryness in vacuo to afford a white solid containing 2-(4-chlorobenzesulfonylamino)-4-hydroxy-4-methyl-pentanoic acid amide. The crude solid was converted to the title

5 compound (98 mg, 34% yield) as in Reaction Scheme 1, method A: LCMS (M+N<sup>+</sup>) 458.2; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.84 (d, 2H, J = 8.6), 7.76 (d, 2H, J = 8.3), 7.62 (d, 2H, J = 8.8), 7.51 (d, 2H, J = 8.3), 7.40 (s, 1H), 7.11 (s, 1H), 4.63 (ABq, 2H, Δν = 5.9, J<sub>a</sub> = 17.6), 4.56 (dd, 1H, J = 8.3, 2.5), 4.54 (s, 1H), 1.95 (dd, 1H, J = 13.7, 8.6), 1.26 (dd, 1H, J = 13.6, 2.4), 1.04 (s, 3H), 0.99 (s, 3H). Anal. Calcd for C<sub>20</sub>H<sub>22</sub>ClN<sub>2</sub>O<sub>5</sub>S: C, 55.10; H, 5.08; N, 9.64. Found: C, 54.96; H, 5.14; N, 9.58.

Exemplification of Reaction Scheme 22




2-[(4-Chlorobenzesulfonylamino)-5-hexenoic acid ethyl ester:

A mixture of (benzhydrylidene-amino)acetic acid ethyl ester (20 g, 74.8 mmol), 4-bromo-1-butene (10.1 g, 74.8 mmol), K<sub>2</sub>CO<sub>3</sub> (31.0 g, 224 mmol), tetrabutylammonium bromide (2.41 g, 7.48 mmol), and acetonitrile (150 mL) was heated at reflux for 6 h. The reaction was cooled to rt and filtered through a sintered glass funnel. The filtrate was concentrated *in vacuo*. The residue was dissolved in diethyl ether (250 mL) and a white solid precipitated. The solid was removed by vacuum filtration. A solution of 1 N HCl (150 mL) was added to the filtrate, which contained the crude product (2-(benzhydrylidene-amino)-hex-5-enoic acid ethyl ester). The resulting biphasic mixture was stirred vigorously for 18 h. The mixture was transferred to a separatory funnel. The aqueous layer was

- 96 -


- collected and concentrated *in vacuo*. The residue was dissolved in toluene (2 x 200 mL) and reconcentrated. The crude amino ester was dissolved in  $\text{CH}_2\text{Cl}_2$  and reacted with 4-chlorobenzenesulfonyl chloride (15.8 g, 74.8 mmol) and  $\text{Et}_3\text{N}$  (31.2 mL, 224 mmol). After 18 h, the resulting mixture was poured into 1 N HCl (500 mL). The organic layer was collected and washed sequentially with 1 N HCl (500 mL) and brine (50 mL). The organic layer was dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. The crude concentrate was purified using silica gel column chromatography (5:1, hexanes/ $\text{EtOAc}$ ) to afford 5.57 g (23% yield over 3 steps) of the title compound: LCMS ( $\text{M}+\text{Na}^+$ ) 354.0;  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  8.47 (d, 1H,  $J = 8.8$ ), 7.76 (d, 2H,  $J = 8.8$ ), 7.66 (d, 2H,  $J = 8.8$ ), 5.69 (m, 1H), 4.95–4.88 (m, 2H), 3.86 (q, 2H,  $J = 7.1$ ), 3.76 (m, 1H), 1.98 (m, 2H), 1.71–1.54 (m, 2H), 1.03 (t, 3H,  $J = 7.1$ ).



- 2-[(4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-hex-5-enoic acid ethyl ester.
- 2-[(4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-hex-5-enoic acid ethyl ester was made in a similar manner to Reaction Scheme 1 starting from 2-(4-chloro-benzenesulfonylamino)-hex-5-enoic acid ethyl ester. 2-[(4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-hex-5-enoic acid ethyl ester was isolated as a crude yellow solid (1.14 g) and used in the next step without further purification.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.71 (d, 2H,  $J = 8.0$ ), 7.61 (d, 2H,  $J = 8.0$ ), 7.53 (d, 2H,  $J = 8.0$ ), 7.46 (d, 2H,  $J = 8.0$ ), 5.54 (m, 2H), 4.90 (m, 2H), 4.74 (d, 1H,  $J = 16.0$ ), 4.48 (m, 2H), 3.90 (m, 1H), 1.95 (m, 2H), 1.81 (m, 1H), 1.48 (m, 1H), 1.11 (t, 3H,  $J = 8.0$ ).


- 97 -

2-[(4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-5-oxo-pentanoic acid ethyl ester




- A mixture of (4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-hex-5-enoic acid ethyl ester (1.14 g, 2.56 mmol), osmium tetroxide (0.030 g, 0.13 mmol), and trimethylamine N-oxide (0.41 g, 5.5 mmol) was dissolved in acetone (50 mL) and stirred for 4 h at rt. Upon completion, the solution was concentrated *in vacuo* and redissolved in 1.5:1 dioxane: $\text{H}_2\text{O}$  (50 mL). To this solution, sodium periodate (0.66 g, 3.07 mmol) was added and stirred at rt for 18 h. The reaction was then diluted with  $\text{EtOAc}$  (500 mL) and washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to give a crude colorless oil. Further purification by flash chromatography ( $\text{SiO}_2$ , 5 to 75 %  $\text{EtOAc}$ /hexanes) afforded 2-[(4-chlorobenzenesulfonyl)-(4-cyanobenzyl)-amino]-5-oxo-pentanoic acid ethyl ester (0.26 g) as a colorless oil in 23 % yield.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.57 (s, 1H), 7.69 (d, 2H,  $J = 8.0$ ), 7.51 (m, 6H), 5.99 ABq, 2H,  $\Delta\nu = 16$ ,  $J_{\text{AB}} = 16.8$ ), 4.47 (m, 1H), 3.89 (m, 2H), 2.53 (m, 1H), 2.32 (m, 1H), 2.11 (m, 1H), 1.61 (m, 1H), 1.06 (t, 3H,  $J = 8.0$ ).

- 98 -



2-[(4-chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-5,5-difluoro-pentanoic acid ethyl ester:

5 2-[(4-chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-5-oxo-pentanoic acid ethyl ester (0.05 g, 0.11 mmol) was slowly added to a solution of DAST (0.020 mL, 0.11 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) at rt and stirred for 16 h. The reaction was diluted with  $\text{CH}_2\text{Cl}_2$  (20 mL) and extracted with  $\text{H}_2\text{O}$  (2 x 25 mL). The combined organic layers were washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to give 2-[(4-chloro-benzesulfonyl)-(4-cyanobenzyl)-amino]-5,5-difluoro-pentanoic acid ethyl ester as a crude yellow residue (61 mg). This crude residue was taken onto the next step without further purification.




2-[(4-chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-5,5-difluoro-pentanoic acid amide (Example 377):

The crude 2-[(4-chlorobenzesulfonyl)-(4-cyanobenzyl)-amino]-5,5-difluoro-pentanoic acid ethyl ester (0.061 g, 0.13 mmol) was dissolved in MeOH (2 mL). To this mixture was added 10 N NaOH (0.052 mL, 0.52 mmol) and the

- 99 -

resulting solution was stirred at rt for 16 h. The reaction was diluted with  $\text{H}_2\text{O}$  (25 mL), acidified with 1 N HCl, and extracted with  $\text{CH}_2\text{Cl}_2$  (4 x 100 mL). The combined organic layers were dried over  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo* to give the carboxylic acid moiety as a crude colorless oil. The carboxylic acid intermediate was then dissolved in DMF (10 mL) and mixed with 1-hydroxybenzotriazole (0.030 g, 0.20 mmol),  $\text{Et}_3\text{N}$  (0.090 mL, 0.52 mmol),  $\text{NH}_4\text{Cl}$  (0.01 g, 0.26 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.04 g, 0.20 mmol) and stirred at rt for 72 h. The reaction was diluted with EtOAc (150 mL) and washed with  $\text{H}_2\text{O}$  (4 x 50 mL). The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo* to give a crude off-white solid. Further purification by flash chromatography ( $\text{SiO}_2$ , 5 to 85 % EtOAc/hexanes) afforded the titled compound (10.7 mg) as a white solid in 19 % yield. LCMS ( $\text{M}+\text{Na}^+$ ) 464.01;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.69 (d, 2H,  $J = 8.3$ ), 7.60 (d, 2H,  $J = 8.3$ ), 7.49 (m, 4H), 6.18 (br s, 1H), 5.67 (t, 1H,  $J = 5.6$ , 4.0), 5.22 (br s, 1H), 4.52 (ABq, 2H,  $\Delta\nu = 16$ ,  $J_{\text{AB}} = 100$ ), 4.34 (m, 1H), 2.03 (m, 1H), 1.68 (m, 1H), 1.38 (m, 1H), 0.86 (m, 1H).

#### Exemplification of Reaction Scheme 23




(2R)-2-[(2-bromo-acetyl)amino]-benzyl-[(4-chlorobenzesulfonyl)-amino]-4-methyl-pentanoic acid amide:

To a solution of (2R)-2-[(4-aminobenzyl)-(4-chlorobenzesulfonyl)amino]-4-methyl-pentanoic acid amide (248 mg, 0.56 mmol) and  $\text{Et}_3\text{N}$  (176 mg, 1.74 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 mL) was added bromoacetylchloride

- 100 -

(105 mg, 0.67 mmol). The reaction mixture was stirred overnight at rt. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (5 mL), washed with 1 N HCl, brine, and dried through a cotton plug. The solvent was removed *in vacuo*. Purification by flash chromatography ( $\text{SiO}_2$ , 10% acetone/ $\text{CH}_2\text{Cl}_2$ ) afforded the title

compound (124 mg) in 42 % yield. MS (ESI), ( $\text{M}+\text{H}^+$ ) 531.86,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  8.78 (br s, NH), 7.95 (d, 2H,  $J=8.0$ ), 7.82 (d, 2H,  $J=8.0$ ), 7.42 (d, 2H,  $J=8.0$ ), 7.33 (d, 2H,  $J=8.0$ ), 6.20 (br s, 1H), 5.20 (br s, 1H), 4.30 (s, 2H), 4.22 (d, 1H,  $J_{\text{ab}}=1.6$ ), 4.14 (d, 1H,  $J_{\text{ab}}=1.6$ ), 3.25 (t, 1H,  $J=6.0$ ), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H,  $J=7.0$ ), 0.94 (d, 3H,  $J=7.0$ )



(2R)-2-((4-Chlorobenzene)sulfonyl)-[4-(2-dimethylamino-ethylamino)-benzyl]-amino]-4-methyl-pentanoic acid amide (Example 308):


To a solution of (2R)-2-[[4-(2-bromo-ethylamino)-benzyl]-4-chlorobenzene)sulfonyl]-amino]-4-methyl-pentanoic acid amide (41 mg, 0.77 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was added excess 2.0 M dimethylamine in THF. The reaction mixture was stirred overnight. The solvent was removed *in vacuo*.

Purification by flash chromatography ( $\text{SiO}_2$ , 10% MeOH/ $\text{CH}_2\text{Cl}_2$ ) afforded the title compound (24 mg) in 63% yield. MS (ESI), ( $\text{M}+\text{H}^+$ ) 495.14,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  8.85 (s, 1H), 8.02 (d, 2H,  $J=8.0$ ), 7.75 (d, 2H,  $J=8.0$ ), 7.38 (d, 2H,  $J=8.0$ ), 7.29 (d, 2H,  $J=8.0$ ), 6.23 (br s, 1H), 5.39 (br s, 1H), 4.62 (m, 4H), 3.25 (t, 1H,  $J=6.0$ ), 2.95 (s, 6H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H,  $J=7.0$ ), 0.94 (d, 3H,  $J=7.0$ )

- 101 -

### Starting Materials

The following  $\alpha$ -amino amides were commercially available or obtained by standard methods from commercially available amino acids:



5,5,5-Trifluoro-2-aminopentanoic acid amide and 6,6,6-trifluoro-2-

aminohexanoic acid were prepared according to: Ojima, I.; Kato, K.; Nakashishi,

K. *J. Org. Chem.* 1989, 54, 4511.

The benzyl bromide used in the synthesis of the compounds of Examples

100 and 155 was prepared according to: Ishihara, Y.; Fujisawa, Y.; Furuyama, N.

PCT Int. Appl. WO 9846590; Senanayake, C.H.; Fang, Q.K.; Wilkinson, S.H.

PCT Int. Appl. WO 9833789.

The aldehydes required for the synthesis of the compounds of Examples

15 91, 248, 249, 289, 290, and 300 (see Reaction Scheme 2) were prepared as

exemplified for 4-(piperidin-1-yl)benzaldehyde. A suspension of 4-

fluorobenzaldehyde (0.48mL, 4 mmol),  $\text{K}_2\text{CO}_3$  (522 mg, 4 mmol), piperidine (340 mg, 4 mmol) in DMSO (5 mL) was heated in a sealed tube at 150°C for 18h. After which time, the reaction was concentrated and purified by silica gel

20 chromatography ( $\text{CH}_2\text{Cl}_2$ , then 2% MeOH/ $\text{CH}_2\text{Cl}_2$ ) to afford 4-(piperidin-1-yl)benzaldehyde, 748 mg, 98% yield.

The aldehydes used in the synthesis of the compounds of Examples 317, 318, and 320 were prepared as exemplified for 4-(piperidin-1-yl)-3-fluorobenzaldehyde. A suspension of 4,3-difluorobenzaldehyde (500 mg, 3.5 mmol),  $K_2CO_3$  (483 mg, 3.5 mmol), piperidine (298 mg, 3.5 mmol) in DMSO (5 mL) was heated in a sealed tube at 130 °C for 18h. The reaction mixture was allowed to cool to rt, concentrated and purified by silica gel chromatography ( $CH_2Cl_2$ , then 2% MeOH/ $CH_2Cl_2$ ) to afford 4-(piperidin-1-yl)-3-fluorobenzaldehyde, 740 mg, 99% yield.

The benzyl chloride used in the preparation of the compounds of Examples 433, 474, 480, and 500 was prepared by the following method. To a solution of 2-[(4-chloromethyl)phenyl]propan-2-ol (769 mg, 4.16 mmol) (ref: Creary, X.; Mehrsheikh-Mohammadi, M.E.; McDonald, S. J. *Org. Chem.* 1987, 52, 3254.) in  $CH_2Cl_2$  (14 mL) at -78 °C was added DAST (0.72 mL, 5.4 mmol). After 1.5 h, the solution was quenched with water and warmed to rt. The mixture was extracted with  $CH_2Cl_2$  (3x). The combined organic layers were dried ( $Na_2SO_4$ ) and concentrated. Flash column chromatography ( $SiO_2$ , 0 to 5% EtOAc/hexanes) provided the chloride as a pale yellow liquid (512 mg, 66%).  $^1H$  NMR ( $CDCl_3$ , 300MHz)  $\delta$  7.30-7.48 (m, 4H), 4.58 (s, 2H), 1.70 (s, 3H), 1.63 (s, 3H).

The preparation of the 2-trimethylsilyl ethyl ester of 4-bromomethyl benzoic acid, which is used in the synthesis of the compound of Example 470 is described by Graffner-Nordberg, M.; Sjoedin, K.; Tunek, A.; Hallberg, A. *Chem. Pharm. Bull.* 1998, 46, 591.

## 25 Conditions for Chromatographic Separation of Enantiomeric Mixtures

Condition 1: Example 345 was separated using the following method. 4.6 X 250 mm, 10  $\mu$ M, Chiralcel OD column, 1.0 mL/min, 85 % Hexane/EtOH 0.1 % DEA, over 20 min.

Condition 2: Example 346 was separated using the following method. 4.6 X 250 mm, 10  $\mu$ M, Chiralpak AD column, 1.0 mL/min, 80 % Hexane/EtOH 0.15 % DEA, over 20 min.


Condition 3: Example 347 was separated using the following method. 4.6 X 250 mm, 10  $\mu$ M, Chiralpak AD column, 1.0 mL/min, 65 % Hexane/TPA 0.1 % DEA, over 18 min.

Condition 4: Examples 365 and 366 were separated using the following method. 4.6 X 250 mm, 10  $\mu$ M, Chiralpak AD column, 1.0 mL/min, 75 % Hexane/EtOH 0.15 % DEA, over 25 min.

Condition 5: Examples 408 and 409 were separated using the following method. 4.6 X 250 mm, 10  $\mu$ M, Chiralcel OD column, 1.0 mL/min, 90 % Hexane/EtOH. 0.15 % DEA, over 36 min.



TABLE 4



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
1				1. 2	white solid	424.95	1.76 Method B	425.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.63 (d, 2H, J=7.0Hz), 7.42 (d, 2H, J=7.0Hz), 7.25 (d, 2H, J=8.0Hz), 6.79 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.35 (s, br, 1H), 4.36 (dd, 2H, J=50Hz, 15Hz), 4.26 (t, 1H, J=7.2Hz), 3.78 (s, 3H), 1.83 (m, 1H), 1.18-1.34 (m, 3H), 0.75 (d, 3H, J=7.0Hz), 0.67 (d, 3H, J=7.0Hz).
2				1	white solid	394.92	1.71 Method B	395.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.81 (d, 2H, J=7.0Hz), 7.60 (d, 2H, J=7.0Hz), 7.50 (s, br, 1H), 7.41 (d, 2H, J=8.0Hz), 7.32 (m, 2H), 7.24 (m, 1H), 7.18 (s, br, 1H), 4.76 (dd, 2H, J=50Hz, 15Hz), 4.36 (t, 1H, J=7.0Hz), 3.33 (s, 3H), 1.20-1.34 (m, 3H), 0.79 (d, 3H, J=6.0Hz), 0.46 (d, 3H, J=6.0Hz).
3				1	white solid	462.92	1.71 Method A	463.1	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=7.0Hz), 7.67 (d, 2H, J=7.0Hz), 7.57-7.62 (m, 4H), 7.06 (s, br, 1H), 4.79 (dd, 2H, J=70Hz, 17Hz), 4.38 (t, 1H, J=6.0Hz), 3.32 (s, 3H), 1.23-1.35 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.50 (d, 3H, J=6.0Hz).

- 104 -

PCT/US02/40605

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
4				1	white solid	429.37	1.69 Method A	429.1	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=7.0Hz), 7.61 (d, 2H, J=7.0Hz), 7.52 (s, br, 1H), 7.41 (d, 2H, J=8.2Hz), 7.37 (d, 2H, J=8.2Hz), 7.03 (s, br, 1H), 4.70 (dd, 2H, J=50Hz, 15Hz), 4.35 (t, 1H, J=7.0Hz), 1.28-1.30 (m, 3H), 0.80 (d, 3H, J=6.0Hz), 0.51 (d, 3H, J=6.0Hz).
5				1	white solid	422.98	1.71 Method A	423.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.76 (d, 2H, J=7.0Hz), 7.61 (d, 2H, J=7.0Hz), 7.42 (s, br, 1H), 7.16-7.20 (m, 5H), 6.99 (s, br, 1H), 4.24 (m, 1H), 3.45-3.51 (m, 1H), 3.10-3.18 (m, 1H), 2.52-2.59 (m, 2H), 1.95-2.05 (m, 1H), 1.69-1.80 (m, 1H), 1.55-1.34 (m, 3H), 0.84 (m, 6H).
6				1	white solid	419.93	1.45 Method A	420.13	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.8 (d, 2H, J=8.0Hz), 7.79 (d, 2H, J=8.0Hz), 7.63 (d, 2H, J=8.0Hz), 7.58 (m, 3H), 7.03 (s, br, 1H), 4.87 (dd, 2H, J=50Hz, 15Hz), 4.32 (t, 1H, J=7.0Hz), 1.28-1.30 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.53 (d, 3H, J=6.0Hz).
7				1	white solid	412.91	1.58 Method A	413.4	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.81 (d, 2H, J=7.0Hz), 7.61 (d, 2H, J=7.0Hz), 7.51 (s, br, 1H), 7.43 (m, 1H), 7.11-7.14 (m, 2H), 7.03 (s, br, 1H), 4.77 (dd, 2H, J=50Hz, 15Hz), 4.33 (t, 1H, J=6.0Hz), 1.21-1.31 (m, 3H), 0.80 (d, 3H, J=6.0Hz), 0.50 (d, 3H, J=6.0Hz).

- 105 -

WO 03/053912

PCT/US02/40605

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
8				1	white solid	412.91	1.58 Method A	413.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.82 (d, 2H, J=8.0Hz), 7.61 (d, 2H, J=8.0Hz), 7.55 (s, br, 1H), 7.39 (m, 1H), 7.05-7.32 (m, 4H), 7.03 (s, br, 1H), 4.78 (dd, 2H, J=50Hz, 15Hz), 4.38 (t, 1H, J=6.0Hz), 1.26-1.32 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.54 (d, 3H, J=6.0Hz)
9				1	white solid	451.03	2.99 Method C	451.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.75 (d, 2H, J=8.5Hz), 7.55 (d, 2H, J=8.5Hz), 7.51 (s, br, 1H), 7.25-7.29 (m, 4H), 7.03 (s, br, 1H), 4.69 (dd, 2H, J=25Hz, 14Hz), 4.35 (m, 1H), 1.21-1.31 (m, 3H), 1.25 (s, 9H), 0.81 (d, 3H, J=6.0Hz), 0.46 (d, 3H, J=6.0Hz)
10				1	white solid	424.95	1.56 Method A	425.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.0Hz), 7.50 (s, br, 1H), 7.21 (m, 1H), 7.04 (m, 1H), 6.80-6.95 (m, 3H), 4.70 (dd, 2H, J=50Hz, 15Hz), 4.37 (m, 1H), 3.70 (s, 3H), 1.30-1.39 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.52 (d, 3H, J=6.0Hz)
11				1	white solid	463.81	1.76 Method A	463	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.61 (d, 2H, J=8.0Hz), 7.55-7.60 (m, 2H), 7.40 (m, 1H), 7.10 (s, br, 1H), 4.72 (dd, 2H, J=50Hz, 15Hz), 4.40 (m, 1H), 1.26-1.40 (m, 3H), 0.83 (d, 3H, J=6.0Hz), 0.60 (d, 3H, J=6.0Hz)













Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
12				1	white solid	462.92	1.68 Method A	463.1	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.68-7.80 (m, 2H), 7.54-7.61 (m, 4H), 7.08 (s, br, 1H), 4.80 (dd, 2H, J=50Hz, 15Hz), 4.38 (t, 1H, J=6.0Hz), 1.26-1.33 (m, 3H), 0.82 (d, 3H, J=6.0Hz), 0.52 (d, 3H, J=6.0Hz)
13				1	white solid	442.94	1.55 Method A	443.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.78 (d, 2H, J=8.0Hz), 7.61 (d, 2H, J=8.0Hz), 7.56 (s, br, 1H), 7.05-7.24 (m, 4H), 4.65 (dd, 2H, J=50Hz, 15Hz), 4.40 (t, 1H, J=6.0Hz), 3.81 (s, 3H), 1.28-1.35 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.56 (d, 3H, J=6.0Hz)
14				1	white solid	410.92	1.51 Method B	411.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.45 (d, 2H, J=8.5Hz), 7.26 (d, 2H, J=8.5Hz), 6.73 (d, 2H, J=8.0Hz), 6.33 (s, br, 1H), 5.24 (s, br, 1H), 4.28 (dd, 2H, J=70Hz, 20Hz), 4.23 (m, 1H), 1.67-1.93 (m, 2H), 1.12-1.32 (m, 2H), 0.77 (d, 3H, J=7.0Hz), 0.68 (d, 3H, J=7.0Hz)
15				1	sticky pale yellow foam	408.5	1.58 Method B	409.3	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.0Hz), 7.74-7.59 (m, 3H), 7.02-7.13 (m, 2H), 6.84 (t, 1H, J=8.4Hz), 6.32 (s, br, 1H), 5.31 (s, br, 1H), 4.48 (dd, 2H, J=50Hz, 17Hz), 4.26 (m, 1H), 3.85 (s, 3H), 1.79-1.84 (m, 1H), 1.25-1.30 (m, 1H), 1.04-1.11 (m, 1H), 0.72 (d, 3H, J=7.0Hz), 0.63 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
16				1	white film	426.49	1.63 Method B	427.3	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.73-7.77 (m, 2H), 6.81-7.18 (m, 5H), 6.29 (s, br, 1H), 5.37 (s, br, 1H), 4.45 (dd, 2H, J=50Hz, 17Hz), 4.26 (m, 1H), 3.89 (s, 3H), 1.76-1.84 (m, 1H), 1.26-1.33 (m, 1H), 1.08-1.17 (m, 1H), 0.74 (d, 3H, J=7.0Hz), 0.66 (d, 3H, J=7.0Hz)
17				1	pale yellow solid	476.49	1.79 Method B	477.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.84 (d, 2H, J=8.0Hz), 7.72 (d, 1H, J=8.0Hz), 7.10 (d, 1H, J=9.5Hz), 6.99 (d, 2H, J=8.5Hz), 6.82 (t, 1H, J=8.5Hz), 6.21 (s, br, 1H), 5.35 (s, br, 1H), 4.46 (dd, 2H, J=50Hz, 15Hz), 4.31 (t, 1H, J=7.5Hz), 3.87 (s, 3H), 1.78-1.85 (m, 1H), 1.30-1.35 (m, 1H), 1.12-1.21 (m, 1H), 0.77 (d, 3H, J=7.0Hz), 0.68 (d, 3H, J=7.0Hz)
18				1	pale yellow solid	476.49	1.76 Method B	477.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.79-7.89 (m, 3H), 7.59-7.64 (m, 1H), 7.01-7.08 (m, 1H), 6.83 (t, 1H, J=8.8Hz), 6.25 (s, br, 1H), 5.42 (s, br, 1H), 4.45 (dd, 2H, J=50Hz, 17Hz), 4.33 (m, 1H), 3.88 (s, 3H), 1.78-1.85 (m, 1H), 1.31-1.35 (m, 1H), 1.17-1.23 (m, 1H), 0.77 (d, 3H, J=7.0Hz), 0.70 (d, 3H, J=7.0Hz)
19				1	white solid	442.94	1.73 Method B	443.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.42-7.63 (m, 4H), 7.02-7.10 (m, 2H), 6.86 (t, 1H, J=8.5Hz), 6.24 (s, br, 1H), 5.42 (s, br, 1H), 4.45 (dd, 2H, J=50Hz, 17Hz), 4.27 (m, 1H), 3.87 (s, 3H), 1.79-1.87 (m, 1H), 1.27-1.33 (m, 1H), 1.14-1.22 (m, 1H), 0.78 (d, 3H, J=7.0Hz), 0.71 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
20				1	pale yellow oil	422.52	1.68 Method B	423.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.52-7.57 (m, 2H), 7.37-7.39 (m, 2H), 7.02-7.12 (m, 2H), 6.84 (t, 1H, J=8.5Hz), 6.34 (s, br, 1H), 5.35 (s, br, 1H), 4.45 (dd, 2H, J=50Hz, 17Hz), 4.25 (m, 1H), 3.86 (s, 3H), 2.38 (s, 3H), 1.78-1.87 (m, 1H), 1.25-1.31 (m, 1H), 1.04-1.11 (m, 1H), 0.72 (d, 3H, J=7.0Hz), 0.65 (d, 3H, J=7.0Hz)
21				1	pale yellow oil	422.52	1.67 Method B	423.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.64 (d, 2H, J=8.0Hz), 7.25-7.30 (m, 2H), 7.02-7.12 (m, 2H), 6.84 (t, 1H, J=8.5Hz), 6.34 (s, br, 1H), 5.32 (s, br, 1H), 4.45 (dd, 2H, J=50Hz, 17Hz), 4.26 (t, 1H, J=10Hz), 3.86 (s, 3H), 2.42 (s, 3H), 1.76-1.85 (m, 1H), 1.25-1.31 (m, 1H), 1.05-1.12 (m, 1H), 0.72 (d, 3H, J=7.0Hz), 0.62 (d, 3H, J=7.0Hz)
22				3	clear oil	455.02	1.94 Method B	455.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.82 (d, 2H, J=8.0Hz), 7.68 (d, 2H, J=8.0Hz), 7.56 (s, br, 1H), 7.45 (m, 1H), 7.16 (m, 1H), 7.04 (s, 1H), 6.86 (m, 1H), 4.25 (m, 1H), 3.94 (m, 1H), 3.40-3.55 (m, 2H), 3.02-3.14 (m, 1H), 1.18-1.69 (m, 10H), 0.76 (s, br, 6H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
23				1	white solid	441.01	1.67 Method A	441.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.79 (d, 2H, J=8.0Hz), 7.59 (s, br, 1H), 7.31 (d, 2H, J=8.0Hz), 7.19 (d, 2H, J=8.0Hz), 7.03 (s, br, 1H), 4.68 (dd, 2H, J=50Hz, 15Hz), 4.35 (t, 1H, J=7.0Hz), 3.32 (s, 3H), 1.24-1.35 (m, 3H), 0.81 (d, 3H, J=6.0Hz), 0.51 (d, 3H, J=6.0Hz)
24				1	white solid	467.86	1.75 Method A	469.1	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.42 (s, br, 1H), 7.01 (s, 1H), 4.25 (m, 1H), 3.35-3.51 (m, 3H), 3.08-3.14 (m, 1H), 1.19-1.82 (m, 11H), 0.86 (d, 6H, J=6.0Hz)
25				3	clear oil	474.07	1.34 Method A	474.4	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.82 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.42 (s, br, 1H), 6.99 (s, 1H), 4.25 (m, 1H), 3.51-3.60 (s, br, 4H), 3.18-3.41 (m, 2H), 2.25-2.35 (s, br, 4H), 2.27 (m, 2H), 1.15-1.62 (m, 9H), 0.80 (d, 6H, J=6.0Hz)
26				3	clear oil	472.09	1.27 Method A	472	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.42 (s, br, 1H), 7.00 (s, 1H), 4.23-4.26 (m, 1H), 3.22-3.45 (m, 1H), 3.11-3.14 (m, 1H), 2.16-2.28 (m, 5H), 1.19-1.52 (m, 18H), 0.86 (m, 6H)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
27				3	clear oil	490.13	1.21 Method A	490	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.44 (s, br, 1H), 7.06 (s, 1H), 4.21-4.25 (m, 1H), 3.55 (s, br, 4H), 3.19-3.40 (m, 4H), 2.43-2.57 (m, 8H), 2.23-2.30 (m, 2H), 1.18-1.59 (m, 9H), 0.82 (m, 6H)
28				3	clear oil	446.01	1.08 Method A	446.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.84 (d, 2H, J=8.0Hz), 7.66 (d, 2H, J=8.0Hz), 7.49 (s, br, 1H), 7.02 (s, 1H), 4.23-4.26 (m, 1H), 3.94 (s, br, 2H), 3.71 (s, br, 2H), 3.52-3.57 (m, 1H), 3.14-3.17 (m, 1H), 3.06 (s, br, 4H), 1.17-1.65 (m, 7H), 0.86 (m, 6H)
29				1	white solid	428.91	1.51 Method A	429.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=9.0Hz), 7.47 (d, 2H, J=9.0Hz), 7.13 (d, 1H, J=11.2Hz), 7.00 (d, 1H, J=8.0Hz), 6.88 (m, 1H), 6.27 (s, br, 1H), 5.49 (s, br, 1H), 5.24 (s, br, 1H), 4.40 (dd, 2H, J=90Hz, 18Hz), 3.20 (m, 1H), 1.09-1.82 (m, 3H), 0.77 (d, 3H, J=7.0Hz), 0.68 (d, 3H, J=7.0Hz)
30				1	white solid	484.59	1.89 Method A	485.0	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.73 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.18-7.27 (m, 4H), 6.25 (s, br, 1H), 5.29 (s, br, 1H), 4.40-4.69 (m, 3H), 1.80-1.88 (m, 1H), 1.29 (s, 9H), 1.25-1.33 (m, 2H), 0.80 (d, 3H, J=7.0Hz), 0.69 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
31				1	white solid	430.61	1.82 Method A	431.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.60 (d, 2H, J=8.2Hz), 7.24-7.39 (m, 6H), 6.34 (s, br, 1H), 5.19 (s, br, 1H), 4.42-4.44 (m, 2H), 4.30 (t, 1H, J=8Hz), 2.41 (s, 3H), 1.74-1.83 (m, 1H), 1.28 (s, 9H), 1.25-1.33 (m, 1H), 0.93-1.01 (m, 1H), 0.72 (d, 3H, J=7.0Hz), 0.60 (d, 3H, J=7.0Hz)
32				3	brown oil	460.04	1.15 Method A	460.2	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.83 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.42 (s, br, 1H) 7.00 (s, 1H), 4.22-4.26 (m, 1H), 3.55 (s, br, 4H), 3.11-3.32 (m, 2H), 2.13-2.37 (m, 6H), 1.12-1.51 (m, 9H), 0.86 (m, 6H)
33				1	white solid	400.97	1.85 Method A	401	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.85 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.32 (s, br, 1H) 7.05 (s, br, 1H), 4.19 (t, 1H, J=7.5Hz), 2.99-3.02 (m, 1H), 1.07-1.65 (m, 14H), 0.78-0.84 (m, 7H)
34				3	clear Oil	476.1	1.28 Method A	476.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.76 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 6.55 (s, br, 1H), 5.41 (s, 1H), 4.17-4.20 (m, 1H), 3.26-3.38 (m, 1H), 3.13-3.17 (m, 1H), 2.62-2.92 (m, 8H), 2.32-2.36 (m, 2H), 1.85-1.87 (m, 1H), 1.25-1.47 (m, 7H), 0.96-0.99 (m, 1H), 0.75 (d, 3H, J=6.6Hz), 0.72 (d, 3H, J=6.6Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
35				3	clear oil	462.08	1.31 Method A	462.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.75 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 6.51 (s, br, 1H), 6.05 (s, 1H), 4.15-4.18 (m, 1H), 3.56-3.78 (m, 4H), 3.45-3.47 (m, 1H), 3.09-3.14 (m, 1H), 2.90-3.08 (m, 4H), 2.61-2.69 (m, 2H), 1.62-2.05 (m, 5H), 1.21-1.29 (m, 1H), 0.80-0.83 (m, 1H), 0.78 (d, 3H, J=6.6Hz), 0.71 (d, 3H, J=6.6Hz)
36				3	clear oil	418.0	1.25 Method A	418.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.5Hz), 7.51 (d, 2H, J=8.5Hz), 6.56 (s, br, 1H), 5.41 (s, 1H), 4.18-4.21 (m, 1H), 3.15-3.32 (m, 2H), 2.25-2.28 (m, 2H), 2.24 (s, 6H), 1.84-1.88 (m, 1H), 1.25-1.49 (m, 7H), 0.97-1.00 (m, 1H), 0.71-0.74 (m, 6H)
37				1	tan solid	428.91	1.5 Method A	429.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.57 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=9.0Hz), 7.03-7.13 (m, 2H), 6.80 (t, 1H, J=8.8Hz), 6.17 (s, br, 1H), 5.39 (s, br, 1H), 4.48 (dd, 2H, J=55Hz, 16Hz), 3.84 (s, 3H), 3.75-3.84 (m, 1H), 1.11-1.30 (m, 1H), 0.91 (d, 3H, J=7.0Hz), 0.54 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
38				1	white solid	412.46	1.46 Method A	413.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65-7.70 (m, 2H), 7.04-7.12 (m, 4H), 6.79 (t, 1H, J=8.5Hz), 6.21 (s, br, 1H), 5.27 (s, br, 1H), 4.48 (dd, 2H, J=50Hz, 15Hz), 3.86 (s, 3H), 3.79-3.85 (m, 1H), 2.16-2.22 (m, 1H), 0.90 (d, 3H, J=7.0Hz), 0.51 (d, 3H, J=7.0Hz).
39				1	white solid	403.48	1.47 Method A	404.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.77-7.80 (m, 2H), 7.59 (d, 2H, J=8.0Hz), 7.49 (d, 2H, J=8.0Hz), 7.17-7.22 (m, 2H), 6.17 (s, br, 1H), 5.20 (s, br, 1H), 4.50 (dd, 2H, J=60Hz, 17Hz), 4.28 (t, 1H, J=10Hz), 1.74-1.83 (m, 1H), 1.25-1.33 (m, 1H), 0.99-1.10 (m, 1H), 0.77 (d, 3H, J=7.0Hz), 0.66 (d, 3H, J=7.0Hz).
40				1	white solid	453.49	1.65 Method A	454.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.88 (d, 2H, J=8.2Hz), 7.78 (d, 2H, J=8.5Hz), 7.59 (d, 2H, J=8.5Hz), 7.49 (d, 2H, J=8.5Hz), 6.10 (s, br, 1H), 5.19 (s, br, 1H), 4.59 (dd, 2H, J=50Hz, 16Hz), 4.33 (t, 1H, J=10Hz), 1.76-1.81 (m, 1H), 1.25-1.35 (m, 1H), 1.02-1.07 (m, 1H), 0.78 (d, 3H, J=7.0Hz), 0.65 (d, 3H, J=7.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
41				1	white solid	399.52	1.53 Method A	400.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.2Hz), 7.47 (d, 2H, J=8.0Hz), 7.31 (d, 2H, J=8.5Hz), 6.24 (s, br, 1H), 5.16 (s, br, 1H), 4.50 (dd, 2H, J=50Hz, 17Hz), 4.27 (t, 1H, J=10Hz), 2.44 (s, 3H), 1.74-1.83 (m, 1H), 1.25-1.33 (m, 1H), 0.93-1.01 (m, 1H), 0.74 (d, 3H, J=7.0Hz), 0.63 (d, 3H, J=7.0Hz).
42				3	clear oil	441.0	1.27 Method A	441.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.74 (d, 2H, J=8.2Hz), 7.67 (s, 1H), 7.50 (d, 2H, J=8.2Hz), 7.10 (s, 1H), 6.93 (s, 1H), 6.51 (s, br, 1H), 5.55 (s, br, 1H), 4.14-4.17 (m, 1H), 3.97 (t, 2H, J=6.0Hz), 3.29-3.35 (m, 1H), 3.09-3.14 (m, 1H), 1.62-1.66 (m, 3H), 1.55-1.62 (m, 2H), 1.25-1.29 (m, 3H), 0.80-0.83 (m, 1H), 0.74 (d, 3H, J=6.6Hz), 0.71 (d, 2H, J=6.6Hz).
43				3	clear oil	446.06	1.29 Method A	446.3	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.76 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 6.61 (s, br, 1H), 5.45 (s, 1H), 4.15-4.18 (m, 1H), 3.09-3.24 (m, 2H), 2.50-2.58 (m, 4H), 2.31-2.39 (m, 2H), 1.92-1.99 (m, 1H), 1.15-1.59 (m, 8H), 1.00-1.04 (m, 7H), 0.71-0.74 (m, 6H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
44				1	tan wax	428.91	1.58 Method A	429.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J=8.0Hz), 7.45 (d, 2H, J=9.0Hz), 7.03-7.15 (m, 2H), 6.84 (t, 1H, J=8.0Hz), 6.25 (s, br, 1H), 5.19 (s, br, 1H), 4.45 (dd, 2H, J=80Hz, 18Hz), 4.19-4.22 (m, 1H), 3.87 (s, 3H), 1.82-1.95 (m, 1H), 0.96-1.30 (m, 3H), 0.72-0.77 (m, 3H).
45				3	clear oil	506.11	1.39 Method A	506.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.75 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.5Hz), 7.01-7.13 (m, 4H), 6.55 (s, br, 1H), 5.39 (s, br, 1H), 4.18 (t, 1H, J=6.0Hz), 3.32-3.60 (m, 1H), 3.16-19 (m, 1H), 2.91 (s, br, 2H), 2.76 (s, br, 2H), 2.52 (s, br, 2H), 1.27-1.86 (m, 8H), 0.97-1.00 (m, 1H), 0.73 (m, 6H).
46				1	white solid	418.94	2.7 Method A	441 (M+Na <sup>+</sup> )	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.89 (d, 2H, J=8.2Hz), 7.65 (d, 2H, J=8.4Hz), 7.52 (s, br, 1H) 7.00 (s, br, 1H), 4.25 (dd, 2H, J=80Hz, 18Hz), 4.10-4.13 (m, 1H), 2.59-2.60 (m, 1H), 1.35 (s, 9H), 1.32-1.35 (m, 2H), 0.860 (d, 3H, J=6.0Hz), 0.75 (d, 3H, J=6.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
47				1	yellow solid	439.92	1.69 Method A	440.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.17 (d, 2H, J=7.0Hz), 7.72 (d, 2H, J=7.0Hz), 7.54 (d, 1H, J=8.8Hz), 7.49 (d, 1H, J=8.8Hz), 6.14 (s, br, 1H), 5.18 (s, br, 1H), 4.60 (dd, 2H, J=70Hz, 18Hz), 4.29-4.34 (m, 1H), 1.74-1.83 (m, 1H), 1.00-1.34 (m, 2H), 0.78 (d, 3H, J=7.0Hz), 0.67 (d, 3H, J=7.0Hz).
48				4	tan solid	409.94	1.26 Method A	410.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.80 (d, 2H, J=8.5Hz), 7.63 (d, 2H, J=8.5Hz), 7.52 (s, br, 1H), 7.46 (d, 1H, J=8.0Hz), 7.26 (d, 1H, J=8.0Hz), 7.02 (s, br, 1H), 4.70 (dd, 2H, J=50Hz, 18Hz), 4.30-4.41 (m, 1H), 3.67 (s, br, 2H), 1.28-1.33 (m, 3H), 0.86 (d, 3H, J=7.0Hz), 0.57 (d, 3H, J=7.0Hz).
49				3	Tan foam	431.99	1.19 Method A	432	<sup>1</sup> H NMR (d <sub>6</sub> DMSO) δ 7.81 (d, 2H, J=8.0Hz), 7.66 (d, 2H, J=8.0Hz), 7.45 (s, br, 1H), 7.02 (s, 1H), 4.27 (m, 1H), 3.56 (s, br, 4H), 3.55-3.57 (m, 1H), 3.08-3.14 (m, 1H), 2.22-2.32 (m, 6H), 1.79-1.82 (m, 11H), 1.17-1.62 (m, 4H), 0.86 (d, 6H, J=6.0Hz).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
50				1	white solid	436.51	1.58 Method A	437.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.95 (d, 2H, J=8.0Hz), 7.74-7.79 (m, 2H), 7.42 (d, 2H, J=8.0Hz), 7.14-7.19 (m, 2H), 6.24 (s, br, 1H), 5.20 (s, br, 1H), 4.50 (dd, 2H, J=50Hz, 17Hz), 4.12 (m, 1H), 3.91 (s, 3H), 1.75-1.82 (m, 1H), 1.25-1.31 (m, 1H), 1.05-1.12 (m, 1H), 0.75 (d, 3H, J=7.0Hz), 0.64 (d, 3H, J=7.0Hz)
51				4	tan solid	423.97	1.21 Method A	424.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.2Hz), 7.47 (d, 2H, J=8.0Hz), 7.31 (d, 2H, J=8.5Hz), 6.24 (s, br, 1H), 5.16 (s, br, 1H), 4.50 (dd, 2H, J=50Hz, 17Hz), 4.27 (t, 1H, J=10Hz), 2.44 (s, 3H), 1.74-1.83 (m, 1H), 1.25-1.33 (m, 1H), 0.93-1.01 (m, 1H), 0.74 (d, 3H, J=7.0Hz), 0.63 (d, 3H, J=7.0Hz)
52				1	white solid	448.90	1.79 Method A	449.0	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.66 (d, 2H, J=8.5Hz), 7.43 (d, 2H, J=8.5Hz), 7.12 (d, 1H, J=8.0Hz), 6.49 (d, 1H, J=8.0Hz), 6.24 (s, br, 1H), 5.22 (s, br, 1H), 4.35 (dd, 2H, J=50Hz, 15Hz), 4.22-4.27 (m, 1H), 2.04 (s, 3H), 1.27-1.89 (m, 3H), 0.74 (d, 3H, J=7.0Hz), 0.68 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
53				1-Method A	white solid	424.95	1.58 Method A	425.2	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.82 (d, 2H, J=8.2), 7.62 (d, 2H, J=8.3), 7.44 (d, 2H, J=7.1), 7.21 (t, 1H, J=6.7), 6.97 (s, 1H), 6.92 (d, 2H, J=6.9), 4.99 (d, 1H, J=17), 4.40 (d, 1H, J=17), 4.33 (br s, 1H), 3.76 (s, 3H), 1.20-1.40 (m, 3H), 0.79 (d, 3H, J=5.2), 0.54 (d, 3H, J=5.2)
54				1-Method A	white solid	400.86	1.81 min Method B	398.94 (M-H <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=6.8Hz), 7.48 (d, 2H, J=6.8Hz), 7.15 (d, 2H, J=10Hz), 7.02 (d, 2H, J=8.0Hz), 6.85 (t, 1H, J=7.5Hz), 6.19 (s, br, 1H), 5.13 (s, br, 1H), 4.31 (dd, 2H, J=50Hz, 15Hz), 4.43-4.45 (m, 1H), 3.87 (s, 3H), 1.17 (d, 3H, J=6.8Hz)
55				1-Method A	white solid	478.92	1.76 Method A	479.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.81 (d, 2H, J=8.5), 7.60 (d, 2H, J=8.2), 7.54 (s, 1H), 7.51 (d, 2H, J=8.5), 7.30 (d, 2H, J=8.2), 7.04 (s, 1H), 4.83 (d, 1H, J=17), 4.71 (d, 1H, J=17), 1.20-1.35 (m, 3H), 0.80 (d, 3H, J=6.1), 0.47 (d, 3H, J=6.2)
56				1-Method A	white solid	434.58	1.95 min Method A	435.24	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67 (dd, 2H, J=5.0, 8.9), 7.20-7.30 (m, 4H), 7.09 (dd, 2H, J=8.6, 8.6), 6.28 (br s, 1H), 5.24 (br s, 1H), 4.44 (s, 2H), 4.34 (t, 1H, J=7.8), 1.75-1.90 (m, 1H), 1.22-1.38 (m, 2H), 1.30 (s, 9H), 0.77 (d, 3H, J=6.4), 0.67 (d, 3H, J=6.4)
























Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
57				3	brown oil	458.07	2.16 Method C	458.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.77 (dd, 2H, J = 1.6, 6.8), 7.53 (dd, 2H, J = 2.0, 6.7), 6.55 (s, 1H), 5.64 (s, 1H), 4.15 (dd, 1H, J = 5.6, 9.2), 3.57 (t, 2H, J = 12), 3.35-3.45 (m, 1H), 3.10-3.17 (m, 1H), 2.87-3.00 (m, 2H), 2.57-2.70 (m, 2H), 2.27-2.40 (m, 2H), 1.80-2.00 (m, 6H), 1.63-1.73 (m, 2H), 1.20-1.50 (m, 4H), 0.85-0.90 (m, 1H), 0.73 (d, 3H, J = 6.4), 0.70 (d, 3H, J = 6.9).
58				5	colorless oil	390.89	1.51 Method A	391.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.91 (dd, 2H, J = 2.0, 6.8), 7.51 (dd, 2H, J = 2.2, 6.9), 7.06 (s, 1H), 5.33 (s, 1H), 4.13-4.25 (m, 2H), 4.03-4.15 (m, 3H), 1.77-1.85 (m, 1H), 1.35-1.45 (m, 1H), 1.30 (t, 3H, J = 7.1), 1.15-1.22 (m, 1H), 0.74 (d, 3H, J = 6.6), 0.71 (d, 3H, J = 6.5).
59				5	white solid	362.83	1.28 Method A	363.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.90 (dd, 2H, J = 2.0, 6.8), 7.65 (dd, 2H, J = 2.0, 6.8), 7.60 (s, 1H), 7.06 (s, 1H), 4.32 (d, 1H, J = 18), 4.12 (t, 1H, J = 8.0), 4.02 (d, 1H, J = 18), 1.55-1.65 (m, 1H), 1.35-1.45 (m, 2H), 0.78 (d, 3H, J = 6.1), 0.73 (d, 3H, J = 6.1).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
60				1-solid support	white solid	408.95	1.82 min Method B	409.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.64 (d, 2H, J=8.0Hz), 7.44 (d, 2H, J=8.0Hz), 7.22 (d, 2H, J=8.0Hz), 7.08 (d, 2H, J=8.0Hz), 6.29 (s, br, 1H), 5.34 (s, br, 1H), 4.53 (d, 1H, J=15.20Hz), 4.34 (d, 1H, J=15.20Hz), 4.27 (t, 1H, J=7.2Hz), 2.32 (s, 3H), 1.84 (m, 1H), 1.30 (m, 1H), 1.21 (m, 1H), 0.75 (d, 3H, J=6.8Hz), 0.67 (d, 3H, J=6.8Hz).
61				1-Method A	white solid	452.96	1.85 min Method A	453.08	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.97 (dd, 2H, J=1.7, 8.4), 7.68 (dd, 2H, J = 2.0, 8.7), 7.41-7.48 (m, 4H), 6.23 (br s, 1H), 5.16 (br s, 1H), 4.64 (d, 1H, J = 15.8), 4.47 (d, 1H, J = 15.9), 4.31 (t, 1H, J = 7.8), 3.92 (s, 3H), 1.76-1.83 (m, 1H), 1.26-1.35 (m, 1H), 1.08-1.13 (m, 1H), 0.76 (d, 3H, J = 6.0), 0.65 (d, 3H, J = 6.0).
62				1-solid support	white solid	430.95	1.81 min Method B	431.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.76 (dd, 2H, J=8.0Hz), 7.49 (d, 2H, J=8.0Hz), 7.24 (m, 1H), 6.95 (m, 2H), 6.25 (s, br, 1H), 5.27 (s, br, 1H), 4.62 (d, 1H, J=16.0Hz), 4.45 (d, 1H, J=16.0Hz), 4.33 (t, 1H, J=6.8Hz), 1.84 (m, 1H), 1.30 (m, 1H), 1.21 (m, 1H), 0.78 (d, 3H, J=6.8Hz), 0.70 (d, 3H, J=6.8Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/Method	M+H <sup>+</sup>	NMR Data
63				1-solid support	pale yellow solid	471.02	2.04 min Method B	471.09	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.37-7.44 (m, 6H), 6.27 (s, br, 1H), 5.33 (s, br, 1H), 4.58 (d, 1H, J=15.2Hz), 4.45 (d, 1H, J=15.2Hz), 4.36 (t, 1H, J=7.2Hz), 1.84 (m, 1H), 1.30 (m, 1H), 1.21 (m, 1H), 0.78 (d, 3H, J=6.8Hz), 0.70 (d, 3H, J=6.8Hz)
64				1-solid support	pale pink solid	372.92	1.82 min Method B	395.11 M+Na	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 6.75 (s, br, 1H), 6.70 (s, br, 1H), 5.04 (m, 1H), 4.32 (t, 1H, J=7.2Hz), 3.91 (d br, 2H, J=7.0Hz), 1.81 (m, 1H), 1.66 (s br, 6H), 1.30 (m, 1H), 1.21 (m, 1H), 0.77 (d, 3H, J=6.5Hz), 0.76 (d, 3H, J=6.5Hz)
65				4	white solid	437.99	1.39 Method A	438.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500 MHz) δ 7.74 (dd, 2H, J=1.9, 6.7), 7.5 (dd, 2H, J=1.9, 6.8), 7.43 (s, 1H), 7.16 (d, 2H, J=8.6), 7.01 (s, 1H), 6.61 (d, 2H, J=8.8), 4.59 (q, 2H, J=16, 25), 4.34 (dd, 1H, J=5.0, 9.3), 2.85 (s, 6H), 1.27-1.47 (m, 3H), 0.80 (d, 3H, J=5.9), 0.52 (d, 3H, J=6.1).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/Method	M+H <sup>+</sup>	NMR Data
66				1-Method A	white solid	473.01	1.58 Method A	437.0	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500 MHz) δ 7.87 (d, 2H, J=8.2), 7.83 (d, 2H, J=8.3), 7.64 (d, 2H, J=8.5), 7.63 (d, 2H, J=8.5), 7.59 (s, 1H), 7.08 (s, 1H), 4.92 (d, 1H, J=17), 4.76 (d, 1H, J=17), 4.39 (t, 1H, J=6.9), 3.35 (br s, 1H), 1.20-1.40 (m, 3H), 0.81 (d, 3H, J=6.2), 0.54 (d, 3H, J=6.2).
67				1-Method A	colorless oil	346.88	1.79 Method A	347.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500 MHz) δ 7.83 (d, 2H, J=8.8), 7.63 (d, 2H, J=8.6), 7.41 (s, 1H), 7.0 (s, 1H), 4.25 (dd, 1H, J=4.8, 8.9), 3.38-3.47 (m, 1H), 3.0-3.13 (m, 1H), 1.55-1.70 (m, 1H), 1.40-1.55 (m, 1H), 1.30-1.40 (m, 1H), 0.87 (t, 3H, J=7.6), 0.73 (d, 3H, J=6.4), 0.72 (d, 3H, J=6.7).
68				3	white oil	448.05	1.25 min Method C	448.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz), δ 7.76 (dd, 2H, J=2.0, 6.7), 7.50 (dd, 2H, J=2.0, 6.7), 6.51 (s, 1H), 5.45 (s, 1H), 4.20 (dd, 1H, J=6.4, 8.2), 3.35-3.45 (m, 1H), 3.20-3.30 (m, 1H), 2.68 (br s, 8H), 2.35 (br s, 2H), 1.70-1.90 (m, 3H), 1.20-1.40 (m, 1H), 0.95-1.05 (m, 1H), 0.75 (d, 3H, J=4.3), 0.73 (d, 3H, J=4.4).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
69				6	white solid	422.48	1.56 min Method A	423.14	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.89-7.93 (m, 4H), 7.84 (br s, 1H), 7.49 (d, 2H, J=8.2), 7.24-7.29 (m, 2H), 6.58 (br s, 1H), 5.12 (d, 1H, J=15.3), 4.23 (dd, 1H, J=4.6, 9.7), 4.05 (d, 1H, J=15.4), 2.04-2.14 (m, 1H), 1.23-1.32 (m, 1H), 0.79-0.88 (m, 1H), 0.72 (d, 3H, J=6.6), 0.67 (d, 3H, J=6.6).
70				1-Method A	white solid	466.89	1.77min Method B	467.03	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J=6.8Hz), 7.52-7.56 (m, 2H), 7.47 (d, 2H, J=6.8Hz), 7.11 (t, 1H, J=8.5Hz), 6.22 (s, br, 1H), 5.24 (s, br, 1H), 4.41 (dd, 2H, J=50Hz, 15Hz), 4.28 (t, 1H, 7.5Hz), 1.80-1.92 (m, 1H), 1.21-1.30 (m, 1H), 0.95-1.19 (m, 2H), 0.76 (t, 3H, J=7.0Hz).
71				1-Method A	white wax	398.89	1.80min Method B	388.0	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J=8.0Hz), 7.54 (d, 2H, J=8.0Hz), 7.42-7.49 (m, 4H), 6.20 (s, br, 1H), 5.21 (s, br, 1H), 4.54 (dd, 2H, J=50Hz, 15Hz), 4.25-4.29 (m, 1H), 1.82-1.95 (m, 1H), 1.26-1.33 (m, 1H), 0.98-1.12 (m, 2H), 0.75 (t, 3H, J=7.0Hz).
72				1-Method A	white solid	482.01	1.79min Method B	482.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.76 (d, 1H, J=7.5Hz), 7.63-7.67 (m, 4H), 7.43-7.50 (m, 8H), 6.24 (s, br, 1H), 5.28 (s, br, 1H), 4.53 (dd, 2H, J=50Hz, 15Hz), 4.27 (t, 1H, J=7.3Hz), 1.87-1.99 (m, 1H), 1.30-1.39 (m, 1H), 1.03-1.11 (m, 2H), 0.76 (t, 3H, J=8.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
73				1-Method A	pale yellow solid	426.90	1.86 min Method A	427.09	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.69 (ddd, 2H, J=2.0, 2.7, 8.7), 7.47 (dd, 2H, J=2.0, 8.7), 7.15 (dd, 1H, J=2.1, 12.0), 7.03 (d, 1H, J=8.4), 6.84 (t, 1H, J=8.5), 6.30 (br s, 1H), 5.25-5.30 (m, 2H), 4.96 (d, 1H, J=1.4), 4.87 (td, 1H, J=1.4, 9.9), 4.57 (d, 1H, J=15.4), 4.23-4.32 (m, 2H), 3.87 (s, 3H), 2.60-2.67 (m, 1H), 2.15-2.24 (m, 1H).
74				1-Method A	colorless oil	436.92	1.88 min Method A	437.09	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.97 (dd 2H, J=1.7, 8.3), 7.71 (dd, 2H, J=2.0, 8.7), 7.43-7.50 (m, 4H), 6.26 (br s, 1H), 5.22-5.35 (m, 1H), 5.18 (br s, 1H), 4.84-4.96 (m, 2H), 4.69 (d, 1H, J=15.8), 4.41 (d, 1H, J=15.8), 4.31 (t, 1H, J=7.5), 3.91 (s, 3H), 2.60-2.67 (m, 1H), 2.11-2.24 (m, 1H).
75				1-Method A	white solid	403.89	1.63 min Method A	404.03	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.71 (ddd, 2H, J=2.0, 2.6, 8.7), 7.60 (dd, 2H, J=1.9, 8.3), 7.49-7.52 (m, 4H), 6.21 (br s, 1H), 5.22-5.33 (m, 1H), 5.17 (br s, 1H), 4.88-4.98 (m, 2H), 4.71 (d, 1H, J=16.2), 4.40 (d, 1H, J=16.1), 4.32 (t, 1H, J=7.6), 2.54-2.63 (m, 1H), 2.09-2.19 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
76				1-Method A	white solid	446.88	2.04 min Method A	447.05	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67 (d, 2H, J = 8.6), 7.45-7.56 (m, 6H), 6.24 (br s, 1H), 5.25-5.39 (m, 1H), 5.19 (br s, 1H), 4.88-4.98 (m, 2H), 4.68 (d, 1H, J = 15.8), 4.42 (d, 1H, J = 15.81), 4.34 (t, 1H, J = 7.5), 2.58-2.68 (m, 1H), 2.13-2.23 (m, 1H).
77				1-Method A	white solid	405.91	1.51min Method B	406.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.70 (d, 2H, J=8.0Hz) 7.68 (d, 2H, J=8.0Hz), 7.47-7.51 (m, 4H), 6.15 (s, br, 1H), 5.16 (s, br, 1H), 4.53 (dd, 2H, J=50Hz, 15Hz), 4.21-4.26 (m, 1H), 1.82-1.87 (m, 1H), 1.20-1.25 (m, 1H), 0.97-1.09 (m, 2H), 0.74 (t, 3H, J=7.0Hz).
78				1-solid support	white foam	408.95	1.84min Method B	431.04 M+Na	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.83 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 7.34 - 7.45 (m, 5H), 5.85(s, br, 2H), 5.17 (q, 1H, 7.2Hz), 3.78 (dd 1H, J=8.4Hz, 4Hz), 2.36 (m, 1H), 1.62 (m, 1H), 1.50 (d, 3H, J=7.2Hz), 1.23 (m, 1H), 0.91 (d, 3H, J=6.6Hz), 0.84 (d, 3H, J=6.6Hz).
79				1-solid support	colorless syrup	408.95	1.89 Method B	431.04 M+Na	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 7.17 - 7.32 (m, 5H), 6.67 (s, br, 1H), 6.15 (s, br, 1H), 5.17 (q, 1H, 7.2Hz), 4.26 (dd 1H, J=6.6Hz), 3.48 (m, 1H), 3.37 (m, 1H), 2.97 (m, 1H), 2.90 (m, 1H), 1.92(m, 1H), 1.33(m, 1H), 1.10(m, 1H), 0.76 (d, 3H, J=6.6Hz), 0.75 (d, 3H, J=6.6Hz).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
80				1-solid support	yellow solid	430.36	1.64min Method B	430.02	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.34 (s, 1H), 7.67 (d, 1H, J=6.8Hz), 7.48 (d, 1H, 6.8Hz), 7.25 (d, 1H, J=6.8Hz), 6.1 (br. s, 1H), 5.29 (br. s, 1H), 4.59 (d, 1H, J=16Hz), 4.39 (d, 1H, J=16Hz), 1.8 (m, 1H), 1.32(m, 1H), 1.06(m, 1H), 0.78 (d, 3H, J=6.4Hz), 0.68 (d, 3H, J=6.4Hz).
81				1-solid support	white solid	435.39	1.87 Method B	456.92 M+Na	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.64 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 6.79 (d, 1H, J=3.7Hz), 6.72 (d, 1H, J=3.7Hz), 4.60 (d, 1H, J=21.9), 4.56(d, 1H, J=21.9Hz), 4.28(t, 1H, J=7.4Hz), 1.86(m, 1H), 1.33(m, 1H), 1.25(m, 1H), 0.76(d, 3H, J=6.5Hz), 0.73(d, 3H, J=6.5Hz).
82				1-Method A	white solid	420.96	1.97 Method A	421.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 7.77 (d, 2H, J = 8.6), 7.45 (d, 2H, J = 9.10), 7.26-7.35 (m, 5H), 6.54 (d, 1H, J = 16), 6.38 (br s, 1H), 6.00-6.07 (m, 1H), 5.34 (br s, 1H), 4.36 (t, 1H, J = 7.2), 4.02-4.15 (m, 2H), 1.83-1.92 (m, 1H), 1.35-1.43 (m, 1 H), 1.25-1.32 (m, 1 H), 0.79 (d, 3H, J = 6.7), 0.77 (d, 3H, J = 6.7).
83				5	white solid	389.90	1.91 Method D	390.2	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500 MHz) δ 8.08 (br s, 1H), 7.97 (d, 2H, J = 8.7), 7.65 (d, 2H, J = 8.5), 7.01 (br s, 1H), 4.46 (d, 1H, J = 18), 4.22 (d, 1H, J = 18), 3.91 (t, 1H, J = 6.3), 3.00 (s, 3H), 2.85 (s, 3H), 1.40-1.50 (m, 2H), 1.30-1.40 (m, 1 H), 0.85 (d, 3H, J = 6.1), 0.64 (d, 3H, J = 6.0).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
84				5	white solid	375.88	1.38 Method A	376.0	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 7.85 (dd, 2H, J = 1.8, 6.8), 7.50 (dd, 2H, J = 2.0, 6.8), 7.40 (br s, 1H), 6.37 (br s, 1H), 5.25 (br s, 1H), 4.26 (dd, 1H, J = 6.2, 8.6), 3.97 (d, 1H, J = 17), 3.85 (d, 1H, J = 17), 2.85 (d, 3H, J = 4.8), 1.75-1.85 (m, 1H), 1.40-1.48 (m, 1H), 0.88 (d, 3H, J = 6.4), 0.87 (d, 3H, J = 6.6).
85				5	white solid	448.01	2.18 Method C	448.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 8.10 (br s, 1H), 7.92 (d, 2H, J = 8.5), 7.50 (d, 2H, J = 8.5), 5.18 (br s, 1H), 4.35 (d, 1H, J = 17), 4.15 (t, 1H, J = 7.4), 3.97-4.05 (m, 1H), 3.95 (d, 1H, J = 17), 3.70-3.89 (m, 3H), 2.55-2.85 (m, 4H), 1.85-1.91 (m, 1H), 1.55-1.85 (m, 1H), 1.30-1.40 (m, 1H), 0.85 (d, 3H, J = 6.4), 0.83 (d, 3H, J = 6.4).
86				5	white solid	429.97	2.26 Method C	430.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 8.59 (br s, 1H), 7.92 (d, 2H, J = 9.1), 7.47 (d, 2H, J = 8.6), 5.16 (br s, 1H), 4.42 (d, 1H, J = 17), 4.23 (dd, 1H, J = 5.6, 8.6), 3.87 (d, 1H, J = 17), 3.55-3.65 (m, 1H), 3.40-3.52 (m, 3H), 1.80-2.00 (m, 1H), 1.45-1.80 (m, 7H), 1.35-1.45 (m, 1H), 0.89 (d, 3H, J = 6.7), 0.86 (d, 3H, J = 6.5).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
87				5	white solid	437.95	2.37 Method C	438.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 8.95 (br s, 1H), 7.83 (d, 2H, J = 8.6), 7.47 (d, 2H, J = 8.1), 7.43 (d, 2H, J = 8.6), 7.33 (t, 2H, J = 8.1), 7.13 (t, 1H, J = 7.6), 6.65 (br s, 1H), 5.45 (br s, 1H), 4.40 (dd, 1H, J = 6.1, 8.6), 4.07 (d, 1H, J = 17), 4.03 (d, 1H, J = 17), 1.70-1.80 (m, 1H), 1.55-1.65 (m, 2H), 0.93 (d, 3H, J = 7.0), 0.90 (d, 3H, J = 6.4).
88				5	white solid	401.92	1.94 Method C	402.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 7.85 (dd, 2H, J = 1.9, 8.9), 7.50 (dd, 2H, J = 2.0, 8.7), 7.40 (br s, 1H), 6.55 (br s, 1H), 6.30 (br s, 1H), 4.23 (dd, 1H, J = 2.9, 8.9), 3.92 (d, 1H, J = 17), 3.83 (d, 1H, J = 17), 2.68-2.73 (m, 1H), 1.75-1.83 (m, 1H), 1.50-1.57 (m, 1H), 1.40-1.49 (m, 1H), 0.88 (d, 3H, J = 6.4), 0.87 (d, 3H, J = 6.7), 0.80 (d, 2H, J = 7.0), 0.51 (t, 2H, J = 4.0).
89				6	white solid	438.93	1.67 min Method A	439.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300 MHz) δ 7.91 (d, 2H, J = 8.2), 7.81-7.84 (m, 3H), 7.56 (d, 2H, J = 8.6), 7.49 (d, 2H, J = 8.2), 6.55 (br s, 1H), 5.10 (d, 1H, J = 15.4), 4.23 (dd, 1H, J = 4.6, 9.7), 4.05 (d, 1H, J = 15.4), 2.04-2.14 (m, 1H), 1.20-1.31 (m, 1H), 0.80-0.89 (m, 1H), 0.74 (d, 3H, J = 6.6), 0.68 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
90				6	white solid	422.89	1.41 min Method A	423.05	<sup>1</sup> H NMR (dmso-d <sub>6</sub> , 300MHz) δ 7.86 (d, 2H, J = 8.2), 7.85 (br s, 1H), 7.81 (d, 2H, J = 8.6), 7.61 (d, 2H, J = 8.6), 7.47 (d, 2H, J = 8.0), 7.10 (br s, 1H), 5.45-5.55 (m, 1H), 4.83-4.95 (m, 3H), 4.71 (d, 1H, J = 17.0), 4.47 (t, 1H, J = 7.4), 2.29-2.37 (m, 1H), 2.13-2.22 (m, 1H).
91				2	tan solid	450.0	1.62min Method B	450.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J = 7.0Hz), 7.42 (d, 2H, J = 7.0Hz), 7.17-7.26 (m, 2H), 6.49-6.58 (m, 2H), 6.18 (s, br, 1H), 5.11 (s, br, 1H), 4.33 (dd, 2H, J = 50Hz, 15Hz), 4.12-4.20 (m, 1H), 3.21-3.30 (m, 4H), 1.91-2.04 (m, 5H), 1.32-1.38 (m, 1H), 0.94-1.09 (m, 2H), 0.75 (t, 3H, J = 8.0Hz).
92				7	white solid	502.08	1.72 Method A	502.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.86 (dd, 2H, J = 2.0, 6.8), 7.65 (dd, 2H, J = 2.0, 6.8), 7.37 (br s, 1H), 7.07 (br s, 1H), 4.19 (t, 1H, J = 7.6), 3.92 (br s, 2H), 3.35 (dd, 1H, J = 15, 6.8), 3.05 (dd, 1H, J = 15, 8.1), 1.85 (br s, 1H), 1.50-1.70 (m, 4H), 1.38 (s, 9H), 1.10-1.20 (m, 1H), 0.80-1.00 (m, 3H), 0.82 (d, 6H, J = 7.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
93				5	white solid	478.01	1.60 Method A	478.1	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 8.05 (s, 1H), 7.98 (d, 2H, J = 7.8), 7.65 (d, 2H, J = 7.8), 7.05 (s, 1H), 4.73 (s, 1H), 4.55-4.65 (m, 2H), 4.37 (t, 1H, J = 16), 3.95 (br s, 1H), 3.70 (br s, 2H), 2.90 (br s, 1H), 2.77 (br s, 1H), 1.00-1.55 (m, 4H), 0.70 (d, 6H, J = 4.1).
94				5	white solid	530.20	1.59 Method A	531.2	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.96 (d, 2H, J = 8.7), 7.65 (d, 2H, J = 8.6), 7.02 (s, 1H), 4.5 (d, 1H, J = 18), 4.27 (d, 2H, J = 18), 3.95 (br s, 1H), 3.35-3.50 (m, 8H), 1.30-1.55 (m, 3H), 1.41 (s, 9H), 0.74 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.0).
95				8	white solid	424.95	1.49 min Method A	425.17	<sup>1</sup> H NMR, 500Hz, (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J = 8.0Hz), 7.46 (d, 2H, J = 8.0Hz), 7.33 (d, 2H, J = 8.0Hz), 7.28 (d, 2H, J = 8.0Hz), 6.26 (s, br, 1H), 5.35 (s, br, 1H), 4.67 (s, br, 2H), 4.56 (d, 1H, J <sub>ax</sub> = 16Hz), 4.36 (d, 1H, J <sub>ax</sub> = 16Hz), 4.26 (t, 1H, J = 7.6Hz), 1.86-1.80 (m, 2H), 1.34-1.28 (m, 1H), 1.16-1.10 (m, 1H), 0.96 (d, 3H, J = 7.0Hz), 0.93 (d, 3H, J = 7.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
96				1-Method A	white solid	452.96	1.75 min Method A	453.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.96-7.90 (m, 2H), 7.64 (A of ABq, 2H, J=8.8Hz), 7.56 (d, 1H, J=7.5Hz), 7.43 (B of ABq, 2H, J=8.8Hz), 7.37 (t, 1H, J=7.5Hz), 6.28 (bs, 1H), 5.25 (bs, 1H), 4.61 (A of ABq, 1H, J=15.7Hz), 4.48 (B of ABq, 1H, J=15.7Hz), 4.36 (t, 1H, J=7.3Hz), 3.91 (s, 3H), 1.86-1.76 (m, 1H), 1.39-1.30 (m, 1H), 1.23-1.13 (m, 1H), 0.78 (d, 3H, J=6.6Hz), 0.68 (d, 3H, J=6.6Hz).
97				1-Method A	white solid	417.92	1.53 min Method A	418.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.71 (d, 2H, J=8.7), 7.60 (d, 2H, J=8.4), 7.47-7.51 (m, 4H), 6.28 (br s, 1H), 5.17 (br s, 1H), 4.73 (d, 1H, J=16.2), 4.41 (d, 1H, J=15.9), 4.32 (dd, 1H, J=1.5, 8.1), 1.65-1.82 (m, 1H), 1.14-1.30 (m, 1H), 0.27-0.40 (m, 2H), 0.12-0.22 (m, 1H), -0.18-0.05 (m, 2H).
98				1-Method A	white solid	460.91	1.76 min Method A	461.05	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67 (d, 2H, J=8.4), 7.44-7.56 (m, 6H), 6.32 (br s, 1H), 5.20 (br s, 1H), 4.70 (d, 1H, J=15.6), 4.43 (d, 1H, J=15.6), 4.35 (t, 1H, J=7.8), 1.70-1.84 (m, 1H), 1.26-1.32 (m, 1H), 0.32-0.40 (m, 2H), 0.16-0.24 (m, 1H), -0.14-0.00 (m, 2H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
99				1-Method A	white solid	450.94	1.63 min Method A	451.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.97 (d, 2H, J=8.3), 7.70 (d, 2H, J=8.7), 7.43-7.48 (m, 4H), 6.32 (br s, 1H), 5.15 (br s, 1H), 4.70 (d, 1H, J=15.8), 4.43 (d, 1H, J=15.8), 4.35 (t, 1H, J=7.8), 3.91 (s, 3H), 1.70-1.82 (m, 1H), 1.23-1.35 (m, 1H), 0.32-0.40 (m, 2H), 0.10-0.21 (m, 1H), -0.25- -0.05 (m, 2H).
100				1-Method A	white solid	509.06	1.94 min Method A	509.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.58 (A of ABq, 2H, J=8.8Hz), 7.40 (B of ABq, 2H, J=8.8Hz), 7.24 (bs, 4H), 6.20 (bs, 1H), 5.23 (bs, 1H), 4.45 (s, 2H), 4.36 (t, 1H, J=7.3Hz), 4.12 (q, 2H, J=7.2Hz), 1.83-1.74 (m, 1H), 1.55 (s, 3H), 1.55 (s, 3H), 1.39-1.20 (m, 2H), 1.19 (t, 3H, J=7.2Hz), 0.78 (d, 3H, J=6.6Hz), 0.66 (d, 3H, J=6.6Hz).
101				6	white solid	508.04	1.48 min Method A	508.22	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.68 (d, 2H, J=8.6), 7.29-7.47 (m, 6H), 6.38 (br s, 1H), 5.75 (br s, 1H), 4.65 (d, 1H, J=16.0), 4.42 (d, 1H, J=16.0), 4.32 (t, 1H, J=7.5), 3.30-3.85 (br m, 8H), 1.69-1.78 (m, 1H), 1.28-1.37 (m, 1H), 1.08-1.14 (m, 1H), 0.76 (d, 3H, J=6.5), 0.63 (d, 3H, J=6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
102				6	white solid	491.59	1.39 min Method A	492.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.75-7.80 (m, 2H), 7.42 (d, 2H, J = 8.2), 7.33 (d, 2H, J = 8.2), 7.14-7.20 (m, 2H), 6.37 (br s, 1H), 5.64 (br s, 1H), 4.64 (d, 1H, J = 16.0), 4.44 (d, 1H, J = 16.0), 4.31 (t, 1H, J = 7.1), 3.20-3.85 (br m, 8H), 1.70-1.78 (m, 1H), 1.28-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.76 (d, 3H, J = 6.5), 0.63 (d, 3H, J = 6.6).
103				6	colorless oil	551.11	1.32 min Method A	551.24	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67-7.72 (m, 4H), 7.42-7.48 (m, 4H), 6.82 (br s, 1H), 6.25 (br s, 1H), 5.31 (br s, 1H), 4.65 (d, 1H, J = 15.9), 4.43 (d, 1H, J = 15.9), 4.30 (t, 1H, J = 7.9), 3.70-3.79 (m, 4H), 3.53-3.59 (m, 2H), 2.53-2.65 (m, 6H), 1.79-1.86 (m, 1H), 1.29-1.38 (m, 1H), 1.08-1.14 (m, 1H), 0.76 (d, 3H, J = 6.5), 0.65 (d, 3H, J = 6.6).
104				6	white solid	534.66	1.22 min Method A	535.28	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.75-7.80 (m, 2H), 7.71 (d, 2H, J = 8.2), 7.43 (d, 2H, J = 8.1), 7.14-7.20 (m, 2H), 6.81 (br s, 1H), 6.28 (br s, 1H), 5.31 (br s, 1H), 4.64 (d, 1H, J = 15.9), 4.44 (d, 1H, J = 15.9), 4.29 (t, 1H, J = 7.6), 3.70-3.79 (m, 4H), 3.53-3.59 (m, 2H), 2.52-2.63 (m, 6H), 1.77-1.85 (m, 1H), 1.29-1.38 (m, 1H), 1.06-1.13 (m, 1H), 0.75 (d, 3H, J = 6.5), 0.65 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
105				6	white solid	607.17	1.70 min Method A	607.29	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.69 (d, 2H, J = 8.6), 7.30-7.47 (m, 6H), 6.38 (br s, 1H), 5.75 (br s, 1H), 4.65 (d, 1H, J = 16.0), 4.43 (d, 1H, J = 16.0), 4.33 (t, 1H, J = 7.5), 3.30-3.85 (br m, 8H), 1.72-1.79 (m, 1H), 1.45 (s, 9H), 1.24-1.39 (m, 1H), 1.05-1.18 (m, 1H), 0.76 (d, 3H, J = 6.5), 0.62 (d, 3H, J = 6.6).
106				6	white solid	554.11	1.75 min Method A	554.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.70 (ddd, 2H, J = 1.9, 2.4, 8.6), 7.47 (ddd, 2H, J = 2.0, 2.3, 8.7), 7.37-7.42 (m, 4H), 7.15-7.20 (m, 4H), 6.31 (br s, 1H), 5.60 (br s, 1H), 4.87 (br s, 1H), 4.65 (d, 1H, J = 15.9), 4.52 (br s, 1H), 4.47 (d, 1H, J = 15.9), 4.33 (t, 1H, J = 7.2), 4.05 (br s, 1H), 2.85-3.01 (br m, 2H), 1.85 (m, 1H), 1.30-1.42 (m, 1H), 1.08-1.18 (m, 1H), 0.79 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.6).
107				1-Method A	clear wax	456.97	1.60min Method B	455.1 (M-H <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.61 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 6.82-7.10 (m, 3H), 6.21 (s, br, 1H), 5.15 (s, br, 1H), 4.35 (dd, 2H, J=50Hz, 15Hz), 4.15-4.22 (m, 1H), 3.89 (s, br, 3H), 2.30-2.33 (m, 1H), 0.86-0.98 (m, 1H), 0.74 (s, 9H).



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
108				1-Method A	white solid	433.96	1.58min Method B	432.14 (M-H)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.0Hz), 7.45-7.49 (m, 4H), 6.19 (s, br, 1H), 5.15 (s, br, 1H), 4.55 (dd, 2H, J=50Hz, 15Hz), 4.20-4.24 (m, 1H), 2.25-2.31 (m, 1H), 0.84-0.88 (m, 1H), 0.74 (s, 9H).
109				1-Method A	white solid	476.95	1.62min Method B	475.12 (M-H)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.61 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 7.40-7.44 (m, 4H), 6.20 (s, br, 1H), 5.21 (s, br, 1H), 4.51 (dd, 2H, J=50Hz, 15Hz), 4.24-4.28 (m, 1H), 2.28-2.32 (m, 1H), 0.91-0.96 (m, 1H), 0.76 (s, 9H).
110				8	tan solid	452.02	1.25min Method A	452.23	<sup>1</sup> H NMR, 400MHz, (CDCl <sub>3</sub> ) δ 7.94 (d, 2H, J=8.0Hz), 7.74 (d, 2H, J=8.0Hz), 7.63 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.44 (d, 1H, J <sub>AB</sub> =16Hz), 3.28-3.23 (m, 3H), 2.17 (s, br, 6H), 1.95 (m, 1H), 1.55 (m, 2H), 0.96 (d, 3H, J=7.0Hz), 0.93 (d, 3H, J=7.0Hz).
111				5	white solid	535.97	2.65 min Method C	536.04	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.80 (dd, 2H, J=2.0, 9.0), 7.45 (dd, 2H, J=2.0, 9.0), 7.32 (dd, 2H, J=1.8, 9.0), 7.24 (br s, 1H), 7.17 (d, 2H, J=9.0), 5.50 (br s, 1H), 4.43 (qd, 2H, J=6.0, 15), 4.22 (t, 1H, J=7.0), 4.05 (d, 1H, J=17), 3.90 (d, 1H, J=17), 1.70-1.80 (m, 1H), 1.42-1.55 (m, 1H), 1.32-1.41 (m, 1H), 0.83 (d, 6H, J=7.9).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
112				5	white foam	487.93	1.52 min Method A	488.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.77 (d, 2H, J=8.3), 7.45 (d, 2H, J=9.0), 7.30 (br s, 1H), 7.05-7.10 (m, 1H), 6.90-7.05 (m, 2H), 5.53 (br s, 1H), 4.39-4.50 (m, 2H), 4.24 (t, 1H, J=7.1), 4.02 (d, 1H, J=17), 3.90 (d, 2H, J=17), 1.70-1.80 (m, 1H), 1.42-1.55 (m, 1H), 1.35-1.42 (m, 1H), 0.83 (d, 6H, J=7.7).
113				5	white oily solid	417.96	1.52 min Method A	418.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.92 (br s, 1H), 7.82 (d, 2H, J=8.2), 7.47 (d, 2H, J=8.2), 7.25 (br s, 1H), 6.23 (br s, 1H), 5.47 (br s, 1H), 4.25 (t, 1H, J=7.2), 3.91 (d, 1H, J=17), 3.75 (d, 1H, J=17), 1.75-1.82 (m, 1H), 1.50-1.62 (m, 1H), 1.38-1.50 (m, 1H), 1.35 (s, 9H), 0.89 (d, 3H, J=5.4), 0.87 (d, 3H, J=5.6).
114				5	white solid	458.02	1.62 min Method A	458.26	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.84 (dd, 2H, J=2.0, 8.8), 7.68 (br s, 1H), 7.47 (dd, 2H, 2.0, 8.3), 6.62 (br t, 1H, J=5.3), 5.45 (br s, 1H), 4.25 (dd, 1H, J=2.3, 6.1), 3.98 (d, 1H, J=17), 3.85 (d, 1H, J=17), 3.03-3.15 (m, 2H), 1.86-1.92 (m, 1H), 1.40-1.85 (m, 7H), 1.05-1.35 (m, 4H), 0.90-0.99 (m, 2H), 0.88 (d, 3H, J=6.6), 0.87 (d, 3H, J=6.4).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
115				5	white solid	452.96	1.14 min Method A	453.22	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.53 (d, 2H, J = 5.2), 8.04 (t, 1H, J = 5.2), 7.80 (dd, 2H, J = 1.8, 8.5), 7.46 (dd, 2H, J = 1.8, 8.7), 7.33 (br s, 1H), 7.29 (d, 2H, J = 5.4), 5.78 (br s, 1H), 4.47 (qd, 2H, J = 6.0, 16, ), 4.21 (t, 1H, J = 7.4), 4.07 (d, 1H, J = 17), 3.92 (d, 1H, J = 17), 1.67-1.77 (m, 1H), 1.30-1.47 (m, 2H), 0.81 (d, 3H, J = 6.5), 0.77 (d, 3H, J = 7.0).
116				5	white foam	419.93	1.29 Min Method A	420.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.87 (dd, 2H, J = 2.0, 8.5), 7.85 (br s, 1H), 7.49 (dd, 2H, J = 2.1, 9.0), 6.73 (br s, 1H), 5.55 (br s, 1H), 4.22 (dd, 1H, J = 6.1, 8.3), 4.02 (d, 1H, J = 17), 3.87 (d, 1H, J = 17), 3.40-3.50 (m, 4H), 3.36 (s, 3H), 1.77-1.86 (m, 1H), 1.46-1.57 (m, 1H), 1.30-1.41 (m, 1H), 0.84 (d, 3H, J = 6.7), 0.83 (d, 3H, J = 6.5).
117				5	white foam	475.01	1.16 min Method A	475.26	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.85 (dd, 2H, J = 2.0, 8.5), 7.64 (br s, 1H), 7.47 (dd, 2H, J = 1.5, 7.1), 6.87 (br s, 1H), 5.55 (br s, 1H), 4.22 (dd, 1H, J = 6.2, 7.9), 4.00 (d, 1H, J = 17), 3.87 (d, 1H, J = 17), 3.72 (t, 1H, J = 4.2), 3.30-3.45 (m, 2H), 2.45-2.55 (m, 6H), 1.75-1.85 (m, 1H), 1.50-1.63 (m, 1H), 1.30-1.41 (m, 1H), 0.86 (d, 6H, J = 9.0).







Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
118				1-Method A	white solid	459.79	1.62min Method B	461.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J=7.0Hz), 7.48 (d, 2H, J=7.0Hz), 7.41 (d, 2H, J=6.5Hz), 7.21 (d, 2H, J=6.5Hz), 6.21 (s, br, 1H), 5.20 (s, br, 1H), 4.43 (dd, 2H, J=50Hz, 15Hz), 4.12-4.24 (m, 1H), 1.88-1.90 (m, 1H), 1.24-1.29 (m, 1H), 0.98-1.08 (m, 2H), 0.74 (t, 3H, J=7.0Hz).
119				1-Method A	yellow solid	437.06	1.84 min Method F	(M+Na) <sup>+</sup> 459.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.83 (d, 2H, J=8.8), 7.80 (d, 2H, J=8.3), 7.64 (d, 2H, J=8.5), 7.59 (d, 2H, J=8.6), 7.48 (s, 1H), 7.15 (s, 1H), 4.79 (ABq, 2H, Δν=22.2, J <sub>AB</sub> =17.4), 4.44 (dd, 1H, J=8.0, 6.3), 2.21 (m, 2H), 1.84 (m, 1H), 1.81 (s, 3H), 1.53 (m, 1H).
120				1-Method A	white solid	480.06	2.08 min Method F	(M+Na) <sup>+</sup> 503.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.8), 7.68 (d, 2H, J=8.6), 7.61 (m, 4H), 7.48 (s, 1H), 7.16 (s, 1H), 4.80 (ABq, 2H, Δν=16.7, J <sub>AB</sub> =17.0), 4.45 (dd, 1H, J=8.2, 6.2), 2.22 (m, 2H), 1.82 (m, 1H), 1.78 (s, 3H), 1.61 (m, 1H).







Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
121				5	white solid	488.19	1.20 min Method A	489.26	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.95 (br s, 1H), 7.85 (dd, 2H, J = 2.5, 9.2), 7.70 (br s, 1H), 7.47 (dd, 2H, 2.0, 8.8), 5.42 (br s, 1H), 4.23 (dd, 1H, J = 6.3, 8.3), 3.94 (d, 1H, J = 17), 3.83 (d, 1H, J = 17), 3.73 (t, 4H, J = 4.7), 3.39-3.48 (m, 1H), 3.22-3.33 (m, 1H), 2.42-2.55 (m, 4H), 1.79-1.88 (m, 2H), 1.63-1.75 (m, 2H), 1.49-1.61 (m, 1H), 1.31-1.45 (m, 1H), 1.05-1.11 (m, 1H), 0.88 (d, 6H, J = 6.6).
122				5	white solid	432.16	1.59 min Method C	433.12	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.04 (br s, 1H), 7.88 (dd, 2H, J = 1.9, 6.9), 7.46 (dd, 2H, J = 1.8, 6.8), 6.93 (br s, 1H), 5.55 (br s, 1H), 4.20 (dd, 1H, J = 6.2, 8.3), 4.01 (d, 1H, J = 17), 3.80 (d, 1H, J = 17), 3.25-3.40 (m, 2H), 2.40-2.50 (m, 2H), 2.25 (s, 6H), 1.75-1.90 (m, 1H), 1.45-1.60 (m, 1H), 1.30-1.45 (m, 1H), 0.84 (d, 3H, J = 6.1), 0.82 (d, 3H, J = 6.4).
123				6	white solid	466.00	1.49 min Method A	466.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.68 (d, 4H, J = 8.6), 7.47 (ddd, 2H, J = 1.6, 2.4, 8.6), 7.41 (d, 2H, J = 8.2), 6.25 (br s, 1H), 6.12 (br s, 1H), 5.30 (br s, 1H), 4.63 (d, 1H, J = 15.8), 4.44 (d, 1H, J = 15.8), 4.30 (t, 1H, J = 6.8), 3.47-3.52 (m, 2H), 1.78-1.84 (m, 1H), 1.30-1.34 (m, 1H), 1.25 (t, 3H, J = 7.2), 1.08-1.13 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.65 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
124				6	colorless oil	546.05	1.65 min Method A	546.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.71 (d, 2H, J = 8.2), 7.67 (d, 2H, J = 8.6), 7.46 (d, 2H, J = 8.6), 7.41 (d, 2H, J = 8.0), 6.52 (br s, 1H), 6.24 (br s, 1H), 5.40 (br s, 1H), 4.63 (d, 1H, J = 15.9), 4.59 (d, 2H, J = 5.6), 4.42 (d, 1H, J = 15.9), 4.29 (t, 1H, J = 6.6), 1.78-1.84 (m, 1H), 1.29-1.34 (m, 1H), 1.25 (t, 3H, J = 7.2), 1.06-1.11 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).
125				6	colorless oil	494.06	1.67 min Method A	494.24	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.68 (dd, 2H, J = 1.7, 8.6), 7.63 (d, 2H, J = 8.2), 7.46 (d, 2H, J = 8.6), 7.39 (d, 2H, J = 8.2), 6.28 (br s, 1H), 5.94 (br s, 1H), 5.35 (br s, 1H), 4.63 (d, 1H, J = 15.8), 4.41 (d, 1H, J = 15.8), 4.29 (t, 1H, J = 6.6), 1.78-1.84 (m, 1H), 1.25 (t, 3H, J = 7.2), 1.06-1.11 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).
126				7	white solid	401.15	1.34 min Method A	402.15	<sup>1</sup> H NMR (DMSO-d <sub>6</sub> , 500MHz) δ 7.87 (d, 2H, J = 8.5), 7.66 (d, 2H, J = 8.6), 7.41 (s, 1H), 7.04 (s, 1H), 4.17 (t, 1H, J = 7.3), 3.40-3.50 (m, 1H), 3.20-3.25 (m, 1H), 3.03-3.10 (m, 1H), 2.65-2.80 (m, 2H), 1.85-2.00 (m, 1H), 1.20-1.85 (m, 2H), 1.45-1.60 (m, 1H), 1.30-1.40 (m, 1H), 1.10-1.30 (m, 4H), 0.75-0.90 (m, 1H), 0.82 (d, 3H, J = 7.3), 0.80 (d, 3H, J = 7.0).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
127				8	white solid	480.07	1.34 min Method A	480.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.60 (d, 2H, J=8.0Hz), 7.39-7.33 (m, 4H), 6.26 (s, br, 1H), 5.40 (s, br, 1H), 4.53 (d, 1H, J <sub>AB</sub> =16Hz), 4.42 (d, 1H, J <sub>AB</sub> =16Hz), 2.58 (q, 4H, J=8.0Hz), 1.94 (m, 1H), 1.59 (m, 2H), 1.06 (t, 6H, J=8.0Hz), 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
128				8	white solid	504.1	1.32 min Method A	504.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.73 (d, 2H, J=8.0Hz), 7.48 (d, 2H, J=8.0Hz), 7.35 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.35 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.44 (d, 1H, J <sub>AB</sub> =16Hz), 3.35 (s, 2H), 3.47-3.42 (m, 1H), 3.01 (s, br, 1H), 1.90 (m, 1H), 1.63 (m, 2H) 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
129				8	white solid	583.2	1.26 min Method A	583.40	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.37 (m, 4H), 7.21 (m, 5H), 6.35 (s, br, 1H), 5.87 (s, br, 1H), 4.72 (d, 1H, J <sub>AB</sub> =16Hz), 4.48 (d, 1H, J <sub>AB</sub> =16Hz), 3.55 (s, 3H), 3.52 (s, 3H), 3.74-3.43 (m, 1H), 2.45 (m, 8H), 1.59 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
130				8	amber glass	496.06	1.32 min Method A	496.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J=8.0Hz), 7.72 (d, 2H, J=8.0Hz), 7.40-7.36 (m, 4H), 6.35 (s, br, 1H), 5.37 (s, br, 1H), 4.59 (d, 1H, J <sub>AB</sub> =16Hz), 4.37 (d, 1H, J <sub>AB</sub> =16Hz), 3.7-3.5 (m, 4H), 3.48 (s, 3H), 3.46 (m, 1H), 2.23-2.1 (m, 4H), 1.85 (m, 1H), 1.55 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
131				8	amber oil	510.12	1.33 min Method A	510.23	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.65 (s, b, 4H), 7.37 (d, 2H, J=8.0Hz), 6.25 (s, b, 1H), 5.36 (s, b, 1H), 4.60 (d, 1H, J <sub>AB</sub> =16Hz), 4.38 (d, 1H, J <sub>AB</sub> =16Hz), 3.47-3.43 (m, 3H), 2.61 (m, 4H), 2.30 (m, 4H), 1.90 (m, 1H), 1.59 (m, 2H), 0.96 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
132				8	amber glass	535.15	1.24 min Method A	535.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.02 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.40 (d, 2H, J=8.0Hz), 7.36 (d, 2H, J=8.0Hz), 6.27 (s, b, 1H), 5.40 (s, b, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.44 (d, 1H, J <sub>AB</sub> =16Hz), 3.61 (s, 2H), 3.45 (m, 1H), 2.72-2.63 (m, 3H), 2.41 (s, 3H), 2.32 (s, 3H), 2.05 (t, 4H, J=12.0Hz), 1.90 (m, 3H), 1.74-1.55 (m, 3H), 0.96 (d, 3H, J=7.0Hz), 0.93 (d, 3H, J=7.0Hz)







Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
133				8	amber glass	540.13	1.28 min Method A	540.34	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.70 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.24-7.14 (m, 3H), 6.88 (m, 1H), 6.25 (s, br, 1H), 5.39 (s, br, 1H), 4.62 (d, 1H, J <sub>AB</sub> =16Hz), 4.40 (d, 1H, J <sub>AB</sub> =16Hz), 3.45 (s, 2H), 3.46 (m, 1H), 3.35-3.19 (m, 2H), 3.93 (m, 1H), 2.71 (m, 1H), 2.61 (m, 1H), 2.49-2.43 (m, 2H), 1.89 (m, 1H), 1.67-1.54 (m, 2H), 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
134				8	white solid	476.04	1.31 min Method A	476.17	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.79 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.85 (s, br, 1H), 4.60 (s, 2H), 4.76 (d, 1H, J <sub>AB</sub> =16Hz), 4.30 (d, 1H, J <sub>AB</sub> =16Hz), 3.72 (s, br, 2H), 3.46 (m, 1H), 2.40 (s, br, 3H), 2.20 (s, 1H), 1.90 (m, 1H), 1.63 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
135				8	white solid	532.14	1.34 min Method A	532.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 7.33 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.59 (d, 1H, J <sub>AB</sub> =16Hz), 4.37 (d, 1H, J <sub>AB</sub> =16Hz), 3.53 (s, 3H), 3.44 (m, 1H), 2.79 (t, 2H, J=8.0Hz), 2.62 (q, 2H, J=8.0Hz), 2.41 (t, 2H, J=8.0Hz), 2.25 (s, 6H), 1.95-1.85 (m, 1H), 1.67-1.54 (m, 2H), 1.07 (t, 3H, J=8.0Hz), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
136				8	white solid	537.17	1.24 min Method A	537.34	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.41-7.36 (m, 4H), 6.57 (s, br, 1H), 5.40 (s, br, 1H), 4.59 (d, 1H, J <sub>AB</sub> =16Hz), 4.37 (d, 1H, J <sub>AB</sub> =16Hz), 3.51 (s, 2H), 3.45 (m, 1H), 2.69 (t, 2H, J=8.0Hz), 2.62 (t, 2H, J=8.0Hz), 2.55 (q, 4H, J=8.0Hz), 2.35 (s, 3H), 1.89 (m, 1H), 1.60 (m, 2H), 1.00 (t, 6H, J=8.0Hz), 0.96 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
137				8	white solid	509.12	1.32 min Method A	509.33	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 8.01 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.40 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.58 (d, 1H, J <sub>ab</sub> =16Hz), 4.36 (d, 1H, J <sub>ab</sub> =16Hz), 3.51 (s, 2H), 3.45 (m, 1H), 2.66 (t, 2H, J=8.0Hz), 2.39 (t, 2H, J=8.0Hz), 2.35 (s, 3H), 2.25 (s, 6H), 1.90 (m, 1H), 1.59 (m, 2H), 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
138				8	clear glass	494.1	1.37 min Method A	494.27	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 d, 2H, J=8.0Hz), 7.74 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.26 (s, br, 1H), 5.39 (s, br, 1H), 4.55 (d, 1H, J <sub>ab</sub> =16Hz), 4.42 (d, 1H, J <sub>ab</sub> =16Hz), 3.70 (s, 3H), 3.46 (m, 1H), 2.38-2.35 (m, 5H), 1.91 (m, 2H), 1.60 (m, 2H), 0.96 (d, 3H, J=7.0Hz), 0.90 (d, 3H, J=7.0Hz)
139				8	white solid	494.1	1.33 min Method A	494.26	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.63 (d, 2H, J=8.0Hz), 7.73 (d, 2H, J=8.0Hz), 7.48 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.38 (s, br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =16Hz), 4.44 (d, 1H, J <sub>ab</sub> =16Hz), 3.47-3.40 (m, 3H), 2.12 (s, 3H), 1.88 (m, 1H), 1.60 (m, 2H), 1.03 (s, 9H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
140				8	amber glass	521.13	1.27 min Method A	521.31	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.39-7.36 (m, 4H), 6.30 (s, br, 1H), 5.35 (s, br, 1H), 4.59 (d, 1H, J <sub>ab</sub> =16Hz), 4.37 (d, 1H, J <sub>ab</sub> =16Hz), 3.55 (s, 2H), 3.45 (m, 1H), 2.84 (m, 4H), 2.48 (q, 2H, J=7.0Hz), 2.28 (m, 4H), 1.88 (m, 1H), 1.60 (m, 2H), 1.20 (t, 3H, J=7.0Hz), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
141				8	white solid	478.06	1.31 min Method A	478.22	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.48 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =16Hz), 4.43 (d, 1H, J <sub>ab</sub> =16Hz), 3.58 (s, br, 2H), 3.45 (m, 1H), 2.66 (m, 4H), 1.86 (m, 5H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
142				8	white foam	468.02	1.58 min Method A	468.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.76 (d, 2H, J=8.0Hz), 7.75 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.38 (s, br, 1H), 4.76 (d, 1H, J <sub>ab</sub> =16Hz), 4.30 (d, 1H, J <sub>ab</sub> =16Hz), 3.75 (s, 2H), 3.45 (m, 1H), 3.36 (s, 3H), 2.61 (s, 3H), 1.88 (m, 1H), 1.59 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
143				8	white solid	482.05	1.28 min Method A	482.24	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.72 (d, 2H, J=8.0Hz), 7.42-7.36 (m, 4H), 6.35 (s, br, 1H), 5.83 (s, br, 1H), 4.58 (d, 1H, J <sub>ab</sub> =16Hz), 4.39 (d, 1H, J <sub>ab</sub> =16Hz), 3.52 (s, 1H), 3.50 (s, 2H), 3.48 (s, 1H), 3.45 (m, 1H), 2.55 (t, 2H, J=8.0Hz), 2.20 (s, 3H), 1.90 (m, 1H), 1.59 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
144				8	clear glass	512.07	1.22 min Method A	512.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=8.0Hz), 7.74 (d, 2H, J=8.0Hz), 7.39 (d, 2H, J=8.0Hz), 7.31 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.72 (d, 1H, J <sub>ab</sub> =16Hz), 4.39 (d, 1H, J <sub>ab</sub> =16Hz), 3.76 (t, 4H, J=8.0Hz), 3.45 (m, 1H), 3.38 (s, br, 2H), 3.15 (s, br, 2H), 3.76 (t, 4H, J=8.0Hz), 1.89 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
145				8	white solid	492.09	1.31 min Method A	492.21	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.63 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.26 (s, br, 1H), 5.39 (s, br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =16Hz), 4.43 (d, 1H, J <sub>ab</sub> =16Hz), 3.47-3.43 (m, 3H), 2.26 (m, 4H), 1.89 (m, 2H), 1.60 (m, 2H), 1.46-1.29 (m, 4H), 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
146				8	white solid	564.15	1.33 min Method A	564.24	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.48 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.40 (s, br, 1H), 4.72 (d, 1H, J <sub>ab</sub> =16Hz), 4.48 (d, 1H, J <sub>ab</sub> =16Hz), 4.03 (q, 2H, J=8.0Hz), 3.47-3.42 (m, 3H), 3.21 (m, 1H), 2.62 (m, 1H), 2.63-2.36 (m, 3H), 2.06-1.95 (m, 1H), 1.93-1.80 (m, 3H), 1.67-1.50 (m, 3H), 1.13 (t, 3H, J=8.0Hz), 0.98 (d, 3H, J=0.7Hz), 0.95 (d, 3H, J=7.0Hz)
147				8	white solid	582.21	1.46 min Method A	582.41	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.24 (t, 1H, J=8.0Hz), 7.03 (t, 2H, J=8.0Hz), 6.95 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =16Hz), 4.44 (d, 1H, J <sub>ab</sub> =16Hz), 3.48-3.43 (m, 3H), 2.76 (m, 2H), 2.61 (m, 2H), 1.88 (m, 3H), 1.69-1.54 (m, 6H), 1.26 (m, 1H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
148				8	white foam	570.16	1.41 min Method A	570.34	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.09 (d, 1H, J=4.0Hz), 8.02 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.41-7.36 (m, 5H), 6.72 (d, 1H, J=12.0Hz), 6.49 (t, 1H, J=8.0Hz), 6.35 (s, br, 1H), 5.87 (s, br, 1H), 4.80 (d, 1H, J <sub>AB</sub> =16Hz), 4.30 (d, 1H, J <sub>AB</sub> =16Hz), 3.84 (m, 4H), 3.55 (s, 2H), 3.46 (m, 1H), 2.57 (m, 4H), 1.89 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 2H, J=7.0Hz)
149				8	white solid	542.15	1.45 min Method A	542.24	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.70 (d, 2H, J=8.0Hz), 7.68 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.29-7.16 (m, 5H), 6.28 (s, br, 1H), 5.38 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.42 (d, 1H, J <sub>AB</sub> =16Hz), 3.54 (s, 2H), 3.45 (m, 1H), 2.62 (q, 2H, J=7.0Hz), 1.89 (m, 1H), 1.60 (m, 2H), 1.08 (t, 3H, J=7.0Hz), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)
150				8	amber film	528.12	1.45 min Method A	528.33	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.68 (d, 2H, J=8.0Hz), 7.54 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 7.35-7.29 (m, 5H), 6.27 (s, br, 1H), 5.37 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.43 (d, 1H, J <sub>AB</sub> =16Hz), 3.55 (s, 2H), 3.45 (m, 1H), 2.17 (s, 3H), 1.89 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.95 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
151				8	clear glass	542.15	1.46 min Method A	542.29	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.45 (d, 2H, J=8.0Hz), 7.39-7.26 (m, 4H), 6.97 (d, 1H, J=8.0Hz), 6.27 (s, br, 1H), 5.38 (s, br, 1H), 4.54 (d, 1H, J <sub>AB</sub> =16Hz), 4.44 (d, 1H, J <sub>AB</sub> =16Hz), 3.43 (s, 2H), 3.25 (t, 1H, J=4.0Hz), 2.77-2.69 (m, 4H), 2.40 (s, 3H), 1.94 (m, 1H), 1.60 (m, 2H), 0.97 (d, 2H, J=7.0Hz), 0.94 (d, 2H, J=7.0Hz)
152				8	amber glass	564.2	1.22 min Method A	564.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.59 (d, 2H, J=8.0Hz), 7.39-7.33 (m, 4H), 6.21 (s, br, 1H), 5.34 (s, br, 1H), 4.76 (d, 1H, J <sub>AB</sub> =16Hz), 4.31 (d, 1H, J <sub>AB</sub> =16Hz), 3.57 (s, 2H), 2.69-2.55 (m, 6H), 2.36-2.18 (m, 12H), 1.95 (m, 1H), 1.66-1.50 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
153				5	white solid	474.21	1.78 min Method C	476.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400 MHz) δ 8.41 (br s, 2H), 7.85 (dd, 2H, J = 2.0, 6.8), 7.46 (dd, 2H, J = 2.0, 6.8), 5.46 (s, 1H), 4.21 (dd, 1H, J = 5.9, 8.8), 3.92 (d, 1H, J = 17), 3.77 (d, 1H, J = 17), 3.27-3.41 (m, 2H), 2.48-2.62 (m, 6H), 1.80-1.90 (m, 1H), 1.50-1.72 (m, 3H), 1.35-1.49 (m, 1H), 1.05 (t, 6H, J = 7.1), 0.87 (d, 3H, J = 6.4), 0.85 (d, 3H, J = 6.7)















Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
154				5	white solid	516.18	1.53 min Method A	517.30	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500 MHz) δ 7.83 (dd, 2H, J = 1.9, 6.8), 7.47 (dd, 2H, J = 2.0, 6.8), 7.31 (br s, 1H), 7.11 (br s, 1H), 5.57 (br s, 1H), 4.23 (dd, 1H, J = 6.8, 8.4), 4.00-4.15 (m, 2H), 3.95 (d, 1H, J = 1.7), 3.83 (d, 1H, J = 1.7), 2.93 (br s, 2H), 1.75-1.95 (m, 4H), 1.60-1.75 (1H, 1.25-1.55 (m, 1H), 1.25 (t, 5H, J = 7.7), 0.85 (d, 3H, J = 6.3), 0.83 (d, 3H, J = 6.3).
155				1-Method A	white solid	495.04	1.88 min Method A	495.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.62-7.59 (m, 2H), 7.43-7.39 (m, 2H), 7.28-7.25 (m, 2H), 7.22-7.19 (m, 2H), 6.22 (bs, 1H), 5.29 (bs, 1H), 4.53-4.43 (m, 2H), 4.42-4.37 (m, 1H), 4.16-4.07 (m, 2H), 1.82-1.73 (m, 1H), 1.48 (d, 3H, J = 7.3 Hz, isomer A), 1.47 (d, 3H, J = 7.3 Hz, isomer B), 1.36-1.22 (m, 2H), 1.21 (t, 3H, J = 7.1 Hz), 0.77 (d, 3H, J = 6.4 Hz, isomer A), 0.77 (d, 3H, J = 6.4 Hz, isomer B), 0.65 (d, 3H, J = 6.7 Hz, isomer A), 0.65 (d, 3H, J = 6.7 Hz, isomer B).
156				1-Method A	white solid	434.07	1.57 min Method B	(M+H) <sup>+</sup> 435.1	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.80 (d, 2H, J = 8.7), 7.66 (d, 2H, J = 8.3), 7.60 (m, 4H), 7.53 (s, 1H), 7.10 (s, 1H), 4.80 (ABq, 2H, Δν = 39.8, J <sub>AB</sub> = 17.2), 4.30 (t, 1H, J = 7.5), 1.60 (m, 1H), 1.39 (m, 1H), 0.71 (t, 3H, J = 7.3).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
157				1-Method A	white solid	391.08	1.32 min Method B	(M+H) <sup>+</sup> 392.1	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.81 (d, 2H, J = 8.7), 7.78 (d, 2H, J = 8.4), 7.62 (d, 2H, J = 8.7), 7.57 (d, 2H, J = 8.3), 7.52 (s, 1H), 7.09 (s, 1H), 4.80 (ABq, 2H, Δν = 45.0, J <sub>AB</sub> = 17.6), 4.28 (t, 1H, J = 7.5), 1.58 (m, 1H), 1.36 (m, 1H), 0.70 (t, 3H, J = 7.3).
158				6	white solid	478.01	1.53 min Method A	478.17	<sup>1</sup> H NMR (CD <sub>3</sub> OD, 300 MHz) δ 7.80 (ddd, 2H, J = 1.9, 2.4, 8.7), 7.73 (d, 2H, J = 8.3), 7.48-7.54 (m, 4H), 4.87 (m, 1H), 4.79 (d, 1H, J = 16.0), 4.49 (t, 1H, J = 6.2), 2.81-2.87 (m, 1H), 1.25-1.43 (m, 3H), 0.83 (d, 3H, J = 6.2), 0.77-0.83 (m, 2H), 0.63-0.66 (m, 2H), 0.57 (d, 3H, J = 6.1).
159				6	white solid	496.03	1.50 min Method A	496.21	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300 MHz) δ 7.70 (d, 2H, J = 8.0), 7.68 (d, 2H, J = 8.6), 7.46 (d, 2H, J = 8.6), 7.41 (d, 2H, J = 8.1), 4.64 (d, 1H, J = 15.9), 4.43 (d, 1H, J = 15.9), 4.30 (t, 1H, J = 6.8), 3.63-3.66 (m, 2H), 3.55-3.58 (m, 2H), 3.39 (s, 3H), 1.76-1.84 (m, 1H), 1.28-1.34 (m, 1H), 1.05-1.11 (m, 1H), 0.75 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.7).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
160				6	white solid	551.15	1.33 min Method A	551.28	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.2), 7.67 (dd, 2H, J = 2.0, 8.7), 7.44 (dd, 2H, J = 1.8, 8.6), 7.39 (d, 2H, J = 8.2), 6.35 (br s, 1H), 5.46 (br s, 1H), 4.60 (d, 1H, J = 15.9), 4.50 (d, 1H, J = 15.9), 4.31 (t, 1H, J = 7.3), 3.54-3.57 (m, 2H), 2.56-2.64 (m, 6H), 1.72-1.80 (m, 3H), 1.28-1.34 (m, 1H), 1.13-1.16 (m, 1H), 1.03 (t, 6H, J = 7.2), 0.74 (d, 3H, J = 6.6), 0.61 (d, 3H, J = 6.6).
161				9	white solid	494.06	1.51 min Method B	494.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J = 7.0Hz), 7.45-7.47 (m, 4H), 7.30 (d, 2H, J = 8.0Hz), 7.12 (s, br, 1H), 6.25 (s, br, 1H), 5.22 (s, br, 1H), 4.40 (dd, 2H, J = 50Hz, 15Hz), 4.25 (t, 1H, J = 7.4Hz), 2.48-2.51 (m, 1H), 1.54-1.86 (m, 1H), 1.17-1.34 (m, 10H), 0.75 (d, 3H, J = 7.0Hz), 0.67 (d, 3H, J = 7.0Hz).
162				9	tan solid	504.01	1.52 min Method B	504.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.10 (s, br, 1H), 7.67 (d, 2H, J = 7.0Hz), 7.58 (d, 2H, J = 7.0Hz), 7.23-7.49 (m, 6H), 6.54-6.57 (m, 1H), 6.27 (s, br, 1H), 5.50 (s, br, 1H), 4.51 (dd, 2H, J = 50Hz, 15Hz), 4.28 (t, 1H, J = 7.4Hz), 1.78-1.85 (m, 1H), 1.12-1.32 (m, 2H), 0.75 (d, 3H, J = 7.0Hz), 0.67 (d, 3H, J = 7.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
163				9	white solid	451.98	1.50 min Method B	450.18 (M-H)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J = 8.0Hz), 7.28-7.46 (m, 6H), 7.12 (s, br, 1H), 6.24 (s, br, 1H), 5.19 (s, br, 1H), 4.48 (dd, 2H, J = 50Hz, 15Hz), 4.27 (t, 1H, J = 7.0Hz), 2.18 (s, 3H), 1.80-2.01 (m, 1H), 1.12-1.32 (m, 2H), 0.75 (d, 3H, J = 7.0Hz), 0.67 (d, 3H, J = 7.0Hz).
164				8	white solid	491.06	1.31 min Method A	491.24	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J = 8.0Hz), 7.64 (d, 2H, J = 8.0Hz), 7.51 (d, 2H, J = 8.0Hz), 7.37 (d, 2H, J = 8.0Hz), 6.25 (s, br, 1H), 5.35 (s, br, 1H), 4.59 (d, 1H, J <sub>AB</sub> = 16Hz), 4.36 (d, 1H, J <sub>AB</sub> = 16Hz), 3.62 (s, 2H), 3.25 (t, 1H, J = 6.0Hz), 2.55-2.48 (m, 7H), 1.94 (m, 1H), 1.60 (m, 2H), 0.97 (d, 3H, J = 7.0Hz), 0.94 (d, 3H, J = 7.0Hz).
165				8	white solid	520.14	1.40 min Method A	520.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J = 8.0Hz), 7.66 (d, 2H, J = 8.0Hz), 7.51 (d, 2H, J = 8.0Hz), 7.37 (d, 2H, J = 8.0Hz), 6.28 (s, br, 1H), 5.39 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> = 16Hz), 4.43 (d, 1H, J <sub>AB</sub> = 16Hz), 3.57 (s, 2H), 3.25 (t, 1H, J = 6.0Hz), 2.59 (t, 2H, J = 8.0Hz), 2.30 (d, 2H, J = 6.0Hz), 2.01-1.90 (m, 1H), 1.68-1.49 (m, 4H), 1.02 (m, 1H), 0.98 (d, 3H, J = 7.0Hz), 0.94 (d, 3H, J = 7.0Hz), 0.89 (t, 3H, J = 8.0Hz), 0.44-0.31 (m, 4H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
166				8	clear glass	506.11	1.41 min Method A	506.31	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.73 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.44 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.56 (d, 1H, J <sub>ab</sub> =16Hz), 4.42 (d, 1H, J <sub>ab</sub> =16Hz), 3.45 (d, 1H, J <sub>ab</sub> =12Hz), 3.28 (d, 1H, J <sub>ab</sub> =12Hz), 3.25 (t, 1H, J=6.0Hz), 2.75 (m, 4H), 2.0-1.53 (m, 7H), 1.35 (m, 1H), 1.17 (m, 1H), 0.97 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz), 0.87 (m, 3H)
167				8	clear glass	506.11	1.42 min Method A	506.30	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.26 (s, br, 1H), 5.40 (s, br, 1H), 4.56 (d, 1H, J <sub>ab</sub> =16Hz), 4.42 (d, 1H, J <sub>ab</sub> =16Hz), 3.48 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.56-2.42 (m, 4H), 1.93 (m, 1H), 1.89-1.48 (m, 6H), 1.34 (m, 1H), 0.98-0.93 (m, 9H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
168				8	clear glass	546.18	1.46 min Method A	546.35	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.42 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =16Hz), 4.43 (d, 1H, J <sub>ab</sub> =16Hz), 3.45 (d, 1H, J <sub>ab</sub> =12Hz), 3.28 (d, 1H, J <sub>ab</sub> =12Hz), 3.25 (t, 1H, J=6.0Hz), 2.87 (m, 1H), 2.70-2.56 (m, 3H), 2.39 (m, 1H), 2.31-2.19 (m, 4H), 1.95 (m, 1H), 1.77 (m, 1H), 1.68-1.59 (m, 2H), 1.54-1.44 (m, 4H), 1.28 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 2H, J=7.0Hz)
169				8	clear glass	508.13	1.39 min Method A	502.28	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.70 (d, 2H, J=8.0Hz), 7.68 (d, 2H, J=8.0Hz), 7.55 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.28 (s, br, 1H), 5.40 (s, br, 1H), 4.57 (d, 1H, J <sub>ab</sub> =16Hz), 4.36 (d, 1H, J <sub>ab</sub> =16Hz), 3.38 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.83 (m, 2H), 1.92 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz), 0.84 (s, 6H), 0.82 (s, 6H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
170				8	clear glass	494.1	1.34 min Method A	494.26	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.67 (d, 2H, J=8.0Hz), 7.44 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.44 (d, 1H, J <sub>AB</sub> =16Hz), 3.40 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.39 (s, 3H), 1.94 (m, 1H), 1.68-1.50 (m, 2H), 1.43 (m, 2H), 1.30 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz), 0.86 (t, 3H, J=7.0Hz)
171				8	clear glass	508.13	1.40 min Method A	508.27	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.4-7.33 (m, 4H), 6.25 (s, br, 1H), 5.35 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =16Hz), 4.43 (d, 1H, J <sub>AB</sub> =16Hz), 3.46 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.62 (t, 2H, J=6.0Hz), 2.52 (q, 2H, J=7.0Hz), 1.94 (m, 1H), 1.68-1.45 (m, 4H), 1.29 (m, 2H), 1.02 (t, 3H, J=7.0Hz), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz), 0.88 (t, 3H, J=7.0Hz)
172				8	amber glass	575.22	1.34 min Method A	575.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.59 (d, 2H, J=8.0Hz), 7.39-7.33 (m, 4H), 6.25 (s, br, 1H), 5.36 (s, br, 1H), 4.53 (d, 1H, J <sub>AB</sub> =16Hz), 4.39 (d, 1H, J <sub>AB</sub> =16Hz), 3.57 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.62-2.32 (m, 9H), 1.94 (m, 1H), 1.68-1.50 (m, 2H), 1.48-1.20 (m, 12H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
173				6	white solid	436.92	1.52 min Method A	437.16	<sup>1</sup> H NMR (dmsd-d <sub>6</sub> , 300MHz) δ 7.85 (d, 2H, J=8.4), 7.82 (dd, 2H, J=1.8, 8.7), 7.61 (dd, 2H, J=1.8, 8.7), 7.54 (br s, 1H), 7.47 (d, 2H, J=8.4), 7.09 (br s, 1H), 4.85 (d, 1H, J=17.4), 4.69 (d, 1H, J=17.1), 4.42 (t, 1H, J=7.2), 1.40-1.48 (m, 1H), 1.27-1.34 (m, 1H), 0.42-0.47 (m, 1H), 0.25-0.30 (m, 2H), 0.00-0.03 (m, 1H), -0.10- -0.07 (m, 1H)
174				1-Method A	off-white solid	492.11	2.13 min Method D	M+Na 514.95	<sup>1</sup> H NMR (DMSO) δ 7.76 (d, 2H, J=6.8Hz), 7.61 (m, 6H), 7.44 (s, br, 1H), 7.13 (s, br, 1H), 4.78 (dd, 2H, J=5.2Hz, 16Hz), 4.58 (t, 1H, J=8.0Hz), 3.47 (d, 2H, J=6.0Hz), 0.88 (s, 9H)
175				1-Method A	white solid	449.12	1.94 min Method D	M+Na 471.97	<sup>1</sup> H NMR (DMSO) δ 7.77 (m, 4H), 7.60 (m, 4H), 7.45 (s, br, 1H), 7.12 (s, br, 1H), 4.78 (dd, 2H, J=5.6Hz, 20Hz), 4.57 (t, 1H, J=8.0Hz), 3.47 (d, 2H, J=6.0Hz), 0.88 (s, 9H)
176				1-solid support		478.90	1.86 Method B	479.02	
177				1-solid support		426.90	1.82 Method B	449.02 M+Na	

- 160 -

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
178				1-solid support		496.00	1.81 Method B	496.06	
179				1-solid support		412.90	1.72 Method B	413.04	
180				1-solid support		423.00	1.86 Method B	445.02 M+Na	
181				1-solid support		501.10	1.94 Method B	523.04 M+Na	
182				1-solid support		413.90	1.53 Method B	414.05	
183				1-solid support		423.00	1.88 Method B	423.08	
184				1-solid support		467.00	1.60 Method B	467.06	
185				1-solid support		505.00	1.89 Method B	505.07	

- 161 -

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
186				1-solid support		478.90	1.84 Method B	479.02	
187				1-solid support		505.00	1.93 Method B	505.06	
188				1-solid support		429.40	1.80 Method B	450.90 M+Na	
189				1-solid support		423.00	1.89 Method B	423.09	
190				1-solid support		425.00	1.92 Method B	425.11	
191				1-solid support		487.00	1.91 Method B	487.04	
192				1-solid support		437.00	1.95 Method B	459.05 M+Na	
193				1-solid support		358.90	1.67 Method B	381.07 M+Na	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
194				1-solid support		473.80	1.86 Method B	472.9	
195				1-solid support		430.9	1.75 Method B	431.04	
196				1-solid support		379.30	1.65 Method B	400.98	
197				1-solid support		421.00	1.82 Method B	443.06 M+Na	
198				1-solid support		379.30	1.64 Method B	400.99	
199				1-solid support		495.00	1.95 Method B	494.98	
200				1-solid support		465.10	2.05 Method B	487.10 M+Na	
201				1-solid support		483.00	1.72 Method B	483.04	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
202				1-solid support		463.80	1.91 Method B	486.96 M+Na	
203				1-solid support		430.90	1.77 Method B	431.04	
204				1-solid support		409.00	1.79 Method B	409.07	
205				1-solid support		462.90	1.81 Method B	463.04	
206				1-solid support		423.00	1.86 Method B	423.10	
207				1-solid support		491.80	1.88 Method B	492.91	
208				1-solid support		409.00	1.78 Method B	431.04 M+Na	
209				1-solid support		419.9	1.58 Method B	442.04 M+Na	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
210				1-solid support		344.90	1.56 Method B	367.05 M+Na	
211				1-solid support		480.90	1.87 Method B	481.02	
212				1-solid support		430.90	1.76 Method B	453.02	
213				1-solid support		455.00	1.71 Method B	455.07	
214				1-solid support		515.00	1.91 Method B	515.09	
215				1-solid support		447.40	1.82 Method B	468.99 M+Na	
216				1-solid support		480.90	1.80 Method B	481.00	
217				1-solid support		402.90	1.600 Method B	403.12	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
218				1-solid support		429.40	1.78 Method B	429.04	
219				1-solid support		448.90	1.78 Method B	471.00	
220				1-solid support		430.90	1.75 Method B	453.03	
221				1-solid support		480.90	1.85 Method B	503.00 M+Na	
222				1-solid support		445.00	1.87 Method B	467.06 M+Na	
223				1-solid support		453.00	1.62 Method B	453.03	
224				1-solid support		453.00	1.63 Method B	453.05	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
225				1-solid support		416.00	1.53 Method B	416.04	
226				1-solid support		401.90	1.45 Method B	401.96	
227				1-solid support		395.90	1.12 Method B	396.01	
228				1-solid support		358.90	1.62 Method B	381.01 M+Na	
229				1-solid support		439.00	1.80 Method B	460.97 M+Na	
230				1-solid support		424.90	1.72 Method B	425.03	
231				1-solid support		463.80	1.85 Method B	464.90	
232				1-solid support		456.90	1.64 Method B	456.02	

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
233				1-solid support		447.40	1.78 Method B	468.92 M+Na	
234				1-Method A	white solid	419.11	1.91 min Method F	(M+Na) <sup>+</sup> 442.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.7), 7.80 (d, 2H, J=8.4), 7.62 (d, 2H, J=8.7), 7.59 (d, 2H, J=8.3), 7.51 (s, 1H), 7.07 (s, 1H), 4.81 (ABq, 2H, Δν=38.0, J <sub>AB</sub> =17.5), 4.31 (t, 1H, J=6.7), 1.54 (m, 1H), 1.29 (m, 1H), 1.03 (m, 3H), 0.85 (m, 1H), 0.66 (t, 3H, J=6.9).
235				1-Method A	white solid	462.10	2.13 min Method F	(M+Na) <sup>+</sup> 485.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.81 (d, 2H, J=8.7), 7.69 (d, 2H, J=8.3), 7.61 (m, 4H), 7.51 (s, 1H), 7.07 (s, 1H), 4.82 (ABq, 2H, Δν=31.9, J <sub>AB</sub> =17.1), 4.32 (t, 1H, J=8.3), 1.53 (m, 1H), 1.31 (m, 1H), 1.05 (m, 3H), 0.88 (m, 1H), 0.63 (t, 3H, J=6.8).
236				1-solid support	white solid	530.92	1.92 Method B	530.99	
237				1-solid support	white solid	416.93	1.61 Method B	417.07	
238				1-solid support	white solid	466.99	1.62 Method B	467.06	





















Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
239				1-solid support	white solid	379.13	1.62	400.98 M+Na	
240				1-solid support	white solid	395.91	1.13	396.01	
241				8	amber glass	592.29	1.69 min Method A	592.39	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.56 (d, 1H, J <sub>AB</sub> =16Hz), 4.40 (d, 1H, J <sub>AB</sub> =16Hz), 3.56 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.40 (t, 4H, J=6.0Hz), 1.95 (m, 1H), 1.68-1.52 (m, 2H), 1.42 (q, 4H, J=6.0Hz), 1.28-1.22 (m, 12H), 0.98 (d, 3H, J=7.0Hz), 0.88 (t, 6H, J=6.0Hz)
242				8	amber glass	648.4	1.88 min Method A	648.43	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.40 (s, br, 1H), 4.56 (d, 1H, J <sub>AB</sub> =16Hz), 4.40 (d, 1H, J <sub>AB</sub> =16Hz), 3.56 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.47 (t, 4H, J=6.0Hz), 1.95 (m, 1H), 1.68-1.50 (m, 2H), 1.43-1.14 (m, 14H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz), 0.88 (t, 6H, J=6.0Hz)







Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
243				9	clear glass	524.08	1.35 min Method A	542.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.34 (s, br, 1H), 4.70 (d, 1H, J <sub>AB</sub> =16Hz), 4.48 (d, 1H, J <sub>AB</sub> =16Hz), 4.24-4.08 (m, 4H), 3.42 (s, 2H), 3.25 (t, 1H, J=6Hz), 2.51 (s, 3H), 1.94 (M, 1H), 1.68-1.54 (m, 2H), 1.22 (t, 3H, J=6.0Hz), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
244				6	white solid	565.14	1.29 min Method B	564.99	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.12 (br s, 1H), 7.77 (d, 2H, J=8.8), 7.70 (d, 2H, J=8.6), 7.47 (d, 2H, J=8.4), 7.45 (d, 2H, J=8.0), 6.25 (br s, 1H), 5.31 (br s, 1H), 4.66 (d, 1H, J=15.7), 4.38 (d, 1H, J=15.8), 4.27 (t, 1H, J=7.1), 3.51-3.89 (m, 4H), 2.35-2.74 (m, 4H), 1.78-1.88 (m, 2H), 1.40-1.65 (m, 4H), 1.24-1.32 (m, 2H), 1.03-1.10 (m, 1H), 0.74 (d, 3H, J=6.5), 0.65 (d, 3H, J=6.6)
245				6	white solid	437.95	1.36 min Method B	438.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.72 (dd, 2H, J=8.3), 7.69 (dd, 2H, J=1.9, 8.7), 7.46 (dd, 2H, J=1.8, 8.7), 7.44 (d, 2H, J=8.6), 6.21 (br s, 1H), 5.98 (br s, 1H), 5.88 (br s, 1H), 5.39 (br s, 1H), 4.66 (d, 1H, J=15.7), 4.41 (d, 1H, J=15.9), 4.29 (t, 1H, J=6.5), 1.77-1.87 (m, 1H), 1.25-1.36 (m, 1H), 1.03-1.11 (m, 1H), 0.75 (d, 3H, J=6.6), 0.65 (d, 3H, J=6.6)







Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
246				6	white solid	549.14	1.34 min Method B	549.00	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.82 (d, 2H, J = 7.8), 7.67 (dd, 2H, J = 2.0, 8.7), 7.40-7.46 (m, 5H), 6.22 (br s, 1H), 5.23 (br s, 1H), 4.61 (d, 1H, J = 15.9), 4.42 (d, 1H, J = 15.7), 4.28 (t, 1H, J = 7.2), 3.60-3.69 (m, 2H), 2.45-2.83 (m, 6H), 1.40-1.85 (m, 7H), 1.24-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.75 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.6).
247				6	white solid	524.11	1.61 min Method B	523.94	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.68 (d, 2H, J = 8.4), 7.46 (d, 2H, J = 8.4), 7.39 (d, 2H, J = 8.1), 7.29 (d, 2H, J = 8.1), 6.20 (br s, 1H), 5.24 (br s, 1H), 4.60 (d, 1H, J = 15.8), 4.44 (d, 1H, J = 15.9), 4.30 (t, 1H, J = 6.9), 3.70-4.05 (br m, 4H), 2.45-2.60 (m, 4H), 1.73-1.80 (m, 1H), 1.28-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.76 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.6).
248				2	red solid	466.00	1.48 min Method B	466.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.65 (d, 2H, J = 7.0Hz), 7.41 (d, 2H, J = 7.0Hz), 7.20 (d, 2H, J = 8.8Hz), 6.79 (d, 2H, J = 8.8Hz), 6.23 (s, br, 1H), 5.20 (s, br, 1H), 4.32 (dd, 2H, J = 50Hz, 15Hz), 4.19-4.27 (m, 1H), 3.84-3.87 (m, 4H), 3.12-3.16 (m, 4H), 1.91-1.95 (m, 1H), 1.35-1.39 (m, 1H), 0.92-1.06 (m, 2H), 0.74 (t, 3H, J = 8.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
249				2	yellow solid	479.05	1.18min Method B	479.02	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.63 (d, 2H, J = 8.0Hz), 7.40 (d, 2H, J = 8.0Hz), 7.19 (d, 2H, J = 8.8Hz), 6.78 (d, 2H, J = 8.8Hz), 6.25 (s, br, 1H), 5.21 (s, br, 1H), 4.36 (dd, 2H, J = 50Hz, 15Hz), 4.20-4.27 (m, 1H), 3.28-3.35 (m, 4H), 2.69-2.76 (m, 4H), 2.48 (s, 3H), 1.93-1.97 (m, 1H), 1.35-1.39 (m, 1H), 0.90-1.07 (m, 2H), 0.72 (t, 3H, J = 8.0Hz).
250				2	tan solid	474.03	1.92min Method B	474.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J = 8.8Hz), 7.43-7.45 (m, 4H), 7.12 (d, 2H, J = 8.8Hz), 6.78 (d, 2H, J = 8.8Hz), 6.19 (s, br, 1H), 5.18 (s, br, 1H), 4.56 (dd, 2H, J = 50Hz, 15Hz), 4.21-4.30 (m, 1H), 2.01 (s, 6H), 1.93-1.97 (m, 1H), 1.35-1.39 (m, 1H), 0.90-1.07 (m, 2H), 0.72 (t, 3H, J = 8.0Hz).
251				6	white solid	482.00	2.01 min Method B	479.07	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67 (ddd, 2H, J = 1.9, 2.4, 8.7), 7.58 (d, 2H, J = 8.1), 7.43 (ddd, 2H, J = 1.5, 2.4, 8.7), 7.37 (d, 2H, J = 8.2), 6.30 (br s, 1H), 5.70 (br s, 1H), 4.62 (d, 1H, J = 15.9), 4.46 (d, 1H, J = 15.9), 4.32 (t, 1H, J = 7.3), 3.51 (s, 3H), 3.32 (s, 3H), 1.73-1.80 (m, 1H), 1.28-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.74 (d, 3H, J = 6.5), 0.61 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
252				1-Method A	white solid	466.98	1.92 min Method A	467.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.92 (d, 1H, J=8.0Hz), 7.79 (A of ABq, 2H, J=8.8Hz), 7.72 (d, 1H, J=7.7Hz), 7.50 (B of ABq, 2H, J=8.8Hz), 7.31 (t, 1H, J=7.7Hz), 6.29 (bs, 1H), 5.21 (bs, 1H), 5.02 (s, 2H), 4.35 (q, 2H, J=7.0Hz), 4.27 (dd, 1H, J=8.6, 5.5Hz), 1.88-1.78 (m, 1H), 1.39 (t, 3H, J=7.0Hz), 1.37-1.29 (m, 1H), 1.02-0.93 (m, 1H), 0.75 (d, 3H, J=6.6Hz), 0.66 (d, 3H, J=6.6Hz).
253				1-Method A	white solid	481.01	1.81 min Method A	481.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (A of ABq, 2H, J=8.8Hz), 7.44 (B of ABq, 2H, J=8.8Hz), 7.27-7.15 (m, 3H), 6.24 (bs, 1H), 5.26 (bs, 1H), 4.55 (A of ABq, 1H, J=15.4Hz), 4.39 (B of ABq, 1H, J=15.4Hz), 4.29 (t, 1H, J=7.0Hz), 4.15 (q, 2H, J=7.2Hz), 1.87-1.78 (m, 1H), 1.37-1.29 (m, 1H), 1.26 (t, 3H, J=7.2Hz), 1.24-1.13 (m, 1H), 0.76 (d, 3H, J=6.2Hz), 0.67 (d, 3H, J=6.6Hz).
254				10	white solid	438.98	1.20 Method B	439.05	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt δ 8.04 (s, 1H), 8.03 (d, 1H, J=9.80Hz), 7.76 (d, 2H, J=7.6 Hz), 7.54 (d, 2H, J=7.6 Hz), 6.83 (d, 1H, J=9.8 Hz), 6.62 (br s, 1H), 6.40 (br s, 1H), 4.64 (d, 1H, J=15.9 Hz), 4.29 (m, 1H), 4.18 (d, 1H, J=15.9 Hz), 3.30 (s, 6H), 1.84 (m, 1H), 1.29 (m, 1H), 0.93 (m, 1H), 0.77 (d, 3H, J=6.5Hz), 0.72 (d, 3H, J=6.5Hz).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
255				1-Method A	light orange residue	450.14	2.02 min Method E	450.98	<sup>1</sup> H NMR (DMSO) δ 7.78 (d, 2H, J=8.4Hz), 7.58 (d, 2H, J=8.8Hz), 7.47 (s, br, 1H), 7.29 (d, 2H, J=8.8Hz), 7.00 (s br, 1H), 6.87 (d, 2H, J=8.8Hz), 6.03 (m, 1H), 5.32 (dd, 2H, J=12Hz, 56Hz), 4.63 (m, 4H), 5.32 (dd, 2H, J=12Hz, 56Hz), 4.35 (m, 1H), 1.33 (m, 3H), 0.80 (d, 3H, J=6.0Hz), 0.50 (d, 3H, J=6.0Hz).
256				7	white solid	548.22	1.87 min Method A	549.00	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.71 (d, 2H, J=8.6), 7.71 (d, 2H, J=8.9), 7.15-7.35 (m, 5H), 6.64 (s, 1H), 5.86 (s, 1H), 4.15 (dd, 1H, J=5.2, 9.5), 3.88 (d, 1H, J=13), 3.76 (d, 1H, J=13), 3.46 (t, 2H, J=6.7), 3.21-3.29 (m, 1H), 2.97 (dd, 1H, J=4.6, 14), 2.65-2.85 (m, 4H), 1.75-1.95 (m, 3H), 1.00-1.30 (m, 5H), 0.75-0.80 (m, 1H), 0.72 (d, 3H, J=6.7), 0.67 (d, 3H, J=6.7).
257				7	white solid	520.19	1.74 min Method A	521.31	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J=8.8), 7.51 (d, 2H, J=8.8), 7.33 (d, 2H, J=7.6), 7.28 (d, 2H, J=7.6), 7.03 (t, 1H, J=7.3), 6.67 (s, 1H), 5.42 (s, 1H), 3.97-4.22 (m, 3H), 3.27-3.35 (m, 1H), 2.78-3.02 (m, 3H), 1.83-1.99 (m, 3H), 1.09-1.42 (m, 4H), 0.75-0.82 (m, 1H), 0.74 (d, 3H, J=6.4), 0.67 (d, 3H, J=6.7).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
258				7	white solid	526.24	1.81 min Method A	527.34	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 8.9), 7.51 (d, 2H, 8.9), 6.65 (s, 1H), 5.37 (s, 1H), 4.15 (dd, 1H, J = 5.1, 6.5), 3.92 (d, 1H, J = 12), 3.82 (d, 1H, J = 14), 3.57-3.67 (m, 1H), 3.26 (dd, 1H, J = 10, 14), 2.98 (dd, 1H, J = 4.5, 14), 2.74 (q, 2H, J = 12, 24), 1.80-1.97 (m, 5H), 1.64-1.72 (m, 3H), 1.00-1.43 (m, 10H), 0.75-0.82 (m, 1H), 0.73 (d, 3H, J = 6.4), 0.67 (d, 3H, 6.7).
259				7	white solid	548.22	1.78 min Method A	549.32	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.71 (d, 2H, J = 8.2), 7.50 (d, 2H, J = 8.5), 7.30 (d, 4H, J = 4.3), 7.20-7.25 (m, 1H), 6.65 (s, 1H), 5.74 (s, 1H), 4.99 (t, 1H, J = 7.02), 4.70-4.77 (m, 1H), 4.10-4.25 (m, 1H), 4.00 (d, 1H, J = 13), 3.90 (d, 1H, J = 13), 3.15-3.35 (m, 1H), 2.90-3.00 (m, 1H), 2.60-2.75 (m, 2H), 1.50-1.95 (m, 5H), 1.46 (d, 3H, J = 6.7), 1.00-1.30 (m, 2H), 0.75-0.83 (m, 1H), 0.73 (d, 3H, J = 6.4), 0.67 (d, 3H, J = 6.4).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
260				7	white solid	588.18	1.90 min Method A	589.25	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 8.9), 7.62 (d, 4H, J = 8.6), 7.45 (d, 2H, J = 8.5), 6.67 (s, 1H), 6.52 (s, 1H), 4.45 (s, 1H), 4.16 (dd, 1H, J = 5.2, 9.8), 4.12 (d, 1H, J = 12), 4.03 (d, 1H, J = 14), 3.80 (dd, 1H, J = 10, 14), 3.00 (dd, 1H, J = 4.5, 14), 2.84-2.92 (m, 2H), 1.85-2.00 (m, 3H), 1.69 (d, 1H, J = 12), 1.10-1.35 (m, 3H), 0.75-0.82 (m, 1H), 0.74 (d, 3H, J = 6.7), 0.68 (d, 3H, J = 6.7).
261				7	white solid	554.14	1.86 min Method A	555.24	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 8.6), 7.52 (d, 2H, J = 8.9), 7.45 (s, 1H), 7.18 (d, 2H, J = 6.7), 6.95-7.02 (m, 1H), 6.65 (s, 1H), 6.50 (s, 1H), 5.50 (s, 1H), 4.16 (dd, 1H, J = 5.2, 9.5), 4.08 (d, 1H, J = 15), 3.99 (d, 1H, J = 14), 3.30 (dd, 1H, J = 10, 15), 2.99 (dd, 1H, J = 4.5, 15), 2.80-2.92 (m, 2H), 1.80-2.00 (m, 3H), 1.67 (d, 1H, J = 13), 1.05-1.40 (m, 4H), 0.75-0.80 (m, 1H), 0.74 (d, 3H, J = 6.7), 0.68 (d, 3H, J = 6.7).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
262				7	white solid	592.21	1.86 min Method A	593.30	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.96 (d, 2H, J=8.9), 7.73 (d, 2H, J=8.9), 7.51 (d, J=8.6), 7.42 (d, 2H, J=8.6), 6.65 (s, 2H), 5.45 (s, 1H), 4.83 (q, 2H, J=7.0), 4.16 (dd, 1H, J=5.2, 9.8), 4.11 (d, 1H, J=13), 4.03 (d, 1H, J=13), 3.30 (dd, 1H, J=10, 14), 3.00 (dd, 1H, J=4.2, 14), 2.81-2.95 (m, 2H), 1.84-2.01 (m, 3H), 1.68 (d, 1H, J=13), 1.37 (t, 3H, J=7.3), 1.08-1.34 (m, 4H), 0.76-0.82 (m, 1H), 0.74 (d, 3H, J=6.7), 0.68 (d, 3H, J=6.7).
263				7	white solid	530.20	2.18 min Method C	531.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J=8.5), 7.51 (d, 2H, J=8.6), 6.67 (s, 1H), 5.41 (s, 1H), 4.97 (s, 1H), 4.20 (q, 2H, J=7.0), 4.15 (dd, 1H, J=5.1, 9.5), 3.90-4.04 (m, 4H), 3.26 (dd, 1H, J=10, 14), 2.99 (dd, 1H, J=4.5, 14), 2.70-2.82 (m, 2H), 1.80-1.95 (m, 3H), 1.28 (t, 3H, J=7.3), 1.05-1.25 (m, 3H), 0.75-0.80 (m, 1H), 0.72 (d, 3H, J=6.7), 0.67 (d, 3H, J=6.4).
264				1-Method A	white solid	462.96	1.47 Method B	462.98	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.81 (d, 2H, J=8.4Hz), 7.75 (d, 2H, J=8.0Hz), 7.55 (d, 2H, J=8.4Hz), 7.5 (d, 2H, J=8.0Hz), 6.86 (s, 1H), 6.44 (s, 1H), 4.96 (d, 1H, J=15.6Hz), 4.36 (dd, 1H, J=5.6Hz, 6.0Hz), 1.99 (m, 1H), 1.29 (m, 1H), 1.06 (m, 1H), 0.77 (d, 3H, J=6.8Hz), 0.74 (d, 3H, J=6.8Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
265				1-Method A	white solid	481.01	1.81 min Method A	481.3	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.61 (A of ABq, 2H, J=8.8Hz), 7.42 (B of ABq, 2H, J=8.8Hz), 7.27 (A of ABq, 2H, J=8.4Hz), 7.19 (B of ABq, 2H, J=8.4Hz), 6.21 (bs, 1H), 5.19 (bs, 1H), 4.52 (A of ABq, 1H, J=15.5Hz, 1H), 4.39 (B of ABq, 1H, J=15.5Hz, 1H), 4.30 (t, 1H, J=7.3Hz), 4.14 (q, 2H, J=7.1Hz), 3.58 (s, 2H), 1.86-1.76 (m, 1H), 1.36-1.27 (m, 1H), 1.14 (t, 3H, J=7.1Hz), 1.23-1.13 (m, 1H), 0.76 (d, 3H, J=6.2Hz), 0.66 (d, 3H, J=6.6Hz).
266				8	white foam	570.16	1.17 min Method A	570.39	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.21 (d, 2H, J=4.0Hz), 8.02 (d, 2H, J=8.0Hz), 7.59 (d, 2H, J=8.0Hz), 7.39-7.33 (m, 4H), 6.83 (d, 2H, J=4.0Hz), 6.23 (s, br, 1H), 5.34 (s, br, 1H), 4.69 (d, 1H, J <sub>ab</sub> =16Hz), 4.39 (d, 1H, J <sub>ab</sub> =16Hz), 3.57 (s, 2H), 3.27-3.15 (m, 5H), 2.58-2.45 (m, 4H), 1.94 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz).
267				8	white foam	508.08	1.28 min Method A	508.21	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.63 (d, 2H, J=8.0Hz), 7.43-7.36 (m, 4H), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.59 (d, 1H, J <sub>ab</sub> =16Hz), 4.36 (d, 1H, J <sub>ab</sub> =16Hz), 2.83 (m, 1H), 2.41 (m, 2H), 2.18 (m, 1H), 1.95 (m, 1H), 1.85 (m, 1H), 1.76-1.52 (m, 5H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
268				8	white foam	508.08	1.26 min Method A	508.18	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.65 (d, 2H, J=8.0Hz), 7.45 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.59 (d, 1H, J <sub>AB</sub> =16Hz), 4.36 (d, 1H, J <sub>AB</sub> =16Hz), 3.77 (m, 1H), 3.47 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 2.80 (m, 4H), 1.95 (m, 1H), 1.77-1.50 (m, 6H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
269				8	white foam	437.99	1.28 min Method A	438.16	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=8.0Hz), 7.66 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.87 (s, br, 1H), 4.76 (d, 1H, J <sub>AB</sub> =16Hz), 4.30 (d, 1H, J <sub>AB</sub> =16Hz), 3.54 (s, 5H), 3.25 (t, 1H, J=6.0Hz), 1.94 (m, 1H), 1.60 (m, 2H), 1.18 (s, br, NH), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
270				8	white solid	468.02	1.28 min Method A	468.16	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.0Hz), 7.67 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.33 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.85 (s, br, 1H), 4.24-4.12 (m, 4H), 3.67 (s, 2H), 3.24 (t, 1H, J=6.0Hz), 3.06 (t, 2H, J=6.0Hz), 2.60 (s, br, NH), 1.95 (m, 1H), 1.68-1.52 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
271				8	clear glass	482.05	1.31 min Method A	482.18	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.79 (d, 2H, J=8.0Hz), 7.69 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.33 (d, 2H, J=8.0Hz), 6.35 (s, br, 1H), 5.85 (s, br, 1H), 4.60 (d, 1H, J <sub>AB</sub> =16Hz), 4.43 (d, 1H, J <sub>AB</sub> =16Hz), 3.67 (s, 2H), 3.61 (t, 2H, J=6.0Hz), 3.35 (s, 3H), 3.29-3.24 (m, 3H), 1.94 (m, 1H), 1.62 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
272				9	white solid	523.1	1.30 min Method A	523.40	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.02 (d, 2H, J=8.0Hz), 7.71 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 7.28 (d, 2H, J=8.0Hz), 6.23 (s, br, 1H), 5.51 (s, br, 1H), 4.46 (s, 2H), 4.70 (d, 1H, J <sub>AB</sub> =16Hz), 4.33 (d, 1H, J <sub>AB</sub> =16Hz), 3.25 (t, 1H, J=6.0Hz), 2.69 (s, 3H), 2.63 (s, 2H), 2.20 (s, 6H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
273				16	white solid	436.96	1.43 min Method B	437.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.87 (d, 2H, J=8.4), 7.67 (dd, 2H, J=1.8, 8.7), 7.42-7.46 (m, 4H), 6.21 (br s, 1H), 5.28 (br s, 1H), 4.64 (d, 1H, J=15.9), 4.45 (d, 1H, J=15.9), 4.31 (t, 1H, J=6.6), 2.58 (s, 3H), 1.73-1.80 (m, 1H), 1.25-1.35 (m, 1H), 1.05-1.14 (m, 1H), 0.74 (d, 3H, J=6.5), 0.65 (d, 3H, J=6.6)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
274				17	yellow foam	549.14	1.38 min Method A	549.16	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.63 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.23 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.35 (s, br, 1H), 4.75 (d, 1H, J <sub>ab</sub> =16Hz), 4.38 (d, 1H, J <sub>ab</sub> =16Hz), 3.25 (t, 1H, J=6.0Hz), 2.65 (t, 2H, J=6.0Hz), 2.56-2.44 (m, 6H), 1.95 (m, 1H), 1.68-1.45 (m, 8H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
275				17	clear glass	564.15	1.21 min Method A	564.32	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 9.30 (s, 1H, NH), 8.02 (d, 2H, J=8.0Hz), 7.61 (d, 2H, J=8.0Hz), 7.36 (d, 2H, J=8.0Hz), 7.23 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.33 (s, br, 1H), 4.72 (d, 1H, J <sub>ab</sub> =16Hz), 4.48 (d, 1H, J <sub>ab</sub> =16Hz), 3.25 (t, 1H, J=6.0Hz), 2.65-2.38 (m, 12H), 2.28 (s, 2H), 1.95 (m, 1H), 1.59 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
276				17	tan foam	509.07	1.33 min Method A	509.17	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 9.31 (s, 1H, NH), 8.02 (d, 2H, J=8.0Hz), 7.63 (d, 2H, J=8.0Hz), 7.36 (d, 2H, J=8.0Hz), 7.23 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.34 (s, br, 1H), 4.71 (d, 1H, J <sub>ab</sub> =16Hz), 4.48 (d, 1H, J <sub>ab</sub> =16Hz), 3.25 (t, 1H, J=6.0Hz), 2.66 (t, 2H, J=8.0Hz), 2.56 (t, 2H, J=8.0Hz), 2.38 (s, 6H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
277				9	tan foam	515.04	1.47 min Method A	515.13	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.95 (s, br, NH), 8.67 (d, 1H, J=8.0Hz), 8.17 (d, 1H, J=8.0Hz), 8.02 (d, 2H, J=8.0Hz), 7.77 (d, 2H, J=8.0Hz), 7.57 (m, 4H), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.72 (d, 1H, J <sub>ab</sub> =16Hz), 4.30 (d, 1H, J <sub>ab</sub> =16Hz), 3.25 (t, 1H, J=6.0Hz), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
278				7	white solid	507.06	2.44 min Method C	509.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 8.66 (br s, 2H), 7.80 (d, 1H, J=8.6), 7.73 (d, 2H, J=8.5), 7.51 (d, 2H, J=7.6), 7.41 (br s, 1H), 6.64 (br s, 1H), 5.35 (br s, 1H), 4.70 (br s, 1H), 4.10 (br s, 1H), 3.71 (br s, 1H), 3.33 (br s, 1H), 3.02 (dd, 2H, J=4.8, 16), 2.70-2.85 (br s, 1H), 1.50-2.09 (m, 5H), 1.18-1.33 (m, 4H), 0.73 (d, 3H, J=6.7), 0.68 (d, 3H, J=6.5).
279				7	white solid	549.14	2.76 min Method C	549.07	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.74 (dd, 2H, J=1.7, 6.7), 7.51 (dd, 2H, J=2.2, 6.9), 7.34 (d, 2H, J=8.0), 6.70 (br s, 1H), 6.60 (br s, 1H), 5.30 (br s, 1H), 4.13 (dd, 1H, J=5.5, 10), 3.35 (dd, 1H, J=11, 14), 3.00 (s, 6H), 2.85 (s, 2H), 1.80-2.00 (m, 3H), 1.50-1.70 (m, 1H), 1.10-1.30 (m, 4H), 0.80-0.90 (m, 1H), 0.745 (d, 3H, J=6.7), 0.67 (d, 3H, J=6.5).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
280				7	white solid	574.07	3.03 Method C	574.03	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 8.2), 7.65 (d, 2H, J = 7.9), 7.51 (d, 2H, J = 8.9), 7.48 (d, 2H, J = 7.6), 6.65 (br s, 1H), 5.45 (br s, 1H), 4.71 (br s, 1H), 4.13 (br s, 1H), 3.65 (br s, 1H), 3.30 (br s, 1H), 2.97 (d, 2H, J = 12), 2.65-2.86 (m, 1H), 1.45-2.07 (m, 6H), 0.98-1.85 (m, 3H), 0.73 (br, s, 3H), 0.67 (br, s, 3H).
281				7	white solid	470.04	2.78 min Method C	470.03	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.5), 7.50 (d, 2H, J = 8.6), 6.78-6.90 (m, 1H), 6.55-6.65 (m, 1H), 6.23 (dd, 1H, J = 1.5, 15), 5.33-5.60 (m, 1H), 4.50-4.75 (m, 1H), 4.09-4.20 (m, 1H), 3.90-4.05 (m, 1H), 2.80-3.25 (m, 3H), 2.40-2.75 (m, 1H), 1.50-2.00 (m, 8H), 1.00-1.40 (m, 3H), 0.73 (br, s, 3H), 0.67 (br, s, 3H).
282				7	white solid	506.07	2.86 min Method C	508.03	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 8.5), 7.51 (d, 2H, J = 8.5), 7.38 (br s, 4H), 6.65 (br s, 1H), 5.35 (br s, 1H), 4.71 (br s, 1H), 4.14 (br s, 1H), 3.76 (br s, 1H), 3.30 (br s, 1H), 2.60-3.05 (m, 3H), 0.99-2.05 (m, 10H), 0.73 (d, 3H, J = 7.8), 0.67 (d, 3H, J = 7.8).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
283				7	white solid	517.09	1.34 min Method A	517.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.5), 7.50-7.65 (m, 2H), 7.50 (d, 2H, J = 7.0), 7.35-7.45 (m, 2H), 6.67 (s, 1H), 5.32 (s, 1H), 4.14 (dd, 1H, J = 5.0, 9.0), 3.52 (br s, 1H), 3.28 (t, 1H, J = 14), 2.97 (dd, 1H, J = 3.5, 14), 2.82 (br s, 1H), 1.00-2.00 (m, 10H), 0.71 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.5).
284				7	white solid	550.12	2.87 min Method C	550.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.6), 7.50 (d, 2H, J = 8.5), 7.13 (d, 2H, J = 8.5), 6.85 (br s, 2H), 6.63 (d, 1H, J = 33), 5.41 (br s, 1H), 4.62 (t, 1H, J = 14), 4.10-4.17 (m, 1H), 3.75-3.90 (m, 4H), 3.65 (s, 3H), 3.14-3.30 (m, 1H), 2.80-2.95 (m, 2H), 2.43-2.60 (m, 1H), 1.45-2.00 (m, 4H), 1.15-1.30 (m, 2H), 0.71 (dd, 3H, J = 7.6, 8.4), 0.65 (dd, 3H, J = 6.0, 8.0).
285				7	white solid	541.50	2.76 min Method C	540.98	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 8.43 (s, 1H), 7.73 (d, 2H, J = 8.5), 7.70 (d, 1H, J = 2.4, 8.4), 7.51 (d, 2H, J = 8.6), 7.87 (d, 1H, J = 8.2), 6.63 (br s, 1H), 5.35 (br s, 1H), 4.68 (br s, 1H), 4.15 (dd, 1H, J = 4.9, 9.8), 3.71 (br s, 1H), 3.31 (br s, 1H), 3.00 (dd, 2H, J = 4.8, 14), 2.65-2.86 (m, 1H), 1.77-2.07 (m, 3H), 1.6-1.76 (m, 1H), 1.00-1.86 (m, 3H), 0.85-0.93 (m, 1H), 0.73 (d, 3H, J = 6.7), 0.67 (d, 3H, J = 6.7).





















Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
286				7	white solid	545.11	1.36 min Method A	545.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 8.11 (d, 2H, J = 8.6), 7.75 (d, 2H, J = 8.6), 7.73 (d, 2H, J = 8.9), 7.50 (d, 2H, J = 8.9), 6.65 (br s, 1H), 5.38 (br s, 1H), 4.14 (dd, 1H, J = 5.5, 9.5), 3.80 (br s, 1H), 3.27 (dd, 1H, J = 10, 14), 2.97 (dd, 2H, J = 4.6, 14), 1.17-2.00 (m, 11H), 0.75-0.81 (m, 1H), 0.73 (d, 3H, J = 6.4), 0.67 (d, 3H, J = 6.7).
287				12	white solid	453.01	1.81 min Method A	453.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.61 (d, 2H, J = 8.7), 7.40 (d, 2H, J = 8.7), 7.37 (d, 2H, J = 8.4), 7.26 (d, 2H, J = 8.4), 6.28 (br s, 1H), 5.25 (br s, 1H), 4.49 (d, 1H, J = 15.9), 4.41 (d, 1H, J = 15.9), 4.33 (t, 1H, J = 6.6), 1.73-1.80 (m, 1H), 1.55 (s, 6H), 1.28-1.35 (m, 1H), 1.20-1.25 (m, 1H), 0.77 (d, 3H, J = 6.5), 0.66 (d, 3H, J = 6.6).
288				7	white solid	493.07	1.25 min Method A	493.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 8.51 (s, 2H), 7.73 (d, 2H, J = 8.5), 7.50 (d, 2H, J = 8.5), 6.67 (s, 1H), 5.83 (s, 1H), 4.15 (dd, 1H, J = 5.2, 8.9), 3.50 (br s, 2H), 3.25 (dd, 1H, J = 8.5, 14), 2.75-3.05 (m, 3H), 1.60-2.10 (m, 6H), 1.10-1.40 (m, 4H), 0.75-0.85 (m, 1H), 0.72 (d, 3H, J = 6.4), 0.67 (d, 3H, J = 6.7).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
289				2	yellow solid	493.07	0.93 min Method B	493.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.67 (d, 2H, J=7.0Hz), 7.44 (d, 2H, J=7.0Hz), 7.19 (d, 2H, J=8.0Hz), 6.46 (d, 2H, J=8.0Hz), 6.21 (s, br, 1H), 5.17 (s, br, 1H), 4.31 (dd, 2H, J=50Hz, 15Hz), 4.15-4.22 (m, 1H), 3.84-3.87 (m, 4H), 3.91-3.99 (m, 1H), 3.51-3.54 (m, 3H), 3.22-3.26 (m, 1H), 2.75 (s, 3H), 2.72 (s, 3H), 2.23-2.36 (m, 2H), 1.91-1.98 (m, 1H), 1.32-1.40 (m, 1H), 0.81-1.04 (m, 2H), 0.73 (t, 3H, 7.2Hz).
290				2	tan solid	464.03	1.16 min Method B	464.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.62 (d, 2H, J=8.8Hz), 7.40 (d, 2H, J=8.0Hz), 7.11-7.20 (m, 2H), 6.79-6.88 (m, 2H), 6.20 (s, br, 1H), 5.13 (s, br, 1H), 4.30 (dd, 2H, J=50Hz, 15Hz), 4.13-4.21 (m, 1H), 3.10-3.19 (m, 4H), 1.92-1.95 (m, 1H), 1.39-1.90 (m, 8H), 1.22-1.26 (m, 1H), 0.97-1.05 (m, 2H), 0.73 (t, 3H, J=8.0Hz).
291				11	orange solid	522.11	1.52 min Method E	522.1	<sup>1</sup> H NMR (DMSO) δ 7.78 (d, 2H, J=8.4Hz), 7.58 (d, 2H, J=8.8Hz), 7.47 (s, br, 1H), 7.27 (d, 2H, J=8.4Hz), 7.00 (s, br, 1H), 6.85 (d, 2H, J=8.8Hz), 4.63 (dd, 2H, J=16Hz, 38Hz), 4.34 (m, 1H), 4.03 (s, 2H), 2.63 (m, 2H), 2.42 (m, 3H), 1.39 (m, 10H), 0.80 (d, 3H, J=6.0Hz), 0.50 (d, 3H, J=6.0Hz).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
292				11	white solid	481.18	1.46 min Method E	482.06	<sup>1</sup> H NMR (DMSO) δ 7.79 (d, 2H, J = 8.8Hz), 7.60 (d, 2H, J = 8.8Hz), 7.49 (s, br, 1H), 7.35 (d, 2H, J = 8.4Hz), 7.01 (s br, 1H), 6.94 (d, 2H, J = 8.8Hz), 4.68 (dd, 2H, J = 16Hz, 5.3Hz), 4.35 (m, 1H), 4.28 (t, 2H, J = 4.8Hz), 3.34 (m, 2H), 2.86 (s, 6H), 1.33 (m, 3H), 0.80 (d, 3H, J = 6.0Hz), 0.50 (d, 3H, J = 6.0Hz)
293				11	white residue	509.21	1.52 min Method E	M+Na 532.03	<sup>1</sup> H NMR (DMSO) δ 7.79 (d, 2H, J = 8.4Hz), 7.60 (d, 2H, J = 8.4Hz), 7.48 (s, br, 1H), 7.35 (d, 2H, J = 8.8Hz), 7.01 (s br, 1H), 6.94 (d, 2H, J = 8.8Hz), 4.68 (dd, 2H, J = 17Hz, 5.4Hz), 4.35 (m, 1H), 4.28 (t, 2H, J = 4.8Hz), 3.51 (m, 3H), 3.21 (m, 3H), 1.29 (m, 9H), 0.80 (d, 3H, J = 6.0Hz), 0.50 (d, 3H, J = 6.0Hz)
294				11	light brown solid	544.12	1.78 min Method E	544.13	<sup>1</sup> H NMR (DMSO) δ 7.78 (d, 2H, J = 8.9Hz), 7.57 (d, 2H, J = 8.0Hz), 7.45 (s, br, 1H), 7.27 (d, 2H, J = 8.0Hz), 7.17 (t, 2H, J = 8.9Hz), 7.00 (s br, 1H), 6.84 (d, 2H, J = 8.0Hz), 6.75 (d, 2H, J = 8.0Hz), 6.62 (t, 1H, J = 8.0Hz), 4.65 (dd, 2H, J = 17Hz, 4.1Hz), 4.34 (m, 1H), 4.10 (t, 2H, J = 5.5Hz), 3.71 (t, 2H, J = 7.9Hz), 2.96 (m, 3H), 1.32 (m, 3H), 0.79 (d, 3H, J = 6.0Hz), 0.49 (d, 3H, J = 6.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
295				11	white powder	558.14	1.62 min Method E	558.10	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J = 8.6Hz), 7.60 (d, 2H, J = 8.6Hz), 7.54 (m, 2H), 7.49 (m, 4H), 7.34 (d, 2H, J = 8.6Hz), 7.01 (s br, 1H), 6.93 (d, 2H, J = 8.6Hz), 4.68 (dd, 2H, J = 17.0Hz, 5.4Hz), 4.47 (m, 1H), 4.36 (m, 4H), 3.50 (m, 2H), 2.80 (s, 3H), 1.34 (m, 3H), 0.80 (d, 3H, J = 6.0Hz), 0.51 (d, 3H, J = 6.0Hz)
296				11	white powder	508.08	1.49 min Method E	508.07	<sup>1</sup> H NMR (DMSO) δ 7.79 (d, 2H, J = 8.6Hz), 7.61 (d, 2H, J = 8.6Hz), 7.49 (s, br, 1H), 7.35 (d, 2H, J = 8.6Hz), 7.00 (s br, 1H), 6.94 (d, 2H, J = 8.6Hz), 4.68 (dd, 2H, J = 17.0Hz, 5.3Hz), 4.35 (m, 1H), 4.27 (m, 2H), 3.59 (m, 4H), 3.13 (m, 2H), 2.03 (m, 2H), 1.89 (m, 2H), 1.34 (m, 3H), 0.80 (d, 3H, J = 6.0Hz), 0.52 (d, 3H, J = 6.0Hz)
297				11	off white solid	524.08	1.46 min Method E	524.09	<sup>1</sup> H NMR (DMSO) δ 7.78 (d, 2H, J = 8.9Hz), 7.57 (d, 2H, J = 8.0Hz), 7.45 (s, br, 1H), 7.27 (d, 2H, J = 8.0Hz), 7.17 (t, 2H, J = 8.9Hz), 7.00 (s br, 1H), 6.84 (d, 2H, J = 8.0Hz), 6.75 (d, 2H, J = 8.0Hz), 6.62 (t, 1H, J = 8.0Hz), 4.65 (dd, 2H, J = 17Hz, 4.1Hz), 4.34 (m, 1H), 4.10 (t, 2H, J = 5.5Hz), 3.71 (t, 2H, J = 7.9Hz), 2.96 (m, 3H), 1.32 (m, 3H), 0.79 (d, 3H, J = 6.0Hz), 0.49 (d, 3H, J = 6.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
298				11	white solid	540.15	1.52 min Method E	540.06	<sup>1</sup> H NMR (DMSO) δ 7.80(d, 2H, J=8.8Hz), 7.60 (d, 2H, J=8.0Hz), 7.48 (s br, 1H), 7.35 (d, 2H, J=8.8Hz), 7.01 (s br, 1H) 6.95 (d, 2H, J=8.8Hz), 4.68(dd, 2H, J=16.4Hz, 5.5Hz), 4.34 (m, 3H), 3.79 (m, 2H), 3.58 (m, 2H), 3.27 (m, 2H), 3.02 (m, 2H), 2.89 (m, 2H), 1.34 (m, 3H), 0.80 (d, 3H, J=6.0Hz), 0.52 (d, 3H, J=6.0Hz)
299				11	light orange solid	537.13	1.43 min Method E	537.13	<sup>1</sup> H NMR (DMSO) δ 7.79(d, 2H, J=8.8Hz), 7.59 (d, 2H, J=8.0Hz), 7.48 (s br, 1H), 7.31 (d, 2H, J=8.8Hz), 7.00 (s br, 1H) 6.88 (d, 2H, J=8.8Hz), 4.66 (dd, 2H, J=16.4Hz, 4.9Hz), 4.35 (m, 1H), 4.11 (m, 2H), 3.18 (m, 9H), 2.78 (m, 3H), 2.63 (m, 1H), 1.34 (m, 3H), 0.80 (d, 3H, J=6.0Hz), 0.51 (d, 3H, J=6.0Hz)
300				2	tan solid	507.10	1.19min Method B	507.2	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.63 (d, 2H, J=8.0Hz), 7.41 (d, 2H, J=8.0Hz), 7.13-7.24 (m, 2H), 6.74-6.80 (m, 2H), 6.23 (s, br, 1H), 5.13 (s, br, 1H), 4.37 (dd, 2H, J=50Hz, 15Hz), 4.11-4.19 (m, 1H), 3.77-3.81 (m, 1H), 3.44 (s, 6H), 3.06-3.13 (m, 8H), 1.93-1.96 (m, 1H), 1.25-1.29 (m, 1H), 0.95-1.09 (m, 2H), 0.71 (t, 3H, J=8.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
301				10	yellow foam	465.02	1.22 Method B	465.25	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt: δ 8.07 (s, 1H), 7.91 (d, 1H, J=9.6Hz), 7.74 (d, 2H, J=6.8Hz), 7.52(d, 2H, J=6.8Hz), 6.64 (d, 1H, J=9.6Hz), 6.36 (s, 1H), 5.77 (s, 1H), 4.52 (d, 1H, J=16.0Hz), 4.28 (dd, 1H, J=5.6Hz, 6.0Hz), 4.21 (d, 1H, J=16.0Hz), 3.62 (m, 4H), 2.13 (m, 4H), 1.84 (m, 1H), 1.32 (m, 1H), 0.96 (m, 1H), 0.79 (d, 3H, J=6.8Hz), 0.72 (d, 3H, J=6.8Hz).
302				10	yellow foam	479.05	1.28 Method B	479.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt: δ 8.01 (s, 1H), 7.95 (d, 1H, J=9.6Hz), 7.75 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.4Hz), 6.88 (d, 1H, J=9.6Hz), 6.43 (s, 1H), 6.04 (s, 1H), 4.53 (d, 1H, J=16.0Hz), 4.28 (dd, 1H, J=5.6Hz, 6.0Hz), 4.20 (d, 1H, J=16.0Hz), 3.65 (m, 4H), 1.82 (m, 1H), 1.74 (m, 6H), 1.31 (m, 1H), 0.95 (m, 1H), 0.78 (d, 3H, J=6.4Hz), 0.71 (d, 3H, J=6.4Hz).
303				10	white solid	481.02	1.16 Method B	481.05	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt: δ 8.25 (s, 1H), 8.06 (d, 1H, J=9.6Hz), 7.74(d, 2H, J=8.0Hz), 7.54 (d, 2H, J=8.0Hz), 6.91(d, 1H, J=9.6Hz), 6.55 (s, 1H), 6.28 (s, 1H), 4.61 (d, 1H, J=16.0Hz), 4.28 (dd, 1H, J=5.2Hz, 6.2Hz), 4.21 (d, 1H, J=16.0Hz), 3.87 (m, 4H), 3.67 (m, 4H), 1.84 (m, 1H), 1.27 (m, 1H), 0.93 (m, 1H), 0.76 (d, 3H, J=6.4Hz), 0.72 (d, 3H, J=6.4Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
304				10	brown solid	493.07	1.357 Method B	493.04	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt: δ 8.04 (s, 1H), 7.93 (d, 1H, J=9.2Hz), 7.73 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 6.69 (d, 1H, J=9.2Hz), 6.38 (s, 1H), 5.98 (s, 1H), 4.50 (d, 1H, J=16.0Hz), 4.30 (dd, 1H, J=5.6Hz, 6.0Hz), 4.22 (d, 1H, J=16.0Hz), 4.20 (m, 2H), 3.87 (m, 4H), 2.24 (m, 2H), 1.90 (m, 2H), 1.84 (m, 1H), 1.36 (d, 3H, J=2.0Hz), 1.34 (d, 3H, J=2.0Hz), 1.32 (m, 1H), 0.98 (m, 1H), 0.79 (d, 3H, J=6.4Hz), 0.72 (d, 3H, J=6.4Hz).
305				7	white solid	513	1.03 min Method A	513.36	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J=8.5), 7.50 (d, 2H, J=7.0), 6.65 (s, 1H), 6.35 (s, 1H), 4.14 (dd, 1H, J=5.5, 9.0), 3.25 (dd, 1H, J=10, 14), 1.35-2.95 (m, 24H), 1.15-1.30 (m, 3H), 0.72 (d, 3H, J=6.5), 0.67 (d, 3H, J=6.7).
306				7	white solid	487	1.43 min Method A	487.019	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.72 (d, 2H, J=8.4), 7.51 (d, 2H, J=8.7), 6.67 (d, 1H, J=24.3), 5.40 (d, 1H, J=12, 6), 4.54 (br s, 1H), 3.90-4.20 (m, 4H), 3.40-3.55 (m, 2H), 3.05-3.35 (m, 2H), 2.85-3.05 (m, 2H), 2.40-2.60 (m, 1H), 2.35 (d, 6H, J=8.1), 0.60-1.95 (m, 11H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
307				1-Method A	clear oil	444.03	1.28 min Method B	444.04	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.72 (d, J=6.8Hz, 2H), 7.52 (d, J=6.8Hz, 2H), 6.75 (s, br, 1H), 5.79 (s, br, 1H), 4.14 (dd, J=9.6Hz, 4.8Hz, 1H), 3.23 (dd, J=14.4Hz, 10.0Hz, 1H), 3.12 (m, 1H), 2.92 (dd, J=14.4Hz, 4.8Hz, 1H), 2.77 (m, 6H), 2.09 (m, 2H), 1.83 (m, 1H), 1.71 (m, 1H), 0.75-1.52 (m, 8H), 0.73 (d, J=6.8Hz, 3H), 0.67 (d, J=6.8Hz, 3H).
308				23	tan foam	495.04	1.31 min Method A	495.14	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.85 (s, 1H, NH), 8.02 (d, 2H, J=8.0Hz), 7.75 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.29 (d, 2H, J=8.0Hz), 6.23 (s, br, 1H), 5.39 (s, br, 1H), 4.62 (m, 4H), 3.25 (t, 1H, J=6.0Hz), 2.95 (s, 6H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz).
309				23	tan foam	535.11	1.34 min Method A	535.29	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.85 (s, 1H, NH), 8.02 (d, 2H, J=8.0Hz), 7.78 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.30 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.36 (s, br, 1H), 4.62 (m, 4H), 3.25 (t, 1H, J=6.0Hz), 2.42 (m, 4H), 1.95 (m, 1H), 1.68-1.38 (m, 8H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
310				23	tan foam	550.12	1.24 min Method A	550.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 8.83 (s, 1H, NH), 8.02 (d, 2H, J=8.0Hz), 7.78 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 7.30 (d, 2H, J=8.0Hz), 6.20 (s, br, 1H), 5.39 (s, br, 1H), 4.71 (d, 1H, J <sub>AB</sub> =16Hz), 4.48 (d, 1H, J <sub>AB</sub> =16Hz), 4.26 (s, 2H), 3.25 (m, 1H), 2.67 (m, 8H), 2.40 (s, 3H), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)
311				7	white solid	492.09	2.30 min Method C	492.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.73 (d, 2H, J=8.4), 7.48 (d, 2H, J=8.4), 7.15-7.38 (m, 5H), 6.67 (s, 1H), 5.37 (s, 1H), 4.14 (dd, 1H, J=5.4, 9.0), 3.47 (s, 2H), 3.24 (dd, 1H, J=10, 14), 2.75-3.05 (m, 3H), 1.45-2.05 (m, 10H), 0.75-0.90 (m, 1H), 0.71 (d, 3H, J=6.6), 0.65 (d, 3H, J=6.6)
312				7	white solid	529.15	2.27 min Method C	529.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.73 (d, 2H, J=8.7), 7.50 (d, 2H, J=8.7), 6.65 (d, 1H, J=5.1), 4.55 (br s, 1H), 4.05-4.30 (m, 1), 3.57-3.95 (m, 1H), 3.10-3.40 (m, 1H), 2.40-3.00 (m, 12H), 1.70-2.00 (m, 3H), 0.90-1.30 (m, 10H), 0.60-0.75 (m, 6H)
313				7	white solid	573.16	1.78 min Method A	573.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.72 (d, 2H, J=7.8), 7.50 (d, 2H, J=8.7), 6.65 (d, 1H, J=14), 5.35 (s, 1H), 4.45-4.65 (m, 1H), 3.67-4.25 (m, 3H), 3.35-3.60 (m, 3H), 3.10-3.25 (m, 1H), 2.80-3.10 (m, 2H), 2.90 (s, 3H), 0.95-2.00 (m, 7H), 1.46 (s, 9H), 0.73 (br, s, 3H), 0.67 (br, s, 3H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
314				7	white solid	532.11	1.47 min Method A	532.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.74 (d, 2H, J=8.8), 7.50 (d, 3H, 8.8), 7.17-7.27 (m, 3H), 6.63 (s, 1H), 5.60 (s, 1H), 4.13 (dd, 1H, J=5.5, 9.2), 3.80 (s, 2H), 3.28 (dd, 1H, J=9.5, 15), 2.80-3.05 (m, 3H), 1.50-2.25 (m, 6H), 1.15-1.45 (m, 5H), 0.72 (d, 3H, J=6.6), 0.66 (d, 3H, J=6.6)
315				7	white solid	444.04	2.11 min Method C	444.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.73 (d, 2H, J=8.7), 7.48 (d, 2H, J=8.4), 6.69 (br s, 1H), 5.45 (br s, 1H), 4.14 (dd, 1H, J=10, 15), 3.15-3.35 (m, 1H), 2.90-3.05 (m, 1H), 2.75-2.90 (m, 1H), 2.60-2.75 (m, 1H), 1.50-2.25 (m, 6H), 1.05-1.40 (m, 3H), 1.00 (d, 6H, J=11), 0.75-0.90 (m, 1H), 0.71 (d, 3H, J=10), 0.66 (d, 3H, J=10)
316				9	yellow foam	515.04	1.50 min Method A	515.08	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.7 (s, 1H, NH), 8.82 (d, 2H, J=4.0Hz), 8.02 (d, 2H, J=8.0Hz), 7.76 (d, 2H, J=8.0Hz), 7.73 (d, 2H, J=4.0Hz), 7.46 (d, 2H, J=8.0Hz), 7.38 (d, 2H, J=8.0Hz), 6.27 (s, br, 1H), 5.39 (s, br, 1H), 4.72 (d, 1H, J <sub>AB</sub> =16Hz), 4.30 (d, 1H, J <sub>AB</sub> =16Hz), 4.26 (s, 2H), 3.25 (t, 1H, J=6.0Hz), 1.95 (m, 1H), 1.60 (m, 2H), 0.98 (d, 3H, J=7.0Hz), 0.94 (d, 3H, J=7.0Hz)



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
323				9	tan solid	465.14	1.52min Method B	466.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J=8.4Hz), 7.46 (d, 2H, J=8.4Hz), 7.43 (d, 2H, J=8.0Hz), 7.10 (d, 2H, J=8.0Hz), 6.19 (s, br, 1H), 5.16 (s, br, 1H), 4.44 (dd, 2H, J=50Hz, 15Hz), 4.31-4.35 (m, 1H), 3.23 (s, 3H), 1.85 (s, 3H), 1.76-1.82 (m, 1H), 1.14-1.35 (m, 2H), 0.78 (d, 3H, J=7.0Hz), 0.65 (d, 3H, J=7.0Hz).
324				7	white powder	473.04	1.54 mins Method A	473.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.73 (d, 2H, J=8.1), 7.50 (d, 2H, J=8.1), 6.67 (s, 1H), 5.85 (s, 1H), 4.15 (dd, 1H, J=5.1, 9.2), 3.65 (d, 2H, J=10), 3.26 (dd, 1H, J=9.9, 15), 2.97 (dd, 1H, J=4.4, 14), 2.60-2.85 (m, 3H), 2.80 (s, 6H), 1.75-1.95 (m, 3H), 1.05-1.30 (m, 3H), 0.72 (d, 3H, J=6.2), 0.65 (d, 3H, J=6.6).
325				7	white solid	507.06	1.41 min Method A	507.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.64 (s, 2H), 7.72 (d, 3H, J=8.4), 7.50 (d, 2H, J=8.8), 7.35 (dd, 1H, J=4.8, 7.7), 6.63 (s, 1H), 5.39 (br s, 1H), 4.69 (br s, 1H), 4.05-4.20 (m, 1H), 3.72 (br s, 1H), 3.25 (m, 1H), 2.50-3.200 (m, 3H), 1.50-2.10 (m, 5H), 1.00-1.40 (m, 3H), 0.72 (d, 3H, J=6.6), 0.66 (d, 3H, J=6.6).
326				1-Method A	white solid	423.90	1.55min Method B	424.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69-7.71 (m, 3H), 7.48-7.56 (m, 4H), 6.11 (s, br, 1H), 5.22 (s, br, 1H), 4.57 (dd, 2H, J=50Hz, 15Hz), 4.22-4.26 (m, 1H), 1.80-1.83 (m, 1H), 0.99-1.23 (m, 3H), 0.74 (t, 3H, J=8.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
327				1-Method A	white solid	459.06	1.57 min Method G	(M+Na) <sup>+</sup> 481.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.84 (d, 2H, J=8.8), 7.80 (d, 2H, J=8.6), 7.65 (d, 2H, J=8.7), 7.59 (d, 2H, J=8.3), 7.47 (s, 1H), 7.25 (s, 1H), 4.79 (ABq, 2H, Δν=5.1, J <sub>AB</sub> =17.4), 4.43 (dd, 1H, J=8.5, 6.6), 2.05 (m, 2H), 1.82 (m, 1H), 1.49 (m, 1H).
328				1-Method A	white solid	502.06	1.83 min Method G	(M+H) <sup>+</sup> 502.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.84 (d, 2H, J=8.8), 7.68 (d, 2H, J=8.0), 7.65-7.60 (m, 4H), 7.47 (s, 1H), 7.26 (s, 1H), 4.79 (s, 2H), 4.45 (dd, 1H, J=8.8, 6.1), 2.03 (m, 2H), 1.82 (m, 1H), 1.52 (m, 1H).
329				1-Method A	white solid	492.07	2.39 min Method G	(M+H) <sup>+</sup> 492.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.91 (d, 2H, J=8.3), 7.85 (d, 2H, J=8.8), 7.64 (d, 2H, J=8.6), 7.54 (d, 2H, J=8.3), 7.43 (s, 1H), 7.23 (s, 1H), 4.79 (ABq, 2H, Δν=3.4, J <sub>AB</sub> =17.2), 4.42 (dd, 1H, J=8.5, 6.1), 3.85 (s, 3H), 2.02 (m, 2H), 1.80 (m, 1H), 1.52 (m, 1H).
330				1-Method A	white solid	473.08	1.60 Method G	(M+Na) <sup>+</sup> 495.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.83 (d, 2H, J=8.8), 7.80 (d, 2H, J=8.3), 7.63 (d, 2H, J=8.6), 7.59 (d, 2H, J=8.3), 7.50 (s, 1H), 7.13 (s, 1H), 4.83 (ABq, 2H, Δν=36.2, J <sub>AB</sub> =17.6), 4.37 (dd, 1H, J=8.5, 6.3), 2.09 (m, 1H), 1.88 (m, 1H), 1.64 (m, 1H), 1.45 (m, 1H), 1.27 (m, 2H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
331				1-Method A	white solid	516.07	1.84 Method G	(M+H) <sup>+</sup> 516.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.8), 7.69-7.60 (m, 6H), 7.49 (s, 1H), 7.13 (s, 1H), 4.85 (ABq, 2H, Δν=27.9, J <sub>AB</sub> =17.1), 4.38 (dd, 1H, J=9.0, 5.9), 2.05 (m, 1H), 1.75 (m, 1H), 1.65 (m, 1H), 1.46 (m, 1H), 1.27 (m, 2H).
332				1-Method A	white solid	506.09	1.67 Method G	(M+H) <sup>+</sup> 506.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.90 (d, 2H, J=8.6), 7.83 (d, 2H, J=8.8), 7.62 (d, 2H, J=8.8), 7.54 (d, 2H, J=8.3), 7.47 (s, 1H), 7.11 (s, 1H), 4.84 (ABq, 2H, Δν=36.3, J <sub>AB</sub> =17.4), 4.36 (dd, 1H, J=8.6, 6.1), 3.85 (s, 3H), 2.04 (m, 1H), 1.82 (m, 1H), 1.62 (m, 1H), 1.45 (m, 1H), 1.26 (m, 2H).
333				19	white solid	437.10	1.48 Method G	(M+Na) <sup>+</sup> 459.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.8), 7.79 (d, 2H, J=8.5), 7.63 (d, 2H, J=8.8), 7.58 (d, 2H, J=8.3), 7.52 (s, 1H), 7.09 (s, 1H), 4.82 (ABq, 2H, Δν=37.2, J <sub>AB</sub> =17.6), 4.34 (dd, 1H, J=8.0, 6.6), 4.25 (dt, 2H, J <sub>e</sub> =47.2, J <sub>r</sub> =5.7), 1.58 (m, 1H), 1.49-1.12 (m, 5H).
334				19	white solid	480.09	1.76 Method G	(M+Na) <sup>+</sup> 502.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.80 (d, 2H, J=8.6), 7.67 (d, 2H, J=8.6), 7.60 (m, 4H), 7.52 (s, 1H), 7.09 (s, 1H), 4.83 (ABq, 2H, Δν=30.1, J <sub>AB</sub> =17.4), 4.36 (dd, 1H, J=8.6, 6.2), 4.22 (dt, 2H, J <sub>e</sub> =47.5, J <sub>r</sub> =6.4), 1.61 (m, 1H), 1.48-1.11 (m, 5H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
335				19	white solid	470.11	1.58 Method G	(M+H) <sup>+</sup> 471.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.90 (d, 2H, J=8.3), 7.82 (d, 2H, J=8.8), 7.62 (d, 2H, J=8.8), 7.53 (d, 2H, J=8.4), 7.50 (s, 1H), 7.07 (s, 1H), 4.82 (ABq, 2H, Δν=39.4, J <sub>AB</sub> =17.4), 4.34 (dd, 1H, J=8.3, 6.6), 4.22 (dt, 2H, J <sub>e</sub> =41.6, J <sub>r</sub> =6.1), 3.85 (s, 3H), 1.58 (m, 1H), 1.46-1.12 (m, 5H).
336				19	white solid	423.08	1.43 Method G	(M+H) <sup>+</sup> 423.9	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.8), 7.78 (d, 2H, J=8.3), 7.63 (d, 2H, J=6.8), 7.57 (d, 2H, J=8.6), 7.53 (s, 1H), 7.14 (s, 1H), 4.81 (ABq, 2H, Δν=36.2, J <sub>AB</sub> =17.6), 4.38 (t, 1H, J=7.6), 4.27 (m, 1H), 4.15 (m, 1H), 1.64 (m, 1H), 1.54-1.36 (m, 3H).
337				19	white solid	466.07	1.72 Method G	(M+Na) <sup>+</sup> 489.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.80 (d, 2H, J=8.8), 7.66 (d, 2H, J=8.1), 7.62-7.57 (m, 4H), 7.54 (s, 1H), 7.15 (s, 1H), 4.81 (ABq, 2H, Δν=29.1, J <sub>AB</sub> =17.1), 4.40 (t, 1H, J=6.9), 4.25 (m, 1H), 4.13 (m, 1H), 1.67 (m, 1H), 1.55-1.39 (m, 3H).
338				19	yellow solid	456.09	1.54 Method G	(M+H) <sup>+</sup> 457.0	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.90 (d, 2H, J=8.3), 7.83 (d, 2H, J=8.8), 7.62 (d, 2H, J=8.8), 7.52 (d, 2H, J=8.3), 7.50 (s, 1H), 7.11 (s, 1H), 4.82 (ABq, 2H, Δν=54.5, J <sub>AB</sub> =17.3), 4.37 (t, 1H, J=8.0), 4.28-4.03 (m, 2H), 3.85 (s, 3H), 1.64 (m, 1H), 1.53-1.36 (m, 3H).



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
339				19	white solid	452.86	1.85 min Method A	452.91	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.70 (d, 2H, J = 8.7), 7.54 (d, 2H, J = 8.4), 7.42-7.49 (m, 4H), 6.31 (br s, 1H), 5.23 (br s, 1H), 4.58-4.63 (m, 2H), 4.33-4.41 (m, 2H), 4.19 (t, 1H, J = 4.5), 2.18-2.37 (m, 1H), 1.54-1.66 (m, 1H).
340				19	white solid	442.89	1.68 min Method A	442.90	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.96 (d, 2H, J = 8.4), 7.72 (d, 2H, J = 8.7), 7.48 (d, 2H, J = 8.7), 7.39 (d, 2H, J = 8.4), 6.32 (br s, 1H), 5.18 (br s, 1H), 4.54-4.63 (m, 2H), 4.30-4.42 (m, 2H), 4.16 (t, 1H, J = 4.5), 3.90 (s, 3H), 2.18-2.37 (m, 1H), 1.54-1.66 (m, 1H).
341				19	white solid	409.87	1.57 min Method A	410.07	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.72 (dd, 2H, J = 1.8, 8.7), 7.59 (d, 2H, J = 8.1), 7.50 (dd, 2H, J = 2.1, 8.7), 7.46 (d, 2H, J = 8.1), 6.30 (br s, 1H), 5.21 (br s, 1H), 4.56-4.68 (m, 2H), 4.31-4.37 (m, 2H), 4.18 (t, 1H, J = 4.8), 2.17-2.37 (m, 1H), 1.48-1.64 (m, 1H).
342				19	white solid	470.85	1.88 min Method A	470.89	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.70 (d, 2H, J = 8.7), 7.56 (d, 2H, J = 8.4), 7.51 (d, 2H, J = 8.4), 7.43 (d, 2H, J = 8.1), 6.32 (br s, 1H), 5.75 (tm, 1H, J <sub>H-F</sub> = 57), 5.34 (br s, 1H), 4.52-4.63 (m, 2H), 4.32 (d, 1H, J = 15.6), 2.51-2.66 (m, 1H), 1.54-1.69 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
343				19	white solid	460.88	1.74 min Method A	461.07	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.97 (dd, 2H, J = 2.0, 8.4), 7.72 (d, 2H, J = 8.7), 7.51 (d, 2H, J = 8.7), 7.38 (d, 2H, J = 8.4), 6.32 (br s, 1H), 5.75 (tm, 1H, J <sub>H-F</sub> = 57), 5.18 (br s, 1H), 4.60 (d, 1H, J = 15.6), 4.50-4.55 (m, 1H), 4.32 (d, 1H, J = 15.6), 3.91 (s, 3H), 2.51-2.66 (m, 1H), 1.54-1.69 (m, 1H).
344				19	white solid	427.85	1.62 min Method A	428.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.70 (d, 2H, J = 8.7), 7.61 (d, 2H, J = 8.1), 7.53 (d, 2H, J = 8.7), 7.45 (d, 2H, J = 7.8), 6.32 (br s, 1H), 5.71 (tm, 1H, J <sub>H-F</sub> = 57), 5.21 (br s, 1H), 4.64 (d, 1H, J = 15.6), 4.51-4.55 (m, 1H), 4.28 (d, 1H, J = 15.6), 2.48-2.60 (m, 1H), 1.54-1.69 (m, 1H).
345				1-Method A, sep cond 1	white solid	502.87	1.99 min Method E	503.02	<sup>1</sup> H NMR (DMSO) δ 7.83 (d, 2H, J = 8.8Hz), 7.64 (m, 6H), 7.46 (s br, 1H), 7.25 (s br, 1H) 4.79 (s, 2H), 4.44 (m, 1H), 2.03 (m, 2H), 1.83 (m, 1H), 1.50 (m, 1H).
346				1-Method A, sep cond 2	white solid	459.88	1.76 min Method E	460.13	<sup>1</sup> H NMR (DMSO) δ 7.84 (d, 2H, J = 8.0Hz), 7.80 (d, 2H, J = 7.7Hz), 7.65 (d, 2H, J = 8.2Hz), 7.59 (d, 2H, J = 7.7Hz), 7.47 (s br, 1H), 7.25 (s br, 1H), 4.78 (AB, 2H, Δν = 5Hz, J <sub>A-B</sub> = 17Hz), 4.43 (m, 1H), 2.05 (m, 2H), 1.83 (m, 1H), 1.48 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
347				1-Method A, sep cond 3	yellow waxy solid	437.92	1.67 min Method E	438.22	<sup>1</sup> H NMR (DMSO) δ 7.81 (m, 4H), 7.61 (m, 4H), 7.53 (s br, 1H), 7.09 (s br, 1H), 4.81 (AB <sub>2</sub> , 2H, Δν=5Hz, J <sub>ab</sub> =17Hz), 4.33 (m, 2H), 4.19 (t, 1H, J=6.0Hz), 1.44 (m, 6H)
348				1-Method A, sep cond 3	white powder	480.91	1.94 min Method E	M+Na 503.14	<sup>1</sup> H NMR (DMSO) δ 7.99 (d, 2H, J=8.8Hz), 7.82 (m, 6H), 7.70 (s br, 1H), 7.27 (s br, 1H), 5.00 (m, 2H), 4.51 (m, 2H), 4.33 (m, 1H), 1.77 (m, 1H), 1.47 (m, 6H)
349				1-Method A	white powder	470.94	1.59 min Method B	493.17 M+Na+	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.94 (d, J=8.0Hz, 2H), 7.72 (d, J=6.8Hz, 2H), 7.46 (d, J=6.8Hz, 2H), 7.38 (d, J=8.0Hz, 2H), 6.25 (s, br, 1H), 5.26 (s, br, 1H), 4.37-4.62 (m, 3H), 3.90 (s, 3H), 2.45 (m, 1H), 1.45 (m, 1H), 1.27 (d, J=21.2Hz, 3H), 1.17 (d, J=21.6Hz, 3H)
350				22	white solid	484.88	1.93 min Method E	485.09	<sup>1</sup> H NMR (DMSO) δ 7.81 (d, 2H, J=8.8Hz), 7.63 (m, 6H), 7.51 (s br, 1H), 7.21 (s br, 1H), 5.85 (t, 1H, 5.6Hz), 4.81 (AB <sub>2</sub> , 2H, Δν=5Hz, J <sub>ab</sub> =15Hz), 4.42 (t, 1H, J=8.0Hz), 1.49 (m, 4H)
351				1-Method A	white solid	437.92	1.49 min Method B	460.13 M+Na+	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.72 (d, J=6.8Hz, 2H), 7.58 (d, J=8.4Hz, 2H), 7.48 (d, J=8.4Hz, 2H), 7.45 (d, J=6.8Hz, 2H), 6.25 (s, br, 1H), 5.28 (s, br, 1H), 4.32-4.64 (m, 3H), 2.45 (m, 1H), 1.38 (m, 1H), 1.24 (d, J=21.2Hz, 3H), 1.21 (d, J=22Hz, 3H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
352				1-Method A	white solid	480.91	1.76 min Method B	503.12 M+Na+	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, J=8.4Hz, 2H), 7.52 (d, J=8.0Hz, 2H), 7.45 (d, J=8.4Hz, 2H), 7.43 (d, J=8.0Hz, 2H), 6.30 (s, br, 1H), 5.44 (s, br, 1H), 4.34-4.66 (m, 3H), 2.49 (m, 1H), 1.46 (m, 1H), 1.26 (d, J=21.6Hz, 3H), 1.22 (d, J=21.6Hz, 3H)
353				18	white solid	466.89	1.41 min Method B	466.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.66 (d, 2H, J=8.4Hz), 7.45-7.55 (m, 6H), 6.17 (s, br, 1H), 5.19 (s, br, 1H), 4.53 (dd, 2H, J=50Hz, 15Hz), 4.34-4.37 (m, 2H), 4.22-4.25 (m, 1H), 2.00-2.05 (m, 1H), 1.43-1.49 (m, 3H)
354				18	white solid	423.90	1.54 min Method B	424.1	<sup>1</sup> H NMR (d <sub>6</sub> -DMSO) δ 7.86 (d, 2H, J=8.0Hz), 7.48-7.75 (m, 6H), 6.56 (s, br, 1H), 5.69 (s, br, 1H), 4.72 (dd, 2H, J=42Hz, 16Hz), 4.51-4.55 (m, 3H), 1.43-2.07 (m, 4H)
355				11	yellow solid	582.02	1.47 min Method E	582.22	<sup>1</sup> H NMR (DMSO) δ 7.81 (d, 2H, J=8.8Hz), 7.63 (d, 2H, J=8.5Hz), 7.43 (s br, 1H), 7.24 (m, 4H), 4.64 (s, 2H), 4.42 (m, 3H), 3.99 (m, 2H), 3.62 (m, 6H), 3.23 (m, 2H), 1.85 (m, 4H)
356				11	brown solid	595.06	1.42 min Method E	595.23	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.8Hz), 7.63 (d, 2H, J=8.5Hz), 7.42 (s br, 1H), 7.17 (m, 4H), 4.62 (s, 2H), 4.41 (m, 1H), 4.18 (m, 2H), 3.96 (s, 1H), 3.39 (m, 2H), 3.05 (m, 7H), 2.78 (s, 3H), 1.97 (m, 3H), 1.58 (m, 1H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
357				21	white solid	437.10	1.50 Method G	(M+Na) <sup>+</sup> 460.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.83 (d, 2H, J=8.5), 7.75 (d, 2H, J=8.3), 7.68 (s, 1H), 7.64 (d, 2H, J=8.6), 7.49 (d, 2H, J=8.1), 7.20 (s, 1H), 4.67 (ABq, 2H, Δν=28.3, J <sub>AB</sub> =17.3), 4.54 (dd, 1H, J=9.3, 3.2), 2.23 (m, 1H), 1.42 (m, 1H), 1.25 (d, 3H, J=21.6), 1.21 (d, 3H, J=21.7).
358				21	white solid	480.09	1.76 Method G	(M+Na) <sup>+</sup> 503.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.80 (d, 2H, J=8.6), 7.69 (s, 1H), 7.61 (m, 4H), 7.50 (d, 2H, J=8.1), 7.22 (s, 1H), 4.68 (ABq, 2H, Δν=2.7, J <sub>AB</sub> =17.1), 4.57 (dd, 1H, J=9.1, 3.0), 2.26 (m, 1H), 1.47 (m, 1H), 1.24 (d, 3H, J=21.5), 1.22 (d, 3H, J=21.5).
359				21	white solid	470.11	1.62 Method G	(M+Na) <sup>+</sup> 493.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.86 (d, 2H, J=8.3), 7.83 (d, 2H, J=8.8), 7.63 (m, 3H), 7.44 (d, 2H, J=8.3), 7.18 (s, 1H), 4.67 (ABq, 2H, Δν=10.3, J <sub>AB</sub> =17.1), 4.53 (dd, 1H, J=9.3, 2.9), 3.85 (s, 3H), 2.22 (m, 1H), 1.46 (m, 1H), 1.22 (d, 3H, J=21.5), 1.19 (d, 3H, J=21.6).
360				18	white solid	409.87	1.53min Method B	407.99 (M-H) <sup>+</sup>	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.4Hz), 7.58 (d, 2H, J=8.4Hz), 7.50 (d, 2H, J=8.4Hz), 7.45 (d, 2H, J=8.4Hz), 6.29 (s, br, 1H), 5.21 (s, br, 1H), 4.19-4.67 (m, 5H), 2.17-2.28 (m, 1H), 1.49-1.61 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
361				18	white solid	452.86	1.56min Method B	452.85	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.4Hz), 7.56 (d, 2H, J=8.4Hz), 7.49 (d, 2H, J=8.4Hz), 7.43 (d, 2H, J=8.4Hz), 6.31 (s, br, 1H), 5.24 (s, br, 1H), 4.19-4.62 (m, 5H), 2.16-2.30 (m, 1H), 1.56-1.63 (m, 1H).
362				18	white solid	456.93	1.86 min Method B	457.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.96 (d, 2H, J=8.4), 7.69 (dd, 2H, J=1.8, 8.4), 7.47 (ddd, 2H, J=1.5, 2.1, 8.7), 7.42 (d, 2H, J=8.4), 6.19 (br s, 1H), 5.18 (br s, 1H), 4.64 (d, 1H, J=15.6), 4.42 (d, 1H, J=15.9), 4.30-4.35 (m, 2H), 4.18 (t, 1H, J=3.6), 3.90 (s, 3H), 1.89-2.08 (m, 1H), 1.38-1.50 (m, 3H).
363				18	white solid	456.97	1.80 min Method B	454.98 (neg. ion)	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.64 (d, 2H, J=8.7), 7.42 (d, 2H, J=8.7), 7.38 (d, 2H, J=8.4), 7.26 (d, 2H, J=8.4), 6.19 (br, 1H), 5.28 (br s, 1H), 4.51 (d, 1H, J=15.6), 4.39 (d, 1H, J=15.3), 4.30-4.35 (m, 2H), 4.18 (t, 1H, J=3.6), 1.92-2.08 (m, 1H), 1.55 (s, 6H), 1.35-1.50 (m, 3H).
364				18	white solid	442.90	1.73 min Method B	443.12	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.81 (d, 2H, J=8.4), 7.59 (dd, 2H, J=1.8, 8.4), 7.29-7.33 (m, 4H), 6.76 (br, 1H), 5.80 (br s, 1H), 4.62 (d, 1H, J=16.2), 4.44 (d, 1H, J=16.2), 4.31 (t, 1H, J=6.9), 3.85-4.10 (m, 2H), 1.70-1.85 (m, 1H), 1.30-1.48 (m, 3H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
365				11, sep cond 4	off-white solid	585.02	1.47 min Method E	582.18	<sup>1</sup> H NMR (DMSO) δ 7.80 (m, 2H), 7.62 (m, 2H), 7.42 (m, 1H), 7.18 (m, 4H), 4.62 (m, 3H), 4.42 (m, 1H), 4.14 (m, 1H), 4.00 (m, 3H), 3.58 (m, 4H), 2.93 (m, 1H), 2.70 (m, 2H), 2.01 (m, 2H), 1.85 (m, 1H), 1.59 (m, 1H)
366				11, sep cond 4	yellow solid	595.06	1.43 min Method E	595.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.72 (d, 2H, J=8.7Hz), 7.52 (d, 2H, J=8.8Hz), 7.14 (m, 1H), 7.01 (m, 1H), 6.87 (m, 1H), 6.24 (s br, 1H), 5.36 (m, 1H), 4.51 (m, 1H), 4.28 (m, 4H), 3.23 (m, 9H), 2.77 (m, 3H), 1.94 (m, 3H), 1.40 (m, 1H)
367				11, sep cond 4	white solid	595.06	1.42 min Method E	595.20	<sup>1</sup> H NMR (DMSO) δ 7.80 (m, 2H), 7.62 (m, 2H), 7.42 (s br, 1H), 7.18 (m, 4H), 4.61 (m, 2H), 4.42 (m, 1H), 4.14 (m, 2H), 3.72 (m, 1H), 3.57 (m, 1H), 3.32 (s, 3H), 2.75 (m, 6H), 2.27 (m, 2H), 1.97 (m, 3H), 1.60 (m, 1H)
368				11, sep cond 4	dark yellow solid	546.04	1.55 min Method E	546.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.5Hz), 7.61 (d, 2H, J=8.5Hz), 7.53 (m, 1H), 7.19 (m, 4H), 4.67 (ABq, 2H, Δν=35, J <sub>ab</sub> =16Hz), 4.30 (m, 6H), 4.01 (m, 4H), 3.48 (m, 4H), 1.68 (m, 1H), 1.49 (m, 4H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
369				11, sep cond 4	orange-yellow solid	559.08	1.29 min Method E	559.22	<sup>1</sup> H NMR (DMSO) δ 7.77 (d, 2H, J=8.3Hz), 7.60 (d, 2H, J=8.3Hz), 7.50 (s br, 1H), 7.16 (m, 4H), 4.65 (ABq, 2H, Δν=20, J <sub>ab</sub> =16Hz), 4.33 (m, 2H), 4.18 (m, 4H), 3.17 (m, 7H), 2.78 (s, 3H), 1.67 (m, 1H), 1.50 (m, 5H)
370				11, sep cond 4	yellow solid	528.05	1.32 min Method E	528.17	<sup>1</sup> H NMR (DMSO) δ 7.74 (d, 2H, J=8.3Hz), 7.60 (d, 2H, J=8.3Hz), 7.44 (s br, 1H), 7.35 (d, 2H, J=8.5Hz), 7.08 (s br, 1H), 6.95 (d, 2H, J=8.5Hz), 4.68 (ABq, 2H, Δν=24, J <sub>ab</sub> =16Hz), 3.77 (m, 13H), 1.66 (m, 1H), 1.45 (m, 5H)
371				11, sep cond 4	light-orange solid	541.09	1.26 min Method E	541.24	<sup>1</sup> H NMR (DMSO) δ 7.77 (d, 2H, J=8.3Hz), 7.59 (d, 2H, J=8.3Hz), 7.43 (s br, 1H), 7.31 (d, 2H, J=8.5Hz), 7.08 (s br, 1H), 6.87 (d, 2H, J=8.8Hz), 4.66 (ABq, 2H, Δν=16, J <sub>ab</sub> =16Hz), 4.22 (m, 5H), 3.17 (m, 8H), 2.77 (m, 3H), 1.65 (m, 1H), 1.46 (m, 5H)
372				18, 8	amber glass	511.05	1.09 min Method A	511.21	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.66 (d, 2H, J=8.0Hz), 7.43 (d, 2H, J=8.0Hz), 7.26 (d, 2H, J=8.0Hz), 7.21 (d, 2H, J=8.0Hz), 6.25 (s br, 1H), 5.45 (s br, 1H), 4.52 (d, 1H, J <sub>ab</sub> =12.0Hz), 4.38 (d, 1H, J <sub>ab</sub> =12.0Hz), 4.29 (m, 2H), 4.18 (t, 1H, J=6.0Hz), 3.72 (m, 1H), 3.47 (s, 2H), 2.29 (s, 3H), 2.0 (m, 2H), 1.83 (m, 2H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
373				18, 8	amber glass	498.01	1.40 min Method A	498.20	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.66 (d, 2H, J=8.0Hz), 7.44 (d, 2H, J=8.0Hz), 7.27 (d, 2H, J=8.0Hz), 7.22 (d, 2H, J=8.0Hz), 6.24 (s, br, 1H), 5.42 (s, br, 1H), 4.52 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.38 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.29 (m, 2H), 4.18 (m, 1H), 3.68 (t, 4H, J=4.0Hz), 3.45 (s, 2H), 2.40 (s, br, 4H), 1.99 (m, 1H), 1.45 (m, 3H)
374				18, 8	amber glass	529.08	1.17 min Method A	529.22	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J=8.0Hz), 7.46 (d, 2H, J=8.0Hz), 7.25 (t, 1H, J=6.0Hz), 7.05 (t, 1H, J=6.0Hz), 6.28 (s, br, 1H), 5.49 (s, br, 1H), 4.54 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.33 (m, 2H), 4.22 (m, 1H), 3.57 (s, 2H), 2.32 (s, 3H), 2.01 (m, 1H), 1.44 (m, 2H)
375				18, 8	amber glass	516.00	1.10 min Method A	516.17	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.68 (d, 2H, J=8.0Hz), 7.45 (d, 2H, J=8.0Hz), 7.28 (t, 1H, J=6.0Hz), 7.06 (t, 2H, J=6.0Hz), 6.24 (s, br, 1H), 5.40 (s, br, 1H), 4.53 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.36 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.34 (m, 2H), 4.21 (m, 1H), 3.69 (t, 4H, J=4.0Hz), 3.53 (s, 2H), 2.44 (s, br, 3H), 2.0 (m, 1H), 1.45 (m, 2H)
376				18	white solid	451.91	1.39min Method B	451.90	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.55 (s, 1H), 8.09 (s, 1H), 7.75 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.4Hz), 7.47-7.51 (m, 4H), 6.36 (s, br, 1H), 5.28 (s, br, 1H), 4.31-4.62 (m, 5H), 2.16-2.30 (m, 1H), 1.56-1.66 (m, 1H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
377				22	white solid	441.89	1.65 min Method E	M+Na 464.01	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.3Hz), 7.60 (d, 2H, J=8.3Hz), 7.49 (m, 4H), 6.18 (s, br, 1H), 5.67 (t, 1H, J=5.6Hz, 4.0Hz), 5.22 (s, br, 1H), 4.52 (AB <sub>2</sub> , 2H, Δν=16, J <sub>AB</sub> =100Hz), 4.34 (m, 1H), 2.03 (m, 1H), 1.68 (m, 1H), 1.38 (m, 1H), 0.86 (m, 1H)
378				18	white solid	450.92	1.52min Method B	450.91	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.91 (s, 1H), 7.71 (m, 3H), 7.63 (d, 2H, J=8.4Hz), 7.50 (d, 2H, J=8.4Hz), 7.41 (d, 2H, J=8.4Hz), 6.47 (s, 1H), 6.34 (s, br, 1H), 5.18 (s, br, 1H), 4.30-4.60 (m, 5H), 2.14-2.29 (m, 1H), 1.56-1.66 (m, 1H)
379				18	white solid	468.96	1.53min Method B	469.04	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.64 (s, 1H), 7.98 (d, 2H, J=8.4Hz), 7.75 (d, 2H, J=8.4Hz), 7.45-7.50 (m, 4H), 6.35 (s, br, 1H), 5.20 (s, br, 1H), 4.22-4.64 (m, 5H), 2.20-2.35 (m, 1H), 1.54-1.62 (m, 1H)
380				18	white solid	464.95	1.54min Method B	464.99	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.54 (s, 1H), 8.10 (s, 1H), 7.73 (d, 2H, J=8.4Hz), 7.62 (d, 2H, J=8.4Hz), 7.48-7.52 (m, 5H), 6.22 (s, br, 1H), 5.18 (s, br, 1H), 4.32-4.69 (m, 5H), 2.09-2.19 (m, 1H), 1.44-1.61 (m, 3H)
381				18	white solid	465.94	1.49min Method B	465.96	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.55 (s, 1H), 8.09 (s, 1H), 7.73 (d, 2H, J=8.4Hz), 7.62 (d, 2H, J=8.4Hz), 7.47-7.51 (m, 4H), 6.36 (s, br, 1H), 5.28 (s, br, 1H), 4.31-4.66 (m, 5H), 2.10-2.39 (m, 1H), 1.56-1.66 (m, 3H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/Method	M+H <sup>+</sup>	NMR Data
382				1-Method A	white solid	501.92	1.53min Method B	502.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.56 (s, 1H), 8.10 (s, 1H), 7.72 (d, 2H, J=8.4Hz), 7.64 (d, 2H, J=8.4Hz), 7.49-7.52 (m, 4H), 6.23 (s, br, 1H), 5.22 (s, br, 1H), 4.32-4.64 (m, 3H), 1.44-2.20 (m, 4H).
383				1-Method A	white solid	500.93	1.66min. Method B	501.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.91 (s, 1H), 7.41-7.78 (m, 9H), 6.47 (s, 1H), 6.23 (s, br, 1H), 5.27 (s, br, 1H), 4.30-4.59 (m, 3H), 1.47-2.21 (m, 4H).
384				18, 11	beige solid	532.01	1.36 min Method E	532.18	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.3Hz), 7.61 (d, 2H, J=8.3Hz), 7.56 (s br, 1H), 7.23 (s br, 1H), 7.19 (m, 3H), 4.64 (ABq, 2H, Δv=16, J <sub>ab</sub> =16Hz), 4.51 (t, 1H, J=8.0Hz), 4.42 (m, 2H), 4.03 (m, 12H), 2.02 (m, 1H), 1.82 (m, 1H).
385				18, 11	pale orange solid	545.05	1.32 min Method E	545.25	<sup>1</sup> H NMR (DMSO) δ 7.79 (d, 2H, J=8.3Hz), 7.60 (d, 2H, J=8.3Hz), 7.54 (s br, 1H), 7.14 (m, 5H), 4.62 (ABq, 2H, Δv=8.0Hz, J <sub>ab</sub> =16Hz), 4.51 (t, 1H, J=8.0Hz), 4.24 (m, 5H), 3.17 (m, 8H), 2.78 (m, 3H), 2.02 (m, 1H), 1.81 (m, 1H).
386				18, 11	yellow residue	514.02	1.33 min Method E	514.20	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.3Hz), 7.61 (d, 2H, J=8.3Hz), 7.48 (s br, 1H), 7.33 (d, 2H, J=8.3Hz), 7.15 (s br, 1H), 6.95 (d, 2H, J=8.3Hz), 4.65 (ABq, 2H, Δv=8.0Hz, J <sub>ab</sub> =16Hz), 3.86 (m, 15H), 2.03 (m, 1H), 1.79 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/Method	M+H <sup>+</sup>	NMR Data
387				18, 11	orange solid	527.06	1.28 min Method E	527.20	<sup>1</sup> H NMR (DMSO) δ 7.79 (d, 2H, J=8.3Hz), 7.60 (d, 2H, J=8.3Hz), 7.47 (s br, 1H), 7.30 (d, 2H, J=8.3Hz), 7.15 (s br, 1H), 6.89 (d, 2H, J=8.3Hz), 4.63 (ABq, 2H, Δv=8.0Hz, J <sub>ab</sub> =16Hz), 4.48 (t, 1H, J=8.0Hz), 4.15 (m, 4H), 3.24 (m, 10H), 2.79 (m, 3H), 2.03 (m, 1H), 1.80 (m, 1H).
388				18, 11	tan solid	560.06	1.65 min Method E	559.14	<sup>1</sup> H NMR (DMSO) δ 7.76 (d, 2H, J=8.3Hz), 7.59 (m, 3H), 7.21 (s br, 1H), 7.08 (s br, 1H), 7.04 (m, 2H), 4.53 (m, 5H), 3.92 (m, 4H), 3.45 (m, 6H), 2.25 (m, 1H), 1.54 (m, 1H), 1.25 (d, 3H, J=20Hz), 1.22 (d, 3H, J=20Hz).
389				18	yellow solid	478.91	1.55min Method B	491.04 (M <sup>+</sup> Na)	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.92 (s, 1H), 7.72-7.76 (m, 3H), 7.65 (d, 2H, J=8.4Hz), 7.52 (d, 2H, J=8.4Hz), 7.40 (d, 2H, J=8.4Hz), 6.47 (s, 1H), 6.30 (s, br, 1H), 5.21 (s, br, 1H), 4.27-4.57 (m, 3H), 2.51-2.60 (m, 1H), 1.53-1.65 (m, 2H).
390				18	white solid	469.90	1.48min Method B	470.1	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.90 (s, 1H), 7.40-7.76 (m, 9H), 6.50 (s, 1H), 6.27 (s, br, 1H), 5.22 (s, br, 1H), 4.25-4.55 (m, 3H), 2.49-2.58 (m, 1H), 1.51-1.62 (m, 2H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
391				18, 6	white solid	469.97	1.59 min Method A	470.09	<sup>1</sup> H NMR (MeOD, 400MHz) δ 7.77 (ddd, 2H, J = 2.0, 2.4, 8.8), 7.71 (d, 2H, J = 8.4), 7.51 (ddd, 2H, J = 2.0, 2.4, 8.8), 7.48 (d, 2H, J = 8.0), 4.80 (d, 1H, J = 16.4), 4.72 (d, 1H, J = 16.4), 4.47 (t, 1H, J = 7.2), 4.05-4.28 (m, 2H), 3.38 (q, 2H, J = 7.2), 1.70-1.85 (m, 1H), 1.30-1.48 (m, 3H), 1.20 (t, 3H, J = 7.2).
392				18, 6	white solid	500.00	1.57 min Method A	500.10	<sup>1</sup> H NMR (MeOD, 400MHz) δ 7.72-7.81 (m, 4H), 7.48-7.52 (m, 4H), 4.80 (d, 1H, J = 16.4), 4.72 (d, 1H, J = 16.4), 4.47 (t, 1H, J = 7.2), 4.05-4.28 (m, 2H), 3.54 (s, 3H), 3.25-2.36 (m, 4H), 1.70-1.85 (m, 1H), 1.30-1.48 (m, 3H).
393				18, 6	white solid	484.00	1.65 min Method A	484.12	<sup>1</sup> H NMR (MeOD, 400MHz) δ 7.79 (d, 2H, J = 8.8), 7.50-7.54 (m, 4H), 7.29-7.33 (m, 2H), 4.79 (d, 1H, J = 16.4), 4.72 (d, 1H, J = 16.4), 4.47 (t, 1H, J = 7.2), 4.05-4.28 (m, 2H), 3.52-3.56 (m, 2H), 3.03 and 2.94 (2 s, 3H), 1.70-1.85 (m, 1H), 1.30-1.48 (m, 3H), 1.15-1.25 (m, 3H).
394				18, 6	white solid	533.03	1.36 min Method A	533.15	<sup>1</sup> H NMR (MeOD, 400MHz) δ 8.48 (d, 1H, J = 4.8), 7.76-7.82 (m, 5H), 7.50-7.53 (m, 4H), 7.41 (d, 1H, J = 8.0), 7.28-7.30 (m, 1H), 4.82 (d, 1H, J = 16.0), 4.72 (d, 1H, J = 16.4), 4.67 (s, 2H), 4.48 (t, 1H, J = 7.2), 4.05-4.28 (m, 2H), 1.70-1.85 (m, 1H), 1.30-1.48 (m, 3H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
395				18, 8	amber glass	501.98	1.17 min Method A	502.25	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 7.42 (t, 1H, J=6.0Hz), 7.26 (d, 1H, J=6.0Hz), 7.14 (d, 1H, J=6.0Hz), 6.35 (s, br, 1H), 6.07 (s, br, 1H), 4.82 (d, 1H, J <sub>ab</sub> =12.0Hz), 4.35 (m, 1H), 4.19 (m, 5H), 3.59 (s, br, 4H), 3.48 (d, br, 1H), 2.90 (d, br, 1H), 2.20 (m, 1H), 1.50 (m, 1H).
396				18, 8	amber glass	515.02	1.21 min Method A	515.27	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.76 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 7.32 (m, 1H), 7.19 (d, 1H, J=8.0Hz), 7.07 (d, 1H, J=8.0Hz), 6.35 (s, br, 1H), 5.67 (s, br, 1H), 4.86 (d, 1H, J <sub>ab</sub> =12.0Hz), 4.70 (s, 2H), 4.56 (m, 2H), 4.17 (m, 6H), 2.80 (s, 3H), 2.22 (m, 1H), 1.54 (m, 1H).
397				18, 8	amber glass	483.98	1.21 min Method A	483.98	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 7.41 (d, 2H, J=8.0Hz), 7.31 (d, 2H, J=8.0Hz), 6.34 (s, br, 1H), 6.05 (s, br, 1H), 4.83 (d, 1H, J <sub>ab</sub> =12.0Hz), 4.51 (m, 1H), 4.20 (m, 4H), 3.94 (m, 5H), 3.51 (d, 1H, J=12.0Hz), 3.40 (d, 1H, J=12.0Hz), 2.89 (t, 1H, J=6.0Hz), 2.76 (t, 1H, J=6.0Hz), 2.20 (m, 1H), 1.51 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
398				18, 8	amber glass	497.03	1.16 min Method A	498.74	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.76 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 7.37 (d, 2H, J=8.0Hz), 7.27 (d, 2H, J=8.0Hz), 6.31 (s, br, 1H), 5.80 (s, br, 1H), 4.82 (d, 1H, J <sub>AB</sub> =12.0Hz), 4.55 (m, 2H), 4.16 (m, 5H), 3.82 (d, 1H), 2.78 (s, 3H), 2.21 (m, 1H), 1.52 (m, 1H)
399				20, 11	colorless residue	573.11	1.63 min Method E		<sup>1</sup> H NMR (DMSO) δ 7.76 (d, 2H, J=8.3Hz), 7.59 (m, 3H), 7.20 (s br, 1H), 7.08 (s br, 1H), 7.03 (m, 2H), 4.53 (m, 11H), 3.92 (m, 4H), 3.45 (m, 3H), 2.24 (m, 1H), 1.54 (m, 1H), 1.25 (d, 3H, J=20Hz), 1.22 (d, 3H, J=20Hz)
400				11	light yellow solid	564.03	1.51min Method E	564.19	<sup>1</sup> H NMR (DMSO) δ 7.81 (d, 2H, J=8.3Hz), 7.63 (d, 2H, J=8.3Hz), 7.36 (m, 3H), 7.20 (s br, 1H), 6.96 (d, 2H, J=8.0Hz), 4.64 (s, 2H), 4.37 (m, 3H), 4.00 (m, 2H), 3.61 (m, 6H), 3.21 (m, 2H), 1.97 (m, 2H), 1.82 (m, 1H), 1.60 (m, 1H)
401				11	pale orange solid	577.07	1.47min Method E	577.20	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.0Hz), 7.33 (m, 3H), 7.20 (s br, 1H), 6.89 (d, 2H, J=8.0Hz), 4.63 (s, 2H), 4.39 (m, 1H), 4.10 (m, 2H), 3.25 (m, 10H), 2.77 (s, 3H), 1.89 (m, 2H), 1.81 (m, 1H), 1.60 (m, 1H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
402				20, 11	pale yellow residue	542.07	1.51min Method E	542.21	<sup>1</sup> H NMR (DMSO) δ 7.77 (d, 2H, J=8.0Hz), 7.59 (d, 2H, J=8.0Hz), 7.55 (s br, 1H), 7.26 (d, 2H, J=8.0Hz), 7.15 (s br, 1H), 6.90 (d, 2H, J=8.0Hz), 4.52 (m, 2H), 4.32 (m, 2H), 4.00 (m, 2H), 3.71 (m, 2H), 3.38 (m, 7H), 2.23 (m, 1H), 1.55 (m, 1H), 1.23 (d, 3H, J=20Hz), 1.21 (d, 3H, J=20Hz)
403				20, 11	light orange solid	555.12	1.45min Method E	555.27	<sup>1</sup> H NMR (DMSO) δ 7.76 (d, 2H, J=8.0Hz), 7.58 (d, 2H, J=8.0Hz), 7.54 (s br, 1H), 7.22 (d, 2H, J=8.0Hz), 7.16 (s br, 1H), 6.84 (d, 2H, J=8.0Hz), 4.52 (m, 2H), 4.13 (m, 2H), 3.25 (m, 11H), 2.79 (m, 3H), 2.20 (m, 1H), 1.56 (m, 1H), 1.23 (d, 3H, J=20Hz), 1.20 (d, 3H, J=20Hz)
404				18, 11	pale yellow residue	532.01	1.43min Method E	532.18	<sup>1</sup> H NMR (DMSO) δ 7.81 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.0Hz), 7.48 (s br, 1H), 7.34 (d, 2H, J=8.0Hz), 7.25 (s br, 1H), 6.96 (d, 2H, J=8.0Hz), 5.72 (t, 1H, J=8.0Hz, 56Hz), 4.61 (s, 2H), 4.52 (t, 1H, J=8.0Hz), 4.33 (t, 2H, J=8.0Hz), 3.96 (m, 2H), 3.56 (m, 2H), 3.22 (m, 2H), 2.24 (m, 1H), 1.95 (m, 1H)



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
405				18, 11	pale yellow solid	545.05	1.37min Method E	545.19	<sup>1</sup> H NMR (DMSO) δ 7.81 (d, 2H, J=8.0Hz), 7.61 (d, 2H, J=8.0Hz), 7.46 (s br, 1H), 7.30 (d, 2H, J=8.0Hz), 7.25 (s br, 1H), 6.89 (d, 2H, J=8.0Hz), 5.64 (m, 1H), 4.60 (s, 2H), 4.52 (t, 1H, J=8.0Hz), 4.11 (m, 2H), 3.17 (m, 10H), 2.77 (m, 3H), 2.24 (m, 1H), 1.96 (m, 1H)
406				18, 11	yellow residue	550.00	1.43min Method E	550.16	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.0Hz), 7.56 (s br, 1H), 7.23 (m, 4H), 5.80 (t, 1H, J=4.0Hz, 56Hz), 3.93 (m, 15H), 2.29 (m, 1H), 1.99 (m, 1H)
407				18, 11	orange solid	563.04	1.37min Method E	563.21	<sup>1</sup> H NMR (DMSO) δ 7.80 (d, 2H, J=8.0Hz), 7.62 (d, 2H, J=8.0Hz), 7.55 (s br, 1H), 7.29 (s br, 1H), 7.15 (m, 3H), 5.78 (t, 1H, J=4.0Hz, 56Hz), 4.56 (m, 3H), 4.21 (m, 2H), 3.22 (m, 10H), 2.79 (m, 3H), 2.30 (m, 1H), 1.97 (m, 1H)
408				18, 10, sep cond 5	clear oil	442.94	0.993 min Method B	443.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.07 (s, 1H), 7.67 (d, J=6.8Hz, 2H), 7.55 (d, J=8.8Hz, 1H), 7.44 (d, J=6.8Hz, 2H), 6.47 (d, J=8.8Hz, 1H), 6.30 (s, br, 1H), 5.45 (s, br, 1H), 4.20-4.46 (m, 5H), 3.12 (s, 6H), 1.22-1.85 (m, 4H)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
409				18, 10, sep cond 5	clear oil	442.94	0.993 min Method B	443.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.07 (s, 1H), 7.67 (d, J=6.8Hz, 2H), 7.55 (d, J=8.8Hz, 1H), 7.44 (d, J=6.8Hz, 2H), 6.47 (d, J=8.8Hz, 1H), 6.30 (s, br, 1H), 5.45 (s, br, 1H), 4.20-4.46 (m, 5H), 3.12 (s, 6H), 1.22-1.85 (m, 4H)
410				20, 8	amber glass	512.04	1.44 min Method B	512.20	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.42 (d, 2H, J=8.0Hz), 7.22 (dd, 4H, J=8.0Hz), 6.27 (s, br, 1H), 5.26 (s, br, 1H), 4.58 (d, 1H), 4.43 (d, 1H, J <sub>AB</sub> =16.0Hz), 4.36 (d, 1H, J <sub>AB</sub> =16.0Hz), 3.68 (m, 6H), 3.46 (s, 2H), 2.45 (m, 6H), 1.53 (m, 1H), 1.26 (d, 3H, J=20.0Hz), 1.17 (d, 3H, J=20.0Hz)
411				20, 8	amber glass	525.08	1.38 min Method B	525.24	<sup>1</sup> H NMR, 400Hz, (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.42 (d, 2H, J=8.0Hz), 7.22 (dd, 4H, J=8.0Hz), 6.26 (s, br, 1H), 5.24 (s, br, 1H), 4.57 (m, 1H), 4.43 (d, 1H, J <sub>AB</sub> =16.0Hz), 4.35 (d, 1H, J <sub>AB</sub> =16.0Hz), 3.48 (s, 2H), 2.47 (m, 7H), 2.30 (s, 3H), 1.26 (d, 3H, J=22.0Hz), 1.18 (d, 3H, J=22.0Hz)
412				18	white solid	427.87	1.39 min Method B	428.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (dd, 2H, J=2.0, 8.8), 7.61 (d, 2H, J=8.4), 7.53 (dd, 2H, J=2.0, 8.4), 7.45 (d, 2H, J=8.0), 6.32 (br s, 1H), 5.73 (m, 1H, J <sub>4,5</sub> =57), 5.22 (br s, 1H), 4.64 (d, 1H, J=16.4), 4.50-4.60 (m, 1H), 4.28 (d, 1H, J=16.4), 2.42-2.60 (m, 1H), 1.50-1.63 (m, 1H)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
413				18	white solid	470.86	1.69 min Method B	471.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.69 (dd, 2H, J = 1.6, 8.8), 7.56 (d, 2H, J = 8.4), 7.51 (ddd, 2H, J = 2.0, 2.4, 8.4), 7.43 (d, 2H, J = 8.4), 6.32 (br s, 1H), 5.75 (m, 1H, J <sub>BF</sub> = 57), 5.25 (br s, 1H), 4.60 (d, 1H, J = 15.6), 4.50-4.60 (m, 1H), 4.32 (d, 1H, J = 15.6), 2.50-2.60 (m, 1H), 1.55-1.70 (m, 1H).
414				18, 13	white solid	480.95	1.41 min Method B	481.22	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.96 (d, 2H, J = 8.4), 7.70 (d, 2H, J = 8.4), 7.47-7.50 (m, 4H), 6.22 (br s, 1H), 5.18 (br s, 1H), 4.66 (d, 1H, J = 15.6), 4.43 (d, 1H, J = 15.6), 4.30-4.35 (m, 2H), 4.19-4.21 (m, 1H), 2.61 (s, 3H), 1.93-2.08 (m, 1H), 1.38-1.50 (m, 3H).
415				18, 10	white foam	428.92	1.34 Method B	429.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt δ 8.10 (s, 1H), 8.04 (d, 1H, J = 9.2Hz), 7.76 (d, 2H, J = 6.8Hz), 7.52 (d, 2H, J = 6.8Hz), 6.80 (d, 2H, J = 9.2Hz), 6.46 (s, 1H), 6.00 (s, 1H), 4.60 (d, 1H, J = 15.6Hz), 4.54 (dd, 1H, J = 5.2Hz, 6.2Hz), 4.30 (m, 1H), 4.18 (m, 1H), 4.07 (d, 1H, J = 15.6Hz), 3.29 (s, 6H), 2.25 (m, 1H), 1.55 (m, 1H).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
416				1-Method A	off-white solid	463.91	1.50 min Method B	464.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.72 (d, 1H, J = 2.4), 7.82 (dd, 1H, J = 2.7, 8.4), 7.55 (d, 2H, J = 8.1), 7.47 (d, 2H, J = 8.1), 7.37 (d, 1H, J = 8.4), 5.97 (br s, 1H), 5.26 (br s, 1H), 4.62 (d, 1H, J = 16.2), 4.54 (d, 1H, J = 15.9), 4.44 (t, 1H, J = 7.5), 1.70-1.77 (m, 1H), 1.35-1.43 (m, 1H), 1.21-1.31 (m, 1H), 0.85 (d, 3H, J = 6.3), 0.67 (d, 3H, J = 6.6).
417				1-Method A	white solid	453.95	1.37 min Method B	454.14	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.71 (d, 1H, J = 3.0), 7.96 (d, 2H, J = 8.1), 7.85 (dd, 1H, J = 2.4, 8.4), 7.37-7.43 (m, 3H), 5.99 (br s, 1H), 5.24 (br s, 1H), 4.61 (d, 1H, J = 15.9), 4.53 (d, 1H, J = 15.9), 4.41 (t, 1H, J = 7.2), 3.91 (s, 3H), 1.70-1.76 (m, 1H), 1.35-1.43 (m, 1H), 1.21-1.31 (m, 1H), 0.82 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).
418				1-Method A	white solid	420.92	1.64 min Method A	421.15	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.74 (d, 1H, J = 2.4), 7.90 (dd, 1H, J = 2.4, 8.4), 7.61 (d, 2H, J = 8.4), 7.50 (d, 2H, J = 8.4), 7.44 (d, 1H, J = 8.4), 5.92 (br s, 1H), 5.22 (br s, 1H), 4.65 (d, 1H, J = 16.5), 4.52 (d, 1H, J = 16.5), 4.40 (t, 1H, J = 7.5), 1.65-1.74 (m, 1H), 1.33-1.40 (m, 1H), 1.18-1.25 (m, 1H), 0.83 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
419				7	white solid	528.12	1.35 min Method A	528.26	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J=8.6), 7.51 (d, 2H, J=8.8), 6.65 (s, 1H), 5.35 (s, 1H), 4.14 (dd, 1H, J=5.1, 9.5), 3.63-3.75 (m, 2H), 3.35-3.55 (m, 3H), 3.25 (dd, 1H, J=9.8, 14), 2.98 (dd, 1H, J=4.5, 14), 2.35-2.87 (m, 8H), 1.77-1.92 (m, 3H), 1.59 (d, 2H, J=13), 1.05-1.30 (m, 3H), 0.75-0.80 (m, 1H), 0.72 (d, 3H, J=6.4), 0.67 (d, 3H, J=6.7).
420				7	white solid	515.08	1.56 min Method A	515.33	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.71 (d, 2H, J=8.5), 7.50 (d, 2H, J=8.5), 6.65 (s, 1H), 5.47 (s, 1H), 4.13 (dd, 1H, J=5.2, 9.2), 3.60-3.75 (m, 6H), 3.15-3.30 (m, 6H), 2.97 (dd, 1H, J=4.6, 14), 2.71 (dd, 2H, J=14, 25), 1.75-1.92 (m, 3H), 1.67 (d, 1H, J=12), 1.05-1.30 (m, 3H), 0.75-0.81 (m, 1H), 0.71 (d, 3H, J=6.7), 0.65 (d, 3H, J=6.7).
421				7	white solid	460.00	1.68 min Method A	460.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J=8.5), 7.50 (d, 2H, J=8.6), 6.65 (s, 1H), 5.35 (s, 1H), 4.15 (dd, 3H, J=5.2, 9.5), 3.67 (s, 3H), 3.25 (t, 1H, J=10), 3.97 (dd, 1H, J=4.8, 14), 2.61-2.80 (m, 2H), 1.74-1.94 (m, 3H), 0.89-1.40 (m, 4H), 0.75-0.80 (m, 1H), 0.72 (d, 3H, J=6.4), 0.67 (m, 3H, J=6.7).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
422				1-Method A	tan solid	491.81	1.84 min Method B	491.04	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J=8.0Hz), 7.50-7.58 (m, 2H), 7.41 (d, 2H, J=8.0Hz), 7.17-7.22 (m, 1H), 6.23 (s, br, 1H), 5.25 (s, br, 1H), 4.41 (dd, 2H, J=50Hz, 15Hz), 4.31-4.35 (m, 1H), 1.80-1.86 (m, 1H), 1.41-1.44 (m, 1H), 1.11-1.15 (m, 1H), 0.81 (d, 3H, J=7.0Hz), 0.71 (d, 3H, J=7.0Hz).
423				1-Method A	white solid	424.09	1.54 min Method F	(M+H) <sup>+</sup> 425.1	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.90 (d, 2H, J=8.1), 7.82 (d, 2H, J=8.5), 7.61 (d, 2H, J=8.5), 7.52 (d, 2H, J=8.1), 7.49 (s, 1H), 7.07 (s, 1H), 4.81 (ABq, 2H, Δν=45.3, J <sub>AB</sub> =17.3), 4.27 (t, 1H, J=7.3), 3.85 (s, 3H), 1.55 (m, 1H), 1.38 (m, 1H), 0.68 (t, 3H, J=7.3).
424				1-Method A	white solid	452.12	1.69 min Method F	(M+H) <sup>+</sup> 453.1	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.90 (d, 2H, J=8.3), 7.83 (d, 2H, J=8.8), 7.62 (d, 2H, J=8.8), 7.54 (d, 2H, J=8.3), 7.48 (s, 1H), 7.04 (s, 1H), 4.82 (ABq, 2H, Δν=41.3, J <sub>AB</sub> =17.3), 4.31 (t, 1H, J=8.1), 3.85 (s, 3H), 1.52 (m, 1H), 1.29 (m, 1H), 1.04 (m, 3H), 0.90 (m, 1H), 0.63 (t, 3H, J=7.3).










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
425				1-Method A	pale yellow solid	460.99	1.48 min Method B	460.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.91 (d, 1H, J = 1.5), 7.67-7.72 (m, 3H), 7.62 (d, 2H, J = 8.7), 7.42-7.48 (m, 4H), 6.46 (t, 1H, J = 2.1), 6.24 (br s, 1H), 5.21 (br s, 1H), 4.62 (d, 1H, J = 15.3), 4.42 (d, 1H, J = 15.6), 4.29 (t, 1H, J = 6.9), 1.80-1.88 (m, 1H), 1.28-1.40 (m, 1H), 1.12-1.21 (m, 1H), 0.75 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).
426				1-Method A	white solid	461.97	1.39 min Method B	462.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.54 (s, 1H), 8.10 (s, 1H), 7.71 (d, 2H, J = 8.4), 7.61 (d, 2H, J = 8.4), 7.46-7.52 (m, 4H), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.66 (d, 1H, J = 15.9), 4.42 (d, 1H, J = 15.9), 4.30 (t, 1H, J = 6.9), 1.79-1.89 (m, 1H), 1.30-1.38 (m, 1H), 1.07-1.14 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.66 (d, 3H, J = 6.6).
427				11	pale yellow solid	542.07	1.48 min Method E	542.25	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J = 8.00Hz), 7.49 (d, 2H, J = 8.0Hz), 7.16 (d, 1H, J = 12.0Hz), 7.05 (d, 1H, J = 8.0Hz), 6.86 (t, 1H, J = 8.0Hz), 6.40 (s, br, 1H), 5.87 (s br, 1H), 4.42 (ABq, 2H, Δν = 16, J <sub>AB</sub> = 164Hz), 4.49 (d, 2H, J = 4.0Hz), 4.27 (t, 1H, J = 8.0Hz), 4.04 (m, 4H), 3.69 (m, 2H), 3.51 (m, 2H), 3.10 (m, 2H), 1.83 (m, 1H), 1.29 (m, 1H), 1.07 (m, 1H), 0.75 (d, 3H, J = 8.0Hz), 0.68 (d, 3H, J = 8.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
428				11	yellow solid	555.12	1.41 min Method E	555.28	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 7.71 (d, 2H, J = 8.00Hz), 7.50 (d, 2H, J = 8.0Hz), 7.16 (d, 1H, J = 12.0Hz), 7.05 (d, 1H, J = 8.0Hz), 6.87 (t, 1H, J = 8.0Hz), 6.38 (s, br, 1H), 5.91 (s br, 1H), 4.41 (ABq, 2H, Δν = 16, J <sub>AB</sub> = 176Hz), 4.45 (m, 2H), 4.27 (t, 1H, J = 8.0Hz), 3.81 (m, 4H), 3.67 (m, 4H), 3.48 (m, 1H), 2.89 (s, 3H), 1.83 (m, 1H), 1.29 (m, 1H), 1.05 (m, 1H), 0.75 (d, 3H, J = 8.0Hz), 0.68 (d, 3H, J = 8.0Hz).
429				7	white solid	545.17	1.84 min Method C	545.36	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J = 9.0), 7.51 (d, 2H, J = 8.0), 6.65 (2 s, 1H), 5.40 (s, 1H), 4.54 (t, 1H, J = 13), 3.85-4.20 (m, 2H), 2.40-3.50 (m, 12H), 1.75-2.00 (m, 3H), 0.92-1.20 (m, 1H), 0.73 (d, 3H, J = 5.8, 6.1), 0.67 (d, 3H, J = 6.1, 6.4).
430				7	white solid	541.16	1.43 min Method A	541.24	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.71 (d, 2H, J = 6.4), 7.50 (d, 2H, J = 8.2), 6.65 (d, 1H, J = 41), 5.44 (s, 1H), 4.55 (t, 1H, J = 13), 4.15 (br s, 1H), 3.95 (br s, 1H), 3.15-3.35 (m, 1H), 2.90-3.00 (m, 2H), 2.60-2.70 (m, 2H), 2.50-2.57 (m, 2H), 2.35-2.50 (m, 6H), 1.85-2.00 (m, 3H), 1.35-1.75 (m, 3H), 1.00-1.30 (m, 5H), 0.90-1.00 (m, 1H), 0.72 (dd, 3H, J = 7.0, 7.3), 0.66 (dd, 3H, J = 6.1, 6.4).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
431				7	white solid	529.10	1.38 min Method A	529.27	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.5), 7.51 (d, 2H, J = 8.9), 6.65 (d, 1H, J = 32), 5.43 (s, 1H), 4.55 (t, 1H, J = 13), 4.14 (dd, 1H, J = 4.8, 9.5), 3.88 (br s, 4H), 3.21-3.84 (m, 2H), 2.90-3.05 (m, 3H), 2.47-2.67 (m, 1H), 1.80-2.05 (m, 3H), 1.40-1.75 (m, 6H), 1.00-1.35 (m, 4H), 0.73 (dd, 3H, J = 3.4, 6.4), 0.67 (dd, 3H, J = 2.7, 6.7).
432				7	white solid	474.02	1.55 min Method A	474.28	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 7.6), 7.50 (d, 2H, J = 8.6), 6.65 (d, 1H, J = 40), 5.46 (s, 1H), 4.56 (t, 1H, J = 13), 4.00-4.20 (m, 2H), 3.80-3.90 (m, 1H), 3.40 (s, 3H), 3.20-3.35 (m, 1H), 2.85-3.05 (m, 2H), 2.40-2.65 (m, 1H), 1.50-2.00 (m, 5H), 1.00-1.45 (m, 2H), 0.83-0.90 (m, 1H), 0.72 (dd, 3H, J = 6.7, 8.2), 0.67 (dd, 3H, J = 5.8, 6.4).
433				1-Method A	white solid	455.00	1.83 min Method B	454.15	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.61 (dd, 2H, J = 1.8, 8.7), 7.40 (ddd, 2H, J = 2.1, 2.4, 8.7), 7.26 (d, 4H, J = 7.2), 6.25 (br s, 1H), 5.19 (br s, 1H), 4.51 (d, 1H, J = 15.6), 4.43 (d, 1H, J = 15.6), 4.34 (t, 1H, J = 7.2), 1.75-1.85 (m, 1H), 1.69 (s, 3H), 1.62 (s, 3H), 1.22-1.35 (m, 2H), 0.78 (d, 3H, J = 6.3), 0.66 (d, 3H, J = 6.3).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
434				7	white solid	542.15	1.73 min Method C	542.46	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.72 (d, 2H, J = 8.5), 7.51 (d, 2H, J = 8.5), 6.65 (d, 1H, J = 39), 5.42 (s, 1H), 4.55 (t, 1H, J = 14), 4.00-4.17 (m, 2H), 3.05-3.33 (m, 3H), 2.85-3.05 (m, 2H), 2.40-2.70 (m, 8H), 2.32 (s, 3H), 1.55-2.10 (m, 6H), 1.00-1.30 (m, 6H), 0.72 (t, 3H, J = 6.7, 6.7), 0.67 (t, 3H, J = 6.1, 6.4).
435				7	white solid	497.02	1.70 min Method A	479.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 8.31 (s, 1H), 7.76 (d, 2H, J = 8.9), 7.56 (d, 2H, J = 8.5), 6.74 (s, 1H), 4.81 (t, 1H, J = 7.6), 4.71 (br s, 1H), 4.20 (br s, 1H), 3.13 (t, 2H, J = 8.6), 2.70-2.95 (m, 2H), 2.05-2.20 (m, 1H), 1.85-2.00 (m, 2H), 1.55-1.85 (m, 4H), 1.10-1.35 (m, 3H), 1.00 (d, 3H, J = 6.4), 0.97 (d, 3H, J = 6.7), 0.87 (t, 1H, J = 7.0).
436				13	white solid	476.99	1.76 min Method B	477.22	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.94 (dd, 2H, J = 1.8, 8.4), 7.69 (dd, 2H, J = 1.8, 8.7), 7.45-7.50 (m, 4H), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.65 (d, 1H, J = 15.9), 4.46 (d, 1H, J = 15.9), 4.31 (dd, 1H, J = 6.6, 7.8), 2.61 (s, 3H), 1.75-1.85 (m, 1H), 1.28-1.35 (m, 1H), 1.08-1.15 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.64 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
437				14	pale yellow solid	476.99	1.92 min Method B	477.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.04 (d, 2H, J = 8.4), 7.70 (dd, 2H, J = 1.8, 8.4), 7.45-7.52 (m, 4H), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.67 (d, 1H, J = 16.2), 4.47 (d, 1H, J = 15.9), 4.31 (t, 1H, J = 7.2), 2.47 (s, 3H), 1.75-1.85 (m, 1H), 1.28-1.35 (m, 1H), 1.08-1.15 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.64 (d, 3H, J = 6.6).
438				6	white solid	480.03	1.81 min Method B	480.26	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.67 (d, 2H, J = 8.7), 7.45 (d, 2H, J = 8.7), 7.38 (d, 2H, J = 8.1), 7.32 (d, 2H, J = 7.5), 6.23 (br s, 1H), 5.19 (br s, 1H), 4.59 (d, 1H, J = 16.2), 4.45 (d, 1H, J = 15.9), 4.30 (t, 1H, J = 6.9), 3.47-3.56 (br m, 1H), 3.15-3.35 (br m, 1H), 2.81-3.09 (br m, 3H), 1.75-1.85 (m, 1H), 1.05-1.40 (m, 5H), 0.76 (d, 3H, J = 6.6), 0.65 (d, 3H, J = 6.6).
439				6	white solid	529.06	1.60 min Method B	529.25	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 8.57 (d, 1H, J = 4.8), 7.80 (d, 2H, J = 8.4), 7.70-7.73 (m, 2H), 7.67 (d, 2H, J = 8.4), 7.45 (d, 2H, J = 8.4), 7.42 (d, 2H, J = 7.8), 7.36 (d, 1H, J = 7.8), 7.28 (br s, 1H), 6.23 (br s, 1H), 5.22 (br s, 1H), 4.77 (d, 2H, J = 4.8), 4.63 (d, 1H, J = 15.9), 4.44 (d, 1H, J = 15.9), 4.29 (t, 1H, J = 7.2), 1.73-1.85 (m, 1H), 1.25-1.38 (m, 1H), 1.06-1.14 (m, 1H), 0.75 (d, 3H, J = 6.3), 0.65 (d, 3H, J = 6.6).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
440				8	amber glass	512.04	1.36 min Method A	512.24	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.70 (d, 2H, J=8.0Hz), 7.74 (d, 2H, J=8.0Hz), 7.31 (d, 2H, J=6Hz), 7.08 (d, 2H, J=8.0Hz), 6.25 (s, br, 1H), 5.41 (s, br, 1H), 4.56 (d, 1H, J <sub>AB</sub> =12Hz), 4.43 (d, 1H, J <sub>AB</sub> =12Hz), 4.35 (t, 1H, J=6.0Hz), 3.72 (t, 4H, J=4.0Hz), 3.56 (s, 2H), 2.48 (t, 4H, J=41.0Hz), 1.79 (m, 1H), 1.36 (m, 1H), 1.18 (m, 1H), 0.79 (d, 3H, J=6.0Hz), 0.69 (d, 3H, J=6.0Hz).
441				8	amber glass	525.08	1.35 min Method A	525.23	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.69 (d, 2H, J=8.0Hz), 7.47 (d, 2H, J=8.0Hz), 7.28 (t, 1H, J=6.0Hz), 7.07 (dd, 2H, J=8.0Hz), 6.29 (s, br, 1H), 5.55 (s, br, 1H), 4.46 (d, 1H, J <sub>AB</sub> =14.0Hz), 4.41 (d, 1H, J <sub>AB</sub> =14.0Hz), 4.34 (t, 1H, J=6.0Hz), 3.58 (s, 2H), 2.33 (s, 3H), 1.78 (m, 1H), 1.18 (m, 1H), 0.79 (d, 3H, J=6.0Hz), 0.68 (d, 3H, J=6.0Hz).
442				1-Method A	amber glass	470.94	1.74 min Method A	471.12	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.78 (t, 1H, J=6.0Hz), 7.72 (d, 2H, J=8.0Hz), 7.51 (d, 2H, J=8.0Hz), 7.19 (m, 2H), 6.21 (s, br, 1H), 5.37 (s, br, 1H), 4.64 (d, 1H, J <sub>AB</sub> =14.0Hz), 4.47 (d, 1H, J <sub>AB</sub> =14.0Hz), 4.34 (t, 1H, J=6.0Hz), 3.95 (s, 3H), 1.80 (m, 1H), 1.37 (m, 1H), 1.01 (m, 1H), 0.80 (d, 3H, J=6.0Hz), 0.69 (d, 3H, J=6.0Hz).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
443				21	white solid	435.10	1.40 min Method D	(M+Na) <sup>+</sup> 458.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.84 (d, 2H, J=8.6), 7.76 (d, 2H, J=8.3), 7.62 (d, 2H, J=8.8), 7.51 (d, 2H, J=8.3), 7.40 (s, 1H), 7.11 (s, 1H), 4.63 (ABq, 2H, Δν=5.9, J <sub>A</sub> =17.6), 4.56 (dd, 1H, J=8.3, 2.5), 4.54 (s, 1H), 1.95 (dd, 1H, J=13.7, 8.6), 1.26 (dd, 1H, J=13.6, 2.4), 1.04 (s, 3H), 0.99 (s, 3H).
444				7	clear oil	465.17	1.38 min Method A	466.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.73 (d, 2H, J=8.8), 7.50 (d, 2H, J=8.9), 6.67 (s, 1H), 5.37 (s, 1H), 4.14 (dd, 1H, J=5.5, 9.5), 3.25 (dd, 1H, J=10, 14), 2.97 (dd, 1H, J=4.5, 14), 2.87-2.95 (m, 2H), 2.65-2.75 (m, 2H), 2.07-2.23 (m, 2H), 1.83-1.90 (m, 1H), 1.50-1.82 (m, 4H), 1.15-1.40 (m, 3H), 0.77-0.85 (m, 1H), 0.72 (d, 3H, J=6.7), 0.66 (d, 3H, J=6.4).
445				6	amber glass	456.92	1.62 min Method A	457.32	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.77 (d, 2H, J=6.0Hz), 7.68 (t, 1H, J=6.0Hz), 7.64 (d, 2H, J=8.0Hz), 7.39 (m, 3H), 6.21 (s, br, 1H), 5.35 (s, br, 1H), 4.67 (d, 1H, J <sub>A</sub> =14.0Hz), 4.38 (d, 1H, J <sub>A</sub> =14.0Hz), 3.44 (m, 1H), 1.88 (m, 1H), 1.59 (m, 2H), 0.79 (d, 3H, J=6.0Hz), 0.69 (d, 3H, J=6.0Hz)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
446				18	off-white solid	417.09	1.53 min Method F	(M+Na) <sup>+</sup> 418.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 7.82 (d, 2H, J=8.8), 7.74 (d, 2H, J=8.5), 7.63 (d, 2H, J=8.8), 7.62 (s, 1H), 7.53 (d, 2H, J=8.5), 7.09 (s, 1H), 4.80 (ABq, 2H, Δν=17.9, J <sub>A</sub> =17.8), 4.70 (s, 1H), 4.62 (t, 1H, J=7.6), 4.60 (s, 1H), 2.34 (dd, 1H, J=14.4, 7.1), 2.02 (dd, 1H, J=14.6, 7.3), 1.57 (s, 3H).
447				18	off-white solid	458.12	1.59 min Method G	(M+Na) <sup>+</sup> 459.2	<sup>1</sup> H NMR (400 MHz, DMSO) δ 8.45 (d, 1H, J=2.2), .82 (d, 2H, J=8.6), 7.72 (m, 3H), 7.60 (d, 2H, J=8.8), 7.57 (s, 1H), 7.44 (d, 2H, J=8.5), 7.09 (s, 1H), 6.54 (t, 1H, J=2.0), 4.74 (ABq, 2H, Δν=25.5, J <sub>A</sub> =16.8), 4.71 (s, 1H), 4.61 (m, 2H), 2.37 (dd, 1H, J=14.2, 7.1), 2.08 (dd, 1H, J=14.7, 7.6), 1.57 (s, 3H).
448				18	off-white solid	442.90	1.74 min Method A	443.05	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.96 (d, 2H, J=8.2), 7.72 (d, 2H, J=8.0), 7.48 (d, 2H, J=8.0), 7.39 (d, 2H, J=8.0), 6.33 (br s, 1H), 5.20 (br s, 1H), 4.55-4.62 (m, 2H), 4.39 (d, 1H, J=15.4), 4.28-4.32 (m, 1H), 4.17-4.21 (m, 1H), 3.90 (s, 3H), 2.17-2.35 (m, 1H), 1.56-1.65 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
449				18, 12	yellow foam	442.94	1.65 min Method A	442.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.67 (ddd, 2H, J = 2.0, 2.6, 8.8), 7.42 (ddd, 2H, J = 2.0, 2.3, 8.8), 7.39 (d, 2H, J = 8.2), 7.24 (d, 2H, J = 8.2), 6.34 (br s, 1H), 5.35 (br s, 1H), 4.56 (dd, 1H, J = 5.8, 8.5), 4.48 (d, 1H, J = 15.5), 4.35 (d, 1H, J = 15.5), 4.30-4.33 (m, 1H), 4.18-4.22 (m, 1H), 2.23-2.39 (m, 1H), 1.56-1.670 (m, 2H), 1.57 (s, 6H).
450				18, 13	white solid	466.92	1.61 min Method A	467.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.96 (ddd, 2H, J = 1.7, 2.0, 8.4), 7.73 (ddd, 2H, J = 1.9, 2.5, 8.7), 7.49 (ddd, 2H, J = 2.0, 2.3, 8.7), 7.46 (d, 2H, J = 8.6), 6.34 (br s, 1H), 5.21 (br s, 1H), 4.64 (d, 1H, J = 15.4), 4.57-4.60 (m, 1H), 4.39 (d, 1H, J = 16.1), 4.30-4.32 (m, 1H), 4.18-4.21 (m, 1H), 2.61 (s, 3H), 2.18-2.36 (m, 1H), 1.55-1.66 (m, 1H).
451				18, 7	clear oil	533.07	1.71 min Method A	533.22	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.73 (d, 2H, J = 8.6), 7.51 (d, 2H, J = 8.8), 6.67 (s, 1H), 5.51 (s, 1H), 4.14-4.52 (m, 4H), 3.76-3.95 (m, 2H), 3.50-3.72 (m, 1H), 3.19-3.27 (m, 1H), 2.86-3.07 (m, 1H), 2.56-2.80 (m, 2H), 1.70-1.99 (m, 7H), 1.36-1.63 (m, 10H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
452				18, 7	clear oil	517.07	2.53 min Method A	517.32	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.74 (d, 2H, J = 8.6), 7.50 (d, 2H, J = 8.5), 6.70 (s, 1H), 5.60 (s, 1H), 4.50 (dd, 1H, J = 4.9, 9.8), 4.22-4.31 (m, 2H), 3.81-4.00 (m, 5H), 3.53-3.66 (m, 2H), 3.40-3.50 (m, 1H), 3.13-3.25 (m, 3H), 2.89 (dd, 1H, J = 4.6, 14), 2.62-2.77 (m, 3H), 1.50-1.95 (m, 6H), 1.01-1.45 (m, 5H).
453				18, 7	clear oil	531.09	2.27 min Method A	531.36	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.73 (d, 2H, J = 8.6), 7.51 (d, 2H, J = 8.6), 6.65 (d, 1H, J = 28), 5.47 (s, 1H), 4.50-4.63 (m, 1H), 4.20-4.45 (m, 1H), 4.05-4.15 (m, 2H), 3.55-3.65 (m, 2H), 3.05-3.35 (m, 6H), 2.85-3.00 (m, 3H), 2.55-2.35 (m, 2H), 1.50-2.00 (m, 6H), 1.00-1.45 (m, 7H).
454				8	amber glass	533.99	1.40 min Method A	534.27	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.78 (d, 2H, J = 8.0Hz), 7.39 (m, 6H), 6.34 (s, br, 1H), 5.80 (s, br, 1H), 4.72 (d, 1H, J <sub>AB</sub> = 14.0Hz), 4.16 (m, 1H), 4.34 (d, 1H, J <sub>AB</sub> = 14.0Hz), 3.60 (m, 4H), 3.84 (s, 2H), 2.62 (m, 2H), 2.25 (m, 4H).
455				18, 8	amber glass	501.98	1.27 min Method A	502.24	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.79 (d, 2H, J = 8.0Hz), 7.39 (m, 6H), 6.33 (s, br, 1H), 5.80 (s, br, 1H), 4.68 (d, 1H, J <sub>AB</sub> = 14.0Hz), 4.22 (d, 1H, J <sub>AB</sub> = 14.0Hz), 3.55 (s, 2H), 2.62 (m, 2H), 2.43 (m, 8H), 2.30 (s, 3H).



Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
456				18, 8	amber glass	515.02	1.22 min Method A	515.31	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.79 (d, 2H, J=8.0Hz), 7.52 (d, 2H, J=8.0Hz), 7.39 (m, 4H), 6.34 (s, br, 1H), 5.78 (s, br, 1H), 5.72 (t, 1H, J=54.0Hz), 4.68 (d, 1H, J <sub>ab</sub> =14.0Hz), 4.34 (d, 1H, J <sub>ab</sub> =14.0Hz), 3.88 (m, 1H), 3.56 (m, 4H), 2.49 (m, 2H), 2.26 (m, 4H)
457				8	amber glass	547.03	1.26 min Method A	547.20	<sup>1</sup> H NMR 400Hz (CDCl <sub>3</sub> ) δ 7.78 (d, 2H, J=8.0Hz), 7.50 (d, 2H, J=8.0Hz), 7.41 (m, 4H), 6.35 (s, br, 1H), 5.80 (s, br, 1H), 5.72 (t, 1H, J=54.0Hz), 4.68 (d, 1H, J <sub>ab</sub> =14.0Hz), 4.35 (d, 1H, J <sub>ab</sub> =14.0Hz), 3.87 (m, 1H), 3.55 (s, 2H), 2.44 (m, 10H), 2.30 (s, 3H)
458				18, 1-Method B	white foam	446.91	1.01 Method B	447.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt δ 8.14 (s, 1H), 7.98 (d, 1H, J=9.6Hz), 7.76 (d, 2H, J=6.8Hz), 7.55 (d, 2H, J=6.8Hz), 6.81 (d, 2H, J=9.6Hz), 6.45 (s, 1H), 6.10 (s, 1H), 5.70 (t, 1H, J=110.0Hz), 4.60 (d, 1H, J=16.0Hz), 4.51 (m, 1H), 4.06 (d, 1H, J=16Hz), 3.30 (s, 6H), 2.55 (m, 1H), 1.60 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
459				18, 1-Method B	white foam	456.97	1.14 Method B	457.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) TFA salt δ 8.11 (s, 1H), 7.95 (d, 1H, J=9.6Hz), 7.77 (d, 2H, J=6.8Hz), 7.51 (d, 2H, J=6.8Hz), 6.76 (d, 2H, J=9.6Hz), 6.34 (s, 1H), 6.02 (s, 1H), 4.58 (d br., 1H, J=8.4Hz), 4.46 (d, 1H, J=16.0Hz), 4.06 (d, 1H, J=16Hz), 3.29 (s, 6H), 2.50 (m, 1H), 1.39 (m, 1H), 1.25 (d, 3H, J=22.0Hz), 1.17 (d, 3H, J=22.0Hz).
460				1-Method B	light yellow gummy solid	478.92	1.89 min 3 X 50 mm ODS-A C-18 column, 4mL/min, 0-100% MeOH/H <sub>2</sub> O 0.1 %TFA 4min gradient	479.21	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.09 (s, 1H), 8.01 (d, 1H, J=9.2Hz), 7.75 (d, 2H, J=8.4Hz), 7.53 (d, 2H, J=8.4Hz), 6.82 (d, 1H, J=9.2Hz), 6.62 (s, br, 1H), 6.12 (s, br, 1H), 4.18-4.58 (m, 3H), 3.30 (s, 6H), 2.15 (m, 1H), 2.05 (m, 1H), 1.85 (m, 1H), 1.40 (m, 1H)
461				18	clear oil	438.28	1.41 min Method B	438.01	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.33 (s, 1H), 7.73 (d, 1H, J=8.4Hz), 7.71 (d, 2H, J=8.8Hz), 7.51 (d, 2H, J=8.8Hz), 7.27 (d, 1H, J=8.4Hz), 6.56 (s, br, 1H), 6.11 (s, br, 1H), 5.69 (m, 1H), 4.21-4.62 (m, 3H), 2.52 (m, 1H), 1.63 (m, 1H)

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
462				1	white solid	470.30	1.53 min Method B	470.02	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.38 (s, 1H), 7.79 (d, 1H, J=8.0Hz), 7.68 (d, 2H, J=8.4Hz), 7.50 (d, 2H, J=8.4Hz), 7.31 (d, 1H, J=8.0Hz), 6.57 (s, br, 1H), 6.25 (s, br, 1H), 4.29-4.64 (m, 3H), 2.12 (m, 1H), 1.98 (m, 1H), 1.81 (m, 1H), 1.43 (m, 1H)
463				18	clear oil	434.32	1.43 min Method B	434.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.37 (s, 1H), 7.70 (d, 1H, J=8.8Hz), 7.68 (d, 1H, J=8.8Hz), 7.67 (d, 2H, J=6.8Hz), 7.48 (d, 2H, J=6.8Hz), 6.25 (s, br, 1H), 5.31 (s, br, 1H), 4.34-4.62 (m, 5H), 1.35-2.05 (m, 4H)
464				18, 7	clear oil	469.14	1.14 min Method A	470.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.73 (d, 2H, J=8.8), 7.51 (d, 2H, J=8.8), 6.65 (s, 1H), 5.39 (s, 1H), 4.05-4.35 (m, 2H), 3.25 (dd, 1H, J=10, 14), 2.85-3.04 (m, 3H), 2.65-2.85 (m, 2H), 2.09-2.29 (m, 2H), 1.93-2.10 (m, 1H), 1.83-1.91 (m, 10H)
465				15	white solid	476.99	1.91 min Method A	477.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.98 (d, 2H, J=8.2), 7.68 (d, 2H, J=8.9), 7.45 (d, 4H, J=8.5), 6.21 (s, 1H), 5.19 (s, 1H), 4.62 (d, 1H, J=15), 4.48 (d, 1H, J=16), 4.31 (t, 1H, J=7.0), 2.65 (s, 3H), 1.75-1.85 (m, 1H), 1.20-1.35 (m, 4H), 1.10-1.17 (m, 1H), 0.85-0.90 (m, 1H), 0.75 (d, 3H, J=6.7), 0.64 (d, 3H, J=6.4)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
466				1-Method B	clear gummy solid	520.96	1.17 min Method B	521.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.16 (s, 1H), 8.06 (d, 1H, J=9.2Hz), 7.75 (d, 2H, J=8.4Hz), 7.50 (d, 2H, J=8.4Hz), 7.31 (d, 1H, J=9.2Hz), 6.67 (s, br, 1H), 6.20 (s, br, 1H), 4.19-4.60 (m, 3H), 3.87 (m, 4H), 3.69 (m, 4H), 2.14 (m, 1H), 1.98 (m, 1H), 1.83 (m, 1H), 1.38 (m, 1H)
467				18, 1-Method B	clear gummy solid	470.95	0.983 min Method B	471.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.15 (s, 1H), 8.10 (d, 1H, J=9.2Hz), 7.77 (d, 2H, J=8.4Hz), 7.53 (d, 2H, J=8.4Hz), 6.93 (d, 1H, J=9.2Hz), 6.65 (s, br, 1H), 6.22 (s, br, 1H), 4.09-4.67 (m, 5H), 3.87 (m, 4H), 3.68 (m, 4H), 2.25 (m, 1H), 1.63 (m, 1H)
468				1-Method B	clear gummy solid	499.01	1.153 min Method B	499.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> ) δ 8.12 (s, 1H), 8.07 (d, 1H, J=9.2Hz), 7.78 (d, 2H, J=8.4Hz), 7.52 (d, 2H, J=8.4Hz), 6.91 (d, 1H, J=9.2Hz), 6.63 (s, br, 1H), 6.19 (s, br, 1H), 4.07-4.62 (m, 3H), 3.86 (m, 4H), 3.68 (m, 4H), 2.46 (m, 1H), 1.31 (m, 1H), 1.25 (d, 3H, J=21.6), 1.17 (d, 3H, J=21.6)
469				15	white solid	511.43	1.90 min Method C	511.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.01 (d, 2H, J=8.3), 7.69 (d, 2H, J=8.8), 7.43-7.51 (m, 4H), 6.20 (s, 1H), 5.15 (s, 1H), 4.74 (s, 2H), 4.63 (d, 1H, J=15), 4.47 (d, 1H, J=16), 4.31 (t, 1H, J=6.8), 1.74-1.87 (m, 1H), 1.04-1.88 (m, 4H), 0.82-0.94 (m, 1H), 0.76 (d, 3H, J=6.6), 0.65 (d, 3H, J=6.6)










Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
470				1	white solid	529.11	2.17 min Method D	529.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.95 (d, 2H, J = 8.4), 7.71 (dd, 2H, J = 1.6, 8.4), 7.48 (dd, 2H, J = 2.4, 8.8), 7.38 (dd, 2H, J = 2.4, 8.8), 6.31 (br s, 1H), 5.22 (br s, 1H), 4.60 (d, 1H, J = 15.6), 4.55-4.58 (m, 1H), 4.38-4.42 (m, 3H), 4.29-4.32 (m, 1H), 4.17-4.21 (m, 1H), 2.18-2.32 (m, 2H), 1.12 (dd, 2H, J = 6.8, 8.4), 0.08 (s, 9H).
471				15, 23	white solid	510.14	1.86 min Method A	511.13	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.01 (d, 2H, J = 8.4), 7.69 (d, 2H, J = 8.8), 7.47 (d, 2H, J = 8.0), 7.46 (d, 2H, J = 8.8), 6.20 (br s, 1H), 5.20 (br s, 1H), 4.74 (s, 2H), 4.64 (d, 1H, J = 15), 4.49 (d, 1H, J = 15), 4.31 (t, 1H, J = 7.0), 1.73-1.87 (m, 1H), 1.20-1.37 (m, 1H), 1.07-1.17 (m, 1H), 0.76 (d, 3H, J = 6.6), 0.65 (d, 3H, J = 6.6).
472				15, 8	white solid	519.17	1.42 min Method A	520.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.02 (d, 2H, J = 7.6), 7.71 (d, 2H, J = 8.4), 7.52 (d, 2H, J = 8.0), 7.48 (d, 2H, J = 8.4), 6.23 (br s, 1H), 5.17 (br s, 1H), 4.70 (d, 1H, J = 16), 4.44 (d, 1H, J = 16), 4.31 (t, 1H, J = 6.4), 3.70 (s, 2H), 3.02 (s, 6H), 1.75-1.90 (m, 1H), 1.00-1.40 (m, 2H), 0.76 (d, 3H, J = 6.8), 0.66 (d, 3H, J = 6.4).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
473				1	off-white foam	460.89	2.11 min Method A	461.08	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.97 (d, 2H, J = 8.4), 7.72 (d, 2H, J = 8.4), 7.51 (dd, 2H, J = 2.0, 8.4), 7.38 (d, 2H, J = 8.0), 6.32 (br s, 1H), 5.71 (tm, 1H, J <sub>H-F</sub> = 55), 5.19 (br s, 1H), 4.60 (d, 1H, J = 15.6), 4.49-4.53 (m, 1H), 4.33 (d, 1H, J = 15.6), 3.91 (s, 3H), 2.42-2.68 (m, 1H), 1.54-1.65 (m, 1H).
474				1	pale yellow foam	462.92	2.24 min Method A	485.07 (M + Na <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.67 (d, 2H, J = 8.4), 7.47 (d, 2H, J = 8.8), 7.25-7.29 (m, 4H), 6.33 (br s, 1H), 5.72 (tm, 1H, J <sub>H-F</sub> = 57), 5.23 (br s, 1H), 4.49-4.53 (m, 1H), 4.46 (d, 1H, J = 15.6), 4.34 (d, 1H, J = 15.0), 2.52-2.60 (m, 1H), 1.55-1.65 (m, 1H), 1.68 (s, 3H), 1.63 (s, 3H).
475				1, 13	white foam	484.91	1.67 min Method D	485.07	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.97 (d, 2H, J = 8.0), 7.73 (dd, 2H, J = 2.0, 8.8), 7.52 (dd, 2H, J = 2.0, 8.8), 7.45 (d, 2H, J = 8.4), 6.34 (br s, 1H), 5.73 (tm, 1H, J <sub>H-F</sub> = 57), 5.20 (br s, 1H), 4.63 (d, 1H, J = 15.6), 4.52-4.55 (m, 1H), 4.32 (d, 1H, J = 15.6), 2.61 (s, 3H), 2.55-2.60 (m, 1H), 1.61-1.65 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
476				1, 12	white foam	460.93	1.72 min Method D	483.07 (M + Na <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.68 (dd, 2H, J = 2.0, 8.8), 7.48 (dd, 2H, J = 8.4), 7.41 (d, 2H, J = 8.4), 7.23 (d, 2H, J = 8.8), 6.30 (br s, 1H), 5.73 (tm, 1H, J <sub>HF</sub> = 57), 5.18 (br s, 1H), 4.48-4.52 (m, 1H), 4.44 (d, 1H, J = 15.2), 4.34 (d, 1H, J = 15.2), 2.50-2.65 (m, 1H), 1.61-1.70 (m, 1H), 1.56 (s, 6H).
477				7	clear oil	455.93	0.99 min Method A	456.20	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72-7.86 (m, 2H), 7.45-7.52 (m, 2H), 6.70 (br s, 1H), 5.44 (br s, 1H), 4.46-4.57 (m, 2H), 4.00-4.37 (m, 2H), 3.24 (dd, 1H, J = 4.4, 9.8), 2.70-3.15 (m, 6H), 2.15-2.34 (m, 2H), 1.80-1.93 (m, 2H), 1.54-1.63 (m, 2H), 1.25-1.35 (m, 2H).
478				7	clear oil	519.04	1.12 min Method A	519.23	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.71-7.84 (m, 2H), 7.39-7.55 (m, 2H), 6.70 (br s, 1H), 5.87 (br s, 1H), 4.88 (m, 1H), 4.50 (d, 1H, J = 9.5), 4.21-4.31 (m, 2H), 4.11-4.20 (m, 2H), 3.80-3.87 (m, 2H), 3.69-3.77 (m, 2H), 3.31-3.50 (m, 2H), 3.00-3.21 (m, 2H), 2.80-3.95 (m, 2H), 2.10-2.50 (m, 4H), 1.85-1.95 (m, 2H), 1.73 (d, 2H, J = 8.5), 1.40-1.50 (m, 2H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
479				7	clear oil	535.06	1.72 min Method A	535.18	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.73 (d, 2H, J = 8.6), 7.53 (d, 2H, J = 8.6), 6.70 (br s, 1H), 5.70 (tm, 1H, J <sub>HF</sub> = 50), 5.55 (br s, 1H), 4.46 (dd, 1H, J = 3.8, 10), 3.80-4.00 (m, 3H), 3.50-3.70 (m, 1H), 3.15-3.30 (m, 1H), 2.80-3.95 (m, 1H), 2.60-2.80 (m, 2H), 2.40-2.60 (m, 1H), 1.05-2.00 (m, 16H).
480				1	off-white foam	444.93	1.89 min Method D	467.09 (M + Na <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.66 (ddd, 2H, J = 2.0, 2.4, 8.8), 7.44 (ddd, 2H, J = 2.0, 2.4, 8.8), 7.25-7.31 (m, 4H), 6.30 (br s, 1H), 5.22 (br s, 1H), 4.56-4.60 (m, 1H), 4.47 (d, 1H, J = 15.6), 4.39 (d, 1H, J = 16.0), 4.32-4.36 (m, 1H), 4.20-4.24 (m, 1H), 2.23-2.39 (m, 1H), 1.70-1.85 (m, 1H), 1.65-1.68 (m, 3H), 1.53 (s, 3H).
481				6	white solid	428.87	1.58 min Method D	429.04	<sup>1</sup> H NMR (CD <sub>3</sub> OD, 400MHz) δ 7.92 (d, 2H, J = 8.4), 7.79 (ddd, 2H, J = 2.0, 2.4, 8.8), 7.52 (ddd, 2H, J = 2.0, 2.8, 8.8), 7.47 (d, 2H, J = 8.0), 4.80 (d, 1H, J = 16), 4.66 (d, 1H, J = 14.5), 4.64 (t, 1H, J = 7.6), 4.10-4.33 (m, 2H), 2.05-2.12 (m, 1H), 1.72-1.80 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
482				6	yellow solid	446.86	1.39 min Method B	447.06	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.89 (d, 2H, J = 8.0), 7.66 (dd, 2H, J = 2.0, 8.0), 7.41 (dd, 2H, J = 2.0, 8.4), 7.31 (d, 2H, J = 8.0), 6.61 (br s, 1H), 5.90 (br s, 1H), 5.76 (tm, 1H, J <sub>H-F</sub> = 57), 4.56 (d, 1H, J = 16.0), 4.49-4.52 (m, 1H), 4.36 (d, 1H, J = 12.0), 2.50-2.65 (m, 1H), 1.61-1.70 (m, 1H).
483				6	white solid	455.94	1.34 min Method B	488.11 (M + Na <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (d, 2H, J = 8.0), 7.68 (d, 2H, J = 8.0), 7.49 (ddd, 2H, J = 2.0, 2.4, 8.4), 7.38 (d, 2H, J = 8.0), 6.32 (br s, 1H), 6.06 (br s, 1H), 5.19 (br s, 1H), 4.59 (d, 1H, J = 15.0), 4.56-4.60 (m, 1H), 4.37 (d, 1H, J = 15.0), 4.29-4.32 (m, 1H), 4.16-4.19 (m, 1H), 3.45-3.49 (m, 2H), 2.15-2.30 (m, 1H), 1.50-1.65 (m, 1H), 1.25 (t, 3H, J = 8.0).
484				6	white solid	469.97	1.40 min Method B	470.15	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (d, 2H, J = 8.0), 7.48 (d, 2H, J = 8.0), 7.30-7.36 (m, 4H), 6.30 (br s, 1H), 5.26 (br s, 1H), 4.54-4.58 (m, 2H), 4.38 (d, 1H, J = 15.2), 4.29-4.32 (m, 1H), 4.18-4.22 (m, 1H), 3.55-3.59 (m, 1H), 3.22-3.27 (m, 1H), 2.90 and 3.05 (2 s, 3H), 2.20-2.33 (m, 1H), 1.50-1.65 (m, 1H), 1.08-1.30 (m, 3H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
485				6	white solid	473.93	1.33 min Method B	474.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (d, 2H, J = 8.8), 7.70 (d, 2H, J = 8.4), 7.51 (d, 2H, J = 8.8), 7.38 (d, 2H, J = 8.0), 6.31 (br s, 1H), 6.05 (br s, 1H), 5.72 (tm, 1H, J <sub>H-F</sub> = 57), 5.20 (br s, 1H), 4.59 (d, 1H, J = 15.6), 4.49-4.52 (m, 1H), 4.31 (d, 1H, J = 15.6), 3.46-3.53 (m, 2H), 2.45-2.65 (m, 1H), 1.57-1.65 (m, 1H), 1.25 (t, 3H, J = 7.2).
486				6	white solid	536.99	1.13 min Method B	537.17	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.57 (d, 1H, J = 4.8), 7.83 (d, 2H, J = 8.4), 7.71-7.78 (m, 3H), 7.71 (br s, 1H), 7.52 (dd, 2H, J = 1.6, 8.8), 7.40 (d, 2H, J = 8.0), 7.30-7.35 (m, 1H), 7.23-7.28 (m, 1H), 6.32 (br s, 1H), 5.72 (tm, 1H, J <sub>H-F</sub> = 57), 5.25 (br s, 1H), 4.75 (d, 2H, J = 4.8), 4.59 (d, 1H, J = 15.6), 4.49-4.53 (m, 1H), 4.33 (d, 1H, J = 15.6), 2.48-2.65 (m, 1H), 1.58-1.65 (m, 1H).
487				6	white solid	519.00	1.10 min Method B	519.35	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.58 (d, 1H, J = 4.4), 7.92 (d, 2H, J = 8.4), 7.71-7.78 (m, 4H), 7.48 (dd, 2H, J = 2.0, 8.8), 7.39-7.44 (m, 3H), 7.24-7.30 (m, 1H), 6.28 (br s, 1H), 5.20 (br s, 1H), 4.78 (d, 2H, J = 4.8), 4.55-4.61 (m, 2H), 4.39 (d, 1H, J = 15.6), 4.29-4.32 (m, 1H), 4.17-4.21 (m, 1H), 2.20-2.33 (m, 1H), 1.50-1.65 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
488				6	white solid	487.96	1.63 min Method D	488.34	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (dd, 2H, J = 2.0, 8.8), 7.51 (dd, 2H, J = 2.0, 8.4), 7.50 (s, 4H), 6.30 (br s, 1H), 5.72 (tm, 1H, J <sub>HF</sub> = 57), 5.24 (br s, 1H), 4.55 (d, 1H, J = 15.5), 4.49-4.52 (m, 1H), 4.33 (d, 1H, J = 15.5), 3.57 (br s, 1H), 3.23 (br s, 1H), 2.90 and 3.05 (2 s, 3H), 2.48-2.62 (m, 1H), 1.57-1.72 (m, 1H), 1.08-1.30 (m, 3H).
489				6	off-white solid	503.96	1.40 min Method A	504.41	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (d, 4H, J = 8.4), 7.52 (d, 2H, J = 8.8), 7.38 (d, 2H, J = 8.4), 6.49 (br s, 1H), 6.31 (br s, 1H), 5.72 (tm, 1H, J <sub>HF</sub> = 57), 5.19 (br s, 1H), 4.59 (d, 1H, J = 15.0), 4.48-4.52 (m, 1H), 4.31 (d, 1H, J = 15.6), 3.62-3.68 (m, 2H), 3.53-3.59 (m, 2H), 3.39 (s, 3H), 2.48-2.62 (m, 1H), 1.57-1.72 (m, 1H).
490				6	white solid	485.97	1.27 min Method B	486.14	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 300MHz) δ 7.69-7.74 (m, 4H), 7.51 (dd, 2H, J = 1.8, 8.4), 7.38 (d, 2H, J = 8.1), 6.48 (br s, 1H), 6.32 (br s, 1H), 5.19 (br s, 1H), 4.54-4.62 (m, 2H), 4.30-4.41 (m, 2H), 4.13-4.19 (m, 1H), 3.61-3.67 (m, 2H), 3.53-3.59 (m, 2H), 3.39 (s, 3H), 2.48-2.62 (m, 1H), 1.57-1.72 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
491				14	off-white solid	484.91	1.54 min Method B	485.12	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.06 (d, 2H, J = 8.4), 7.74 (d, 2H, J = 8.8), 7.53 (d, 2H, J = 8.8), 7.48 (d, 2H, J = 8.4), 6.34 (br s, 1H), 5.72 (tm, 1H, J <sub>HF</sub> = 57), 5.20 (br s, 1H), 4.65 (d, 1H, J = 15.4), 4.51-4.56 (m, 1H), 4.33 (d, 1H, J = 15.8), 2.48-2.65 (m, 1H), 2.47 (s, 3H), 1.57-1.65 (m, 1H).
492				14	off-white solid	466.92	1.65 min Method A	467.19	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.04 (d, 2H, J = 8.0), 7.74 (dd, 2H, J = 1.6, 8.4), 7.48-7.51 (m, 4H), 6.32 (br s, 1H), 5.17 (br s, 1H), 4.65 (d, 1H, J = 16.0), 4.56-4.60 (m, 1H), 4.40 (d, 1H, J = 16.0), 4.29-4.32 (m, 1H), 4.18-4.21 (m, 1H), 2.47 (s, 3H), 2.18-2.38 (m, 1H), 1.50-1.65 (m, 1H).
493				6	white solid	478.88	1.59 min Method A	479.15	<sup>1</sup> H NMR (dmso-d <sub>6</sub> , 400MHz) δ 8.88 (d, 2H, J = 8.0), 7.85 (d, 2H, J = 8.8), 7.63 (d, 2H, J = 8.8), 7.51 (d, 2H, J = 8.0), 7.42 (br s, 1H), 7.24 (br s, 1H), 4.77 (d, 2H, J = 2.0), 4.40-4.44 (m, 1H), 1.97-2.05 (m, 2H), 1.75-1.84 (m, 1H), 1.42-1.50 (m, 1H).
494				1	white solid	492.91	1.66 min Method A	493.11	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.97 (d, 2H, J = 8.4), 7.69 (dd, 2H, J = 1.6, 8.4), 7.50 (dd, 2H, J = 2.0, 8.4), 7.40 (d, 2H, J = 8.4), 6.22 (br s, 1H), 5.19 (br s, 1H), 4.58 (d, 1H, J = 16.0), 4.43 (d, 1H, J = 15.6), 4.31-4.35 (m, 1H), 3.91 (s, 3H), 2.08-2.20 (m, 1H), 1.88-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.38-1.47 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
495				6	off-white solid	519.97	1.48 min Method B	520.21	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.70 (d, 2H, J = 8.4), 7.50 (d, 2H, J = 8.8), 7.37 (d, 2H, J = 8.4), 7.33 (d, 2H, J = 8.0), 6.20 (br s, 1H), 5.26 (br s, 1H), 4.45-4.52 (m, 2H), 4.33 (t, 1H, J = 7.2), 3.61-3.70 (m, 1H), 3.22-3.27 (m, 1H), 2.90 and 3.06 (2 s, 3H), 2.05-2.18 (m, 1H), 1.90-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.38-1.47 (m, 1H), 1.11-1.28 (m, 3H).
496				6	off-white solid	569.01	1.21 min Method B	569.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.57 (d, 1H, J = 4.8), 7.82 (d, 2H, J = 8.4), 7.68-7.72 (m, 4H), 7.50 (d, 2H, J = 8.8), 7.42 (d, 2H, J = 8.4), 7.35 (d, 1H, J = 8.0), 7.24 (br s, 1H), 6.24 (br s, 1H), 5.26 (br s, 1H), 4.76 (d, 2H, J = 4.8), 4.58 (d, 1H, J = 15.6), 4.41 (d, 1H, J = 15.6), 4.31-4.35 (m, 1H), 2.10-2.19 (m, 1H), 1.90-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.38-1.47 (m, 1H).
497				6	white solid	535.97	1.53 min Method A	536.16	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.72 (d, 2H, J = 8.0), 7.69 (d, 2H, J = 8.4), 7.50 (d, 2H, J = 8.4), 7.40 (d, 2H, J = 8.4), 6.48 (br s, 1H), 6.22 (br s, 1H), 5.20 (br s, 1H), 4.57 (d, 1H, J = 15.6), 4.41 (d, 1H, J = 15.6), 4.30-4.34 (m, 1H), 3.63-3.67 (m, 2H), 3.55-3.57 (m, 2H), 3.39 (s, 3H), 2.10-2.20 (m, 1H), 1.88-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.38-1.47 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
498				6	white solid	505.95	1.55 min Method A	506.10	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.69 (dd, 2H, J = 1.6, 8.4), 7.50 (d, 2H, J = 8.4), 7.39 (d, 2H, J = 8.0), 7.25-7.28 (m, 2H), 6.22 (br s, 1H), 6.03 (br s, 1H), 5.19 (br s, 1H), 4.57 (d, 1H, J = 15.6), 4.41 (d, 1H, J = 15.6), 4.30-4.33 (m, 1H), 3.48-3.53 (m, 2H), 2.10-2.20 (m, 1H), 1.88-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.38-1.47 (m, 1H), 1.25 (t, 3H, J = 7.2).
499				14	Beige solid	516.93	1.65 min Method B	517.15	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 8.06 (dd, 2H, J = 1.6, 8.4), 7.71 (dd, 2H, J = 2.0, 8.4), 7.49-7.52 (m, 4H), 6.22 (br s, 1H), 5.20 (br s, 1H), 4.63 (d, 1H, J = 15.6), 4.44 (d, 1H, J = 15.6), 4.33-4.37 (m, 1H), 2.47 (s, 3H), 2.10-2.20 (m, 1H), 1.91-2.03 (m, 1H), 1.73-1.86 (m, 1H), 1.40-1.51 (m, 1H).
500				1	off-white foam	494.94	1.74 min Method B	475.27 (M-HF+H <sup>+</sup> )	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 500MHz) δ 7.65 (dd, 2H, J = 2.0, 8.5), 7.46 (ddd, 2H, J = 2.0, 2.5, 7.5), 7.25-7.31 (m, 4H), 6.18 (br s, 1H), 5.24 (br s, 1H), 4.47 (d, 1H, J = 15.5), 4.41 (d, 1H, J = 15.0), 4.35 (t, 1H, J = 7.7), 2.08-2.20 (m, 1H), 1.88-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.68 (s, 3H), 1.63 (s, 3H), 1.38-1.47 (m, 1H).

Ex. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Reaction Scheme	Appearance	Calc. MW	Ret. Time/ Method	M+H <sup>+</sup>	NMR Data
501				12	off-white solid	492.95	2.20 min Method A	475.96 (- H <sub>2</sub> O) 492.01 (neg ion)	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400MHz) δ 7.65 (d, 2H, J = 8.8), 7.46 (d, 2H, J = 8.4), 7.40 (d, 2H, J = 8.4), 7.25-7.27 (m, 2H), 6.19 (br s, 1H), 5.23 (br s, 1H), 4.46 (d, 1H, J = 15.4), 4.41 (d, 1H, J = 15.4), 4.34 (t, 1H, J = 7.6), 2.08-2.20 (m, 1H), 1.88-2.03 (m, 1H), 1.69-1.82 (m, 1H), 1.56 (s, 3H), 1.54 (s, 3H), 1.38-1.47 (m, 1H).

Method A = 4.6 X 33 mm ODS-A C-18 column, 5mL/min, 10:90:0.1 (MeOH/H<sub>2</sub>O/TFA) to 90:10:0.1 (MeOH/H<sub>2</sub>O/TFA), 2min gradient

Method B = 3 X 50 mm ODS-A C-18 column, 5mL/min, 10:90:0.1 (MeOH/H<sub>2</sub>O/TFA) to 90:10:0.1 (MeOH/H<sub>2</sub>O/TFA), 2min gradient

Method C = 3 X 50 mm ODS-A C-18 column, 5mL/min, 10:90:0.1 (MeOH/H<sub>2</sub>O/TFA) to 90:10:0.1 (MeOH/H<sub>2</sub>O/TFA), 3min gradient

Method D = 4.6 X 50 mm Phenomenex Luna C-18 S5 column, 5mL/min, 0-100% MeOH/H<sub>2</sub>O, 0.1%TFA, 2min gradient

Method E = 4.6 X 50 mm Xterra C18 S5 column, 5mL/min, 0-100% MeOH/H<sub>2</sub>O, 0.1 % TFA, 2min gradient


Method F = 4.6 X 50 mm Phenomenex Luna C-18 S5 column, 5mL/min, 10:90:0.1 (MeOH/H<sub>2</sub>O/TFA) to 90:10:0.1 (MeOH/H<sub>2</sub>O/TFA), 2min gradient

Method G = 3.0 X 50 mm Xterra C18 S7 column, 5mL/min, 10:90:0.1 (MeOH/H<sub>2</sub>O/TFA) to 90:10:0.1 (MeOH/H<sub>2</sub>O/TFA), 2min gradient

- 246 -

What is claimed is:

1. A compound of formula I; or an optical isomer thereof



wherein:

R<sup>1</sup> is selected from the group consisting of

(a) a straight or branched-chain C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl optionally substituted with substituents selected from the group consisting of hydroxy, C<sub>2-7</sub> cycloalkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkylthio, and halogen;

(b) C<sub>2-7</sub> cycloalkyl optionally substituted with hydroxy or halogen; is hydrogen or R<sup>1</sup> and R taken together is C<sub>2-6</sub> alkylene;

R<sup>2</sup> is selected from the group consisting of


(a) a straight or branched-chain C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub> alkoxy, and NR<sup>4</sup>R<sup>5</sup>;

(b) C<sub>2-7</sub> cycloalkylmethyl optionally substituted with substituents selected from the group consisting of amino, (C<sub>1-6</sub> alkyl)NH<sub>2</sub>, di(C<sub>1-6</sub> alkyl)N<sup>+</sup>, C<sub>1-4</sub> alkylC(=O)NH<sub>2</sub>, and C<sub>1-4</sub> alkylOC(=O)NH<sub>2</sub>;

(c) a straight or branched-chain C<sub>1-6</sub> alkyl-C(=O)-A;

(d) -B-naphthyl;

(e)



25

D and E are each independently a direct bond, a straight or branched-chain C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, or C<sub>2-7</sub> cycloalkyl;

- 247 -



- 248 -

Z is selected from the group consisting of hydrogen,  $C_{1-4}alkyl$ ,  $C_{1-4}alkoxy$ , halogen, cyano, hydroxy,  $-OCHF_2$ ,  $-OCF_3$ ,  $-CF_3$ , and  $-CHF_2$ ;

5 X and Y are each independently selected from the group consisting of hydrogen, hydroxy, halogen, (halogen) $_2C$ , (halogen) $_2CH$ ,  $C_{1-4}alkylS$ -,

$C_{1-4}alkylS(O)$ -,  $C_{1-4}alkylSO_2$ -, nitro,  $F_3S$ -, and cyano;

$-OR^4$ ;

$-NR^4R^5$ ;

$-NR^4C(=O)R^4$ ;

$-NR^4C(=O)OR^4$ ;

$-NHSO_2C_{1-4}alkyl$ ;

$-N(SO_2C_{1-4}alkyl)_2$ ;

$-C(=O)W$  wherein W is selected from the group consisting of

hydroxy,  $C_{1-4}alkyl$ ,  $C_{1-4}alkoxy$ , phenoxy, and  $-NR^4R^5$ ;

$-OC(=O)C_{1-4}alkyl$ ;

15 -phenyl in which said phenyl is optionally substituted with cyano, halogen,  $C_{1-4}alkoxy$ ,  $C_{1-4}alkylS$ -,  $CH_2C(=O)$ ,  $C_{1-4}alkylS(O)$ -, or

$C_{1-4}alkylSO_2$ -, and

20 heterocyclic group, in which said heterocyclic group is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl,

oxazolyl, isoxazolyl, thiadiazolyl, and thiazolyl, wherein said

heterocyclic group is optionally substituted with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}alkyl$ ,

25 (halogen) $C_{1-4}alkyl$ , and  $CO_2C_{1-4}alkyl$ ;

(f) -B-(heterocycle), in which said heterocycle is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, oxazolyl, isoxazolyl, thiadiazolyl and thiazolyl wherein said heterocycle is optionally substituted with substituents selected from the group consisting of cyano, halogen,  $C_{1-4}alkyl$ ,  $CO_2C_{1-4}alkyl$ , amino,

- 249 -

,  $(C_{1-4}alkyl)NH$ -,  $di(C_{1-4}alkyl)N$ -, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperidin-1-yl, and 4-

$(C_{1-4}alkyl)piperazin-1-yl$ ;

(g) -B-(piperidin-4-yl), in which said piperidin-4-yl is optionally

5 substituted with substituents selected from the group consisting of a

straight or branched-chain  $C_{1-4}alkyl$ ,  $CH_2C(=O)phenyl$ , phenyl and phenylmethyl in which said  $C_{1-4}alkyl$  and said phenyl are optionally

substituted with substituents selected from the group consisting of

cyano, halogen, benzimidazol-2-yl, pyridyl and tetrahydrofuran-2-yl;

10 and  $-C(=O)W'$  wherein  $W'$  is selected from the group consisting of

$C_{1-4}alkoxy$ ,  $R^6$ , and  $-NR^4R^5$ ;

A is hydroxy,  $C_{1-4}alkoxy$  or  $NR^4R^5$ ;

B is a straight or branched-chain  $C_{1-4}alkyl$  or  $C_{2-6}alkenyl$ ;

15  $R^3$  is phenyl or pyridyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy,  $C_{1-4}alkoxy$ ,  $C_{1-4}alkyl$ ,

(halogen) $_2C$ -, (halogen) $_2CH$ -, and halogen $CH_2$ -,

$R^4$  and  $R^5$  each are independently hydrogen, a straight or branched-chain  $C_{1-6}$

alkyl,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_3cycloalkyl$ ,  $C_3cycloalkylmethyl$ ,

$C_{1-4}alkoxy$ , phenyl, benzyl, pyridyl, piperidin-4-yl, indan-1-yl, indan-2-yl,

20 tetrahydrofuran-3-yl, or pyrrolidin-3-yl; in which each is optionally

substituted with substituents selected from the group consisting of

hydroxy, cyano, halogen, (halogen) $_2C$ -, (halogen) $_2CH$ -, halogen $CH_2$ -,

hydroxymethyl, benzyloxymethyl, phenyl, pyridyl,  $C_{1-4}alkyl$ ,  $C_{1-4}alkoxy$ ,

(halogen) $_2C-O$ -, (halogen) $_2CH-O$ -,  $C_{1-4}alkylthio$ , amino,  $(C_{1-4}alkyl)NH$ -,

25  $di(C_{1-4}alkyl)N$ -, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl,

piperidin-1-yl, piperazin-1-yl, 4-( $C_{1-4}alkyl$ )piperazin-1-yl, 4-

phenylpiperazin-1-yl, 4-benzylpiperazin-1-yl, 4-pyridylpiperazin-1-yl,

$CO_2H$ ,  $CO_2C_{1-4}alkyl$ ,  $C(=O)NH(C_{1-4}alkyl)$ , and  $C(=O)N(C_{1-4}alkyl)_2$ ;

$R^4$  and  $R^5$  taken together may be morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-

30 1-yl, 1,2,3,4-tetrahydroisquinolin-2-yl, decalhydroquinolin-1-yl,

piperidin-1-yl, piperazin-1-yl, [1,4]-oxazepan-4-yl, azetidin-1-yl, 2,3-

- 250 -

- 5 dihydro-1*H*-isoindol-2-yl, or 2,3-dihydro-1*H*-indol-1-yl, in which each is optionally substituted with substituents selected from the group consisting of hydroxy, cyano, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-, phenyl, pyridyl, benzyl, C<sub>1-4</sub>alkyl, C<sub>2-7</sub>cycloalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylthio, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1-4</sub>alkyl, C(=O)NHC<sub>1-4</sub>alkyl, and C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

- 10 R<sup>4</sup> is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-8</sub> alkenyl, benzyl, or phenyl in which each is optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, (C<sub>1-4</sub>alkyl)(phenyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl;

R<sup>5</sup> is hydrogen, a straight or branched-chain C<sub>1-4</sub>alkyl;


- 15 R<sup>6</sup> is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-7</sub>cycloalkyl, phenyl, pyridyl, or furanyl; in which each is optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl;

- 20 R<sup>7</sup> is a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-8</sub> alkenyl, benzyl, phenyl, oxazolyl or pyridyl; in which each is optionally substituted with substituents selected from the group consisting of halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl;

- 25 or a non-toxic pharmaceutically acceptable salt thereof.

- 251 -

2. The compound of Claim 1 having the formula



wherein:

- 5 R<sup>1</sup> is selected from the group consisting of  
 (a) a straight or branched-chain C<sub>1-8</sub>alkyl or C<sub>2-8</sub>alkenyl optionally substituted with substituents selected from the group consisting of hydroxy, C<sub>2-7</sub>cycloalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylthio, and halogen;  
 (b) C<sub>2-7</sub>cycloalkyl optionally substituted with hydroxy or halogen;  
 10 R<sup>2</sup> is selected from the group consisting of  
 (a) a straight or branched-chain C<sub>1-4</sub>alkyl or C<sub>2-8</sub>alkenyl optionally substituted with substituents selected from the group consisting of halogen, C<sub>1-4</sub>alkoxy, and NR<sup>5</sup>;


(b) C<sub>2-7</sub>cycloalkylmethyl optionally substituted with substituents selected from the group consisting of amino, (C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, C<sub>1-4</sub>alkyl(C(=O)NH-, and C<sub>1-4</sub>alkylOC(=O)NH-;

15

(c) a straight or branched-chain C<sub>1-4</sub>alkyl-C(=O)-A<sub>1</sub>;

(d) -B-naphthyl;

(e)



20

D and E are each independently a direct bond, a straight or branched-chain C<sub>1-4</sub>alkyl, C<sub>2-8</sub> alkenyl, or C<sub>2-7</sub>cycloalkyl;

25 Z is selected from the group consisting of hydrogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, halogen, cyano, hydroxy, -OCHF<sub>2</sub>, -OCF<sub>3</sub>, -CF<sub>3</sub>, and -CHF<sub>2</sub>;

- 252 -

X and Y are each independently selected from the group consisting of hydrogen, hydroxy, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, C<sub>1-4</sub>alkylS-, C<sub>1-4</sub>alkylS(O)-, C<sub>1-4</sub>alkylSO<sub>2</sub>-, nitro, F, S<sub>2</sub>, and cyano;

-OR<sup>5</sup>;

5 -NR<sup>4</sup>R<sup>5</sup>;

-NR<sup>2</sup>C(=O)R<sup>4</sup>;

-NR<sup>2</sup>C(=O)OR<sup>4</sup>;

-NHSO<sub>2</sub>C<sub>1-4</sub>alkyl;

-N(SO<sub>2</sub>C<sub>1-4</sub>alkyl)<sub>2</sub>;

10 -C(=O)W wherein W is selected from the group consisting of

hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, phenoxy, and -NR<sup>4</sup>R<sup>5</sup>;

-OC(=O)C<sub>1-4</sub>alkyl;

-phenyl in which said phenyl is optionally substituted with cyano,

halogen, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkylS-, CH<sub>2</sub>C(=O), C<sub>1-4</sub>alkylS(O)-, or

15 C<sub>1-4</sub>alkylSO<sub>2</sub>-; and

heterocyclic group, in which said heterocyclic group is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl,

oxazolyl, isoxazolyl, thiadiazolyl, and thiazolyl, wherein said

20 heterocyclic group is optionally substituted with substituents selected from the group consisting of cyano, halogen, C<sub>1-4</sub>alkyl,

(halogen)C<sub>1-4</sub>alkyl, and CO<sub>2</sub>C<sub>1-4</sub>alkyl;

(f) -B-(heterocycle), in which said heterocycle is selected from the group

25 consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, oxazolyl,

isoxazolyl, thiadiazolyl and thiazolyl wherein said heterocycle is

optionally substituted with substituents selected from the group

consisting of cyano, halogen, C<sub>1-4</sub>alkyl, CO<sub>2</sub>C<sub>1-4</sub>alkyl, amino,

(C<sub>1-4</sub>alkyl)NH-, di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl,

30 pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-

(C<sub>1-4</sub>alkyl)piperazin-1-yl;

- 253 -

(g) -B-(piperidin-4-yl), in which said piperidin-4-yl is optionally

substituted with substituents selected from the group consisting of a

straight or branched-chain C<sub>1-4</sub>alkyl, CH<sub>2</sub>C(=O)phenyl, phenyl and

phenylmethyl in which said C<sub>1-4</sub>alkyl and said phenyl are optionally

substituted with substituents selected from the group consisting of

cyano, halogen, benzimidazol-2-yl, pyridyl and tetrahydrofuran-2-yl;

and -C(=O)W' wherein W' is selected from the group consisting of

5 C<sub>1-4</sub>alkoxy, R<sup>6</sup>, and -NR<sup>4</sup>R<sup>5</sup>;

10 A is hydroxy, C<sub>1-4</sub>alkoxy or NR<sup>4</sup>R<sup>5</sup>;

B is a straight or branched-chain C<sub>1-4</sub>alkyl or C<sub>3-6</sub>alkenyl;

15 R<sup>1</sup> is phenyl or pyridyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkyl,

(halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, and halogenCH<sub>2</sub>-;

R<sup>4</sup> and R<sup>5</sup> each are independently hydrogen, a straight or branched-chain C<sub>1-4</sub>

15 alkyl, C<sub>3-6</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>2-3</sub>cycloalkyl, C<sub>2-3</sub>cycloalkylmethyl,

C<sub>1-4</sub>alkoxy, phenyl, benzyl, pyridyl, piperidin-4-yl, indan-1-yl, indan-2-yl,

tetrahydrofuran-3-yl, or pyrrolidin-3-yl; in which each is optionally

substituted with substituents selected from the group consisting of

hydroxy, cyano, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-,

hydroxymethyl, benzyloxyethyl, phenyl, pyridyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy,

20 (halogen)<sub>2</sub>C-O-, (halogen)<sub>2</sub>CH-O-, C<sub>1-4</sub>alkylthio, amino, (C<sub>1-4</sub>alkyl)NH-,

di(C<sub>1-4</sub>alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl,

piperidin-1-yl, piperazin-1-yl, 4-(C<sub>1-4</sub>alkyl)piperazin-1-yl, 4-

phenylpiperazin-1-yl, 4-benzylpiperazin-1-yl, 4-pyridylpiperazin-1-yl,

25 CO<sub>2</sub>H, CO<sub>2</sub>C<sub>1-4</sub>alkyl, C(=O)NHC<sub>1-4</sub>alkyl, and C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

R<sup>4</sup> and R<sup>5</sup> taken together may be morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, 1,2,3,4-tetrahydroisquinolin-2-yl, decahydroquinolin-1-yl,

piperidin-1-yl, piperazin-1-yl, [1,4]-oxazepan-4-yl, azetidin-1-yl, 2,3-

30 dithydro-1H-isindol-2-yl, or 2,3-dihydro-1H-indol-1-yl; in which each is

optionally substituted with substituents selected from the group consisting

of hydroxy, cyano, halogen, (halogen)<sub>2</sub>C-, (halogen)<sub>2</sub>CH-, halogenCH<sub>2</sub>-,

- 254 -


- phenyl, pyridyl, benzyl,  $C_{1-6}$ alkyl,  $C_{3-7}$  cycloalkyl,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio, amino,  $(C_{1-6}$ alkyl)NH-, di( $C_{1-6}$ alkyl)N-,  $CO_2H$ ,  $CO_2C_{1-6}$ alkyl,  $C(=O)NHC_{1-6}$ alkyl, and  $C(=O)N(C_{1-6}$ alkyl);
- $R^6$  is a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-6}$  alkenyl, benzyl, or phenyl in which each is optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy, amino,  $(C_{1-6}$ alkyl)NH-, di( $C_{1-6}$ alkyl)N-,  $(C_{1-6}$ alkyl)(phenyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4- $(C_{1-6}$ alkyl)piperazin-1-yl;
- 10  $R^7$  is hydrogen, a straight or branched-chain  $C_{1-6}$ alkyl;
- $R^8$  is a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-7}$  cycloalkyl, phenyl, pyridyl, or furanyl; in which each is optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $(C_{1-6}$ alkyl)NH-, di( $C_{1-6}$ alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4- $(C_{1-6}$ alkyl)piperazin-1-yl;
- $R^9$  is a straight or branched-chain  $C_{1-6}$ alkyl,  $C_{3-6}$  alkenyl, benzyl, phenyl, oxazolyl or pyridyl; in which each is optionally substituted with substituents selected from the group consisting of halogen, (halogen) $C_{1-6}$ alkoxy,  $CH_2$ -, halogen $CH_2$ -,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy, amino,  $(C_{1-6}$ alkyl)NH-, di( $C_{1-6}$ alkyl)N-, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4- $(C_{1-6}$ alkyl)piperazin-1-yl; or a non-toxic pharmaceutically acceptable salt thereof.
- 25 3. The compound of Claim 2 in which  $R^1$  is a straight or branched-chain  $C_{1-6}$ alkyl or  $C_{3-6}$ alkenyl optionally substituted with substituents selected from the group consisting of hydroxy,  $C_{3-7}$  cycloalkyl,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkylthio, and halogen.
- 30 4. The compound of Claim 2 in which  $R^1$  is  $C_{3-7}$  cycloalkyl optionally substituted with hydroxy or halogen.

- 255 -

5. The compound of Claim 3 in which  $R^1$  is a straight or branched-chain  $C_{1-6}$ alkyl optionally substituted with  $C_{3-7}$  cycloalkyl.
6. The compound of Claim 3 in which  $R^1$  is a straight or branched-chain  $C_{1-6}$ alkyl optionally substituted with halogen.
7. The compound of Claim 2 in which  $R^1$  is phenyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkyl, (halogen) $C_{1-6}$ alkoxy, and halogen $CH_2$ -. 10
8. The compound of Claim 2 in which  $R^1$  is pyridyl optionally substituted with substituents selected from the group consisting of halogen, hydroxy,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkyl, (halogen) $C_{1-6}$ alkoxy, and halogen $CH_2$ -. 15
9. The compound of Claim 7 in which  $R^3$  is phenyl optionally substituted with halogen.
10. The compound of Claim 2 in which  $R^2$  is a straight or branched-chain  $C_{1-6}$ alkyl or  $C_{3-6}$ alkenyl optionally substituted with substituents selected from the group consisting of halogen,  $C_{1-6}$ alkoxy, and  $NR^4R^5$ .
11. The compound of Claim 2 in which  $R^2$  is  $C_{3-7}$  cycloalkylmethyl optionally substituted with substituents selected from the group consisting of amino,  $(C_{1-6}$ alkyl)NH-, di( $C_{1-6}$ alkyl)N-,  $C_{1-6}$ alkyl $C(=O)NH$ -, and  $C_{1-6}$ alkyl $OC(=O)NH$ -. 25
12. The compound of Claim 2 in which  $R^2$  is a straight or branched-chain  $C_{1-6}$ alkyl- $C(=O)$ -A. 30
13. The compound of Claim 2 in which  $R^2$  is -B-naphthyl.

- 256 -

14. The compound of Claim 2 in which R<sup>2</sup> is



- 5
15. The compound of Claim 2 in which R<sup>2</sup> is B-(heterocycle), in which said heterocycle is selected from the group consisting of furanyl, thiofuranyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, oxadiazolyl, oxazolyl, isoxazolyl, thiadiazolyl and thiazolyl wherein said heterocycle is optionally substituted with substituents selected from the group consisting of cyano, halogen, C<sub>1-6</sub>alkyl, CO<sub>2</sub>C<sub>1-6</sub>alkyl, amino, (C<sub>1-6</sub>alkyl)NH<sub>2</sub>, di(C<sub>1-6</sub>alkyl)N<sub>2</sub>, morpholin-4-yl, thiomorpholin-4-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and 4-(C<sub>1-6</sub>alkyl)piperazin-1-yl.
- 10

- 15
16. The compound of Claim 2 in which R<sup>2</sup> is -B-(piperidin-4-yl), in which said piperidin-4-yl is optionally substituted with substituents selected from the group consisting of a straight or branched-chain C<sub>1-6</sub>alkyl, CH<sub>2</sub>C(=O)phenyl, phenyl or phenylmethyl in which said C<sub>1-6</sub>alkyl and said phenyl are optionally substituted with substituents selected from a group consisting of cyano, halogen, benzimidazol-2-yl, pyridyl and tetrahydrofuran-2-yl; and -C(=O)W' wherein W' is selected from the group consisting of C<sub>1-6</sub>alkoxy, R<sup>9</sup>, and -NR<sup>9</sup>R<sup>9</sup>.
- 20

- 25
17. The compound of Claim 14 in which B is straight-chain C<sub>1-6</sub>alkyl.
18. The compound of Claim 17 wherein Z is hydrogen.

- 257 -

19. The compound of Claim 17 wherein X is C(=O)W, E is a direct bond and Y is hydrogen.

20. The compound of Claim 17 wherein X is -NR<sup>4</sup>R<sup>5</sup>, E is a direct bond and Y is hydrogen.

21. The compound of Claim 17 wherein X is -OR<sup>6</sup>, E is a direct bond and Y is hydrogen.

22. The compound of Claim 17 wherein X is -NR<sup>3</sup>C(=O)R<sup>4</sup>, E is a direct bond and Y is hydrogen.

23. A pharmaceutical composition for the treatment of disorders responsive to the inhibition of  $\beta$ -amyloid peptide production comprising a therapeutically effective amount of a compound of claim 1 in association with a pharmaceutically acceptable carrier or diluent.

24. A method for the treatment of disorders responsive to the inhibition of  $\beta$ -amyloid peptide production in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound of claim 1.

25. A method of claim 24 wherein said disorder is Alzheimer's Disease and Down's Syndrome.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US02/40605

## A. CLASSIFICATION OF SUBJECT MATTER

IPC Class. : C07C 237/04, 237/14; A61K 31/18; A61P 25/28  
US Cl. : 544/80, 84, 88; 544/106; 544/310; 546/184; 546/131, 134; 514/618, 222.2, 315  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELD(S) SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 544/310

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE, BAST

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98/03166 A1 (MONSANTO COMPANY) 29 January 1998 (29.01.1998), see entire document.	1-25

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:	
*A* documents defining the general state of the art which is not considered to be of particular relevance	*-*
*B* earlier application or patent published on or after the international filing date	*X*
*L* documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim(s) or other special reason (as specified)	*Y*
*O* documents referring to an oral disclosure, use, exhibition or other means	
*P* documents published prior to the international filing date but later than the priority date claimed	*Z*

Date of the actual completion of the international search

31 March 2003 (31.03.2003)

Date of mailing of the international search report  
17 APR 2003

Name and mailing address of the ISA/US  
Combustion of Patents and Trademarks  
Box PCT  
Washington, D.C. 20531

Author/Officer  
William D. Roberts for  
Combustion of Patents and Trademarks

Facsimile No. (703)308-3230

Telephone No. (703)308-1235

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☒ OTHER: Side text

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**